

**Using CA ViaR Models with Implied Volatility
for Value at Risk Estimation**

Jooyoung Jeon
School of Enterprise and the Environment, University of Oxford

and

James W. Taylor
Saïd Business School, University of Oxford

Journal of Forecasting, 2013, Vol. 32, pp. 62-74.

Address for Correspondence:

Jooyoung Jeon
Smith School of Enterprise & the Environment
University of Oxford
Hayes House, 75 George St,
Oxford, OX1 2BQ
joo.jeon@smithschool.ox.ac.uk

Using CAViaR Models with Implied Volatility for Value at Risk Estimation

Abstract

This paper proposes VaR estimation methods that are a synthesis of conditional autoregressive value at risk (CAViaR) time series models and implied volatility. The appeal of this proposal is that it merges information from the historical time series and the different information supplied by the market's expectation of risk. Forecast combining methods, with weights estimated using quantile regression, are considered. We also investigate plugging implied volatility into the CAViaR models, a procedure that has not been considered in the VaR area so far. Results for daily index returns indicate that the newly proposed methods are comparable or superior to individual methods, such as the standard CAViaR models and quantiles constructed from implied volatility and the empirical distribution of standardised residual. We find that the implied volatility has more explanatory power as the focus moves further out into the left tail of the conditional distribution of S&P500 daily returns.

JEL classification: C22, C53, G17

Key words: Value at Risk; CAViaR; Implied Volatility; Quantile Regression; Combining.

1. Introduction

Value at risk (VaR) has become a standard tool for measuring market risk. It involves the estimation of the maximum potential loss of the market value of an asset or a portfolio over a certain time horizon at a given confidence level, which is typically chosen to be 1% or 5%. Thus, estimating the VaR involves forecasting tail quantiles of the conditional distribution of returns. The accurate assessment of the exposure to market risk of a financial institution is of great importance for internal risk control. Despite its conceptual simplicity and popularity as an industrial standard, no consensus has been reached as to the best method for estimating VaR.

The conditional autoregressive value at risk (CAViaR) models of Engle and Manganelli (2004) provide an appealing approach to VaR estimation. These models avoid distributional assumptions by modelling the quantile directly using quantile regression. They are autoregressive in structure, which is intuitively attractive, as series of financial returns tend to exhibit volatility clustering. A variety of alternative time series modelling approaches have been presented, including the use of GARCH volatility models, extreme value theory and exponentially weighted quantile regression (see Manganelli and Engle, 2004; Kuester *et al.*, 2006; Taylor, 2008b). Empirical evidence has shown that CAViaR models are competitive with other VaR models (Bao *et al.*, 2006; Yu *et al.*, 2010). An approach that contrasts with these time series methods is to base VaR estimation on the implied volatility, which is the expectation of volatility implied by the options market (see Chong, 2004; Christoffersen and Mazzotta, 2005; Giot, 2005). This approach constructs quantile forecasts using the implied volatility and a distributional assumption.

In this paper, we take the view that there is valuable and different information provided by both CAViaR statistical time series models and the methods based on implied volatility. This leads us to consider approaches for combining the two VaR methods. Forecast combining is a well-established procedure for synthesizing the different information

supplied by two or more separate forecasting methods in order to improve forecasting accuracy (Bunn, 1989). However, the focus in the literature has largely been on the combination of point forecasts. Although a number of studies have investigated the combination of volatility forecasts, there are very few papers on the combination of quantile forecasts. This is perhaps a little surprising, as it is now more than twenty years since Granger (1989) and Granger *et al.* (1989) originally proposed combining quantile forecasts. In this paper, we consider similar combining methods to those proposed by Granger *et al.* However, in contrast to their application to monthly economic time series, our focus is VaR estimation for daily financial returns data, and we use different individual quantile forecasting methods. In addition to the combining methods, we also evaluate the worth of including, in a CAViaR model, an additional regressor that is a quantile predictor based on implied volatility. We are not aware of any other studies that have considered the synthesis of CAViaR and implied volatility for VaR estimation.

In Section 2, we briefly review the literature on VaR estimation. Section 3 discusses the use of implied volatility for the prediction of volatility and VaR. In Section 4, we describe our approaches to combining VaR estimates. Section 5 presents CAViaR models that include, as an extra regressor, a quantile estimator based on implied volatility. Section 6 is an empirical study in which we evaluate the proposed approaches to combining VaR forecasts for daily stock index returns. The final section summarises the paper and provides concluding comments.

2. Value at Risk

Manganelli and Engle (2004) classify the existing VaR models into three categories: *parametric*, *semiparametric* and *nonparametric*. *Parametric* approaches involve a parameterization of the behaviour of prices, with conditional quantiles estimated using a conditional volatility forecast and an assumption for the shape of the distribution. An

example is a GARCH volatility model with a Student- t distribution or perhaps an asymmetric t distribution (see, for example, Mittnik and Paoletta, 2000). A notable benefit of a parametric method is the complete formation of the conditional returns distribution. A significant pitfall of a parametric approach is that the specification of the variance equation and the choice of distribution may be wrong.

The most widely used *nonparametric* method is historical simulation. With this method, the VaR is estimated as the quantile of the empirical distribution of historical returns from a moving window of the most recent periods. The advantage of historical simulation is that it requires no distributional assumption and that it is easy to compute. However, the VaR estimation can be poor and slow to converge to the actual VaR, especially for the extreme quantiles. Another difficulty, recognised by Boudoukh *et al.* (1998), is in the choice of the number of observations to include in the moving window. A moving window that is too small leads to large sampling errors, while too many observations in the moving window results in sluggish adaptation to the dynamic changes in the true distribution. Boudoukh *et al.* (1998), Mittnik and Paoletta (2000) and Taylor (2008b) attempt to overcome this issue through their exponentially weighted approaches to VaR estimation.

The *semiparametric* VaR category includes applications of extreme value analysis and methods based on quantile regression, such as the CAViaR models introduced by Engle and Manganelli (2004). Using an autoregressive framework, CAViaR models aim to derive the evolution of the desired quantile rather than extracting the quantile from an estimate of a complete distribution or from a volatility estimate. The approach has the advantage of allowing the shape of the conditional returns distribution to be time-varying, and for the time-variation to be different for different quantiles of the distribution. The four CAViaR models introduced by Engle and Manganelli (2004) and the AR(1)-GARCH(1,1) CAViaR model proposed by Kuester *et al.* (2006) are presented in the following expressions:

Symmetric Absolute Value CAViaR:

$$Q_t(\theta) = \beta_1 + \beta_2 Q_{t-1}(\theta) + \beta_3 |y_{t-1}|$$

Asymmetric Slope CAViaR:

$$Q_t(\theta) = \beta_1 + \beta_2 Q_{t-1}(\theta) + \beta_3 |y_{t-1}| I(\theta > 0) + \beta_4 |y_{t-1}| I(\theta < 0)$$

Indirect GARCH(1,1) CAViaR:

$$Q_t(\theta) = (1 - 2I(\theta < 0.5)) \left(\beta_1 + \beta_2 Q_{t-1}(\theta)^2 + \beta_3 y_{t-1}^2 \right)^{\frac{1}{2}}$$

Indirect AR(1)-GARCH(1,1) CAViaR:

$$Q_t(\theta) = \beta_4 y_{t-1} + (1 - 2I(\theta < 0.5)) \left(\beta_1 + \beta_2 (Q_{t-1}(\theta) - \alpha y_{t-2})^2 + \beta_3 (y_{t-1} - \alpha y_{t-2})^2 \right)^{\frac{1}{2}}$$

Adaptive CAViaR:

$$Q_t(\theta) = Q_{t-1}(\theta) + \beta_1 \left\{ \theta - [1 + \exp(G[y_{t-1} - Q_{t-1}(\theta)])]^{-1} \right\}$$

where $Q_t(\theta)$ is the θ quantile conditional upon Ψ_{t-1} , the information set up to time $t-1$; I is an indicator function which returns a value of one if the argument is true, and zero otherwise; G is some sizeable positive number; and the β_i are parameters. Note that we are modelling here a residual term, y_t , defined as $y_t = r_t - E(r_t | \Psi_{t-1})$, where r_t is the return and $E(r_t | \Psi_{t-1})$ is the conditional expectation, which is often assumed to be zero or a constant. The parameters are determined by the quantile regression (Koenker and Bassett, 1978) minimisation, which is of the following form:

$$\min_{\beta} \left[\sum_{t|y_t \geq Q_t(\theta)} \theta |y_t - Q_t(\theta)| + \sum_{t|y_t < Q_t(\theta)} (1 - \theta) |y_t - Q_t(\theta)| \right] \quad (1)$$

The structures of the first three models have similarities to GARCH models. The asymmetric slope CAViaR model is designed specifically to capture the asymmetric leverage effect. This effect is the tendency for volatility to be greater following a negative return than a positive return of equal size. Both models rely on the magnitude of the error rather than the squared error as in GARCH models. The indirect GARCH(1,1) CAViaR model and the GARCH(1,1) volatility model are similar in form, but different in that the indirect GARCH CAViaR model is estimated by quantile regression rather than by maximum likelihood, which is

used for the GARCH volatility model. The indirect AR(1)-GARCH(1,1) CAViaR model allows for the conditional mean to be time-varying. The second term in the adaptive CAViaR model is a function that forces the quantile to take a lower value if y_t falls below the quantile, and a higher value otherwise. As $G \rightarrow \infty$, the function converges to the hit variable, defined as $H_t = \theta - I(y_t \leq Q_t(\theta))$. Engle and Manganelli (2004) note that the structure of the adaptive CAViaR model is such that the estimator learns nothing from the extent to which the quantile has or has not been exceeded since it considers only whether $Q_t(\theta)$ is larger than y_t or not.

3. Using Implied Volatility for Predicting Volatility and VaR

3.1. Implied Volatility

Implied volatility represents the volatility of the underlying security that is implicit in the market price of an option according to a particular model. In other words, the implied volatility is the level of volatility in the Black-Scholes formula that delivers an option price equal to the current market price. In contrast to historical volatility, which is a measure of price changes in the past, implied volatility reflects the market's expectations regarding future volatility. There has been a growing trend towards the use of implied volatility as a key variable in financial investment decisions, risk management, derivative pricing, market making, market timing and portfolio selection.

In this paper, we do not use implied volatility for volatility estimation, but instead we use it for VaR estimation. We consider the benefit of combining quantile forecasts based on implied volatility and CAViaR time series models. Before we present this idea in more detail, let us first briefly review the related literature.

3.2. Comparison and Synthesis of Volatility Forecasting Approaches

Although there have been many studies on the issue, no consensus has been reached on the usefulness of implied volatility as a predictor for future volatility in comparison with

predictions from time series models. Although many studies conclude in favour of implied volatility, others find no significant benefit in using implied volatility over predictions from a time series model.

Looking at the relatively recent literature, we find that Szakmary *et al.* (2003) conclude that implied volatility outperforms historical volatility as a predictor of the realised volatility in a large majority of 35 futures markets including equity indices, interest rates, currencies, commodities and crude oil. Pong *et al.* (2004) find historical intraday returns have significant incremental information in exchange rates, beyond the implied volatility information, for horizons up to one week, while for longer horizons, implied volatility provides more informative forecasts than historical volatility. Corredor and Santamaria (2004) report that implied volatility clearly dominates all of the time series models such as the GARCH family. Giot and Laurent (2007) find that implied volatility delivers more relevant volatility information content than historical volatility for stock indices such as the S&P100 and S&P500. However, there is also recent evidence suggesting that there is value in both implied volatility and volatility based on the historical time series of returns, and that the superiority of each depends on the financial time series considered and the time series method used. Noh and Kim (2006) conclude that both implied volatility and historical volatility using high-frequency returns can outperform each other in forecasting volatility. In their empirical test, historical volatility from high frequency returns performed better in the FTSE100 futures, which tends to be relatively close to normally distributed, while the result of implied volatility was better in the S&P500 futures, which displays excess skewness even with volatilities from high frequency returns.

If it is not clear which of two forecasts performs better, a combination can be the best option. Since the seminal work of Bates and Granger (1969), a large literature has evolved on combining forecasts for the level (or mean) of a series. Examples of popular combining methods are the simple average and the use of least squares regression to estimate the

weights in a linear combination of individual forecasts. In contrast to the literature on combining forecasts of the level, research into combining volatility forecasts is far less well developed (Poon and Granger, 2003). Of the studies that do exist, there is general support for the idea of forming a linear combination of historical volatility and implied volatility (see Kroner *et al.*, 1995; Doidge and Wei, 1998; Pong *et al.*, 2004). It is also worth noting that there is evidence in favour of combining volatility forecasts constructed from the same information set, such as two GARCH models (see, for example, Amendola and Storti, 2007).

An approach that is related to combining forecasts is to include implied volatility as an exogenous variable in the time series model, such as a GARCH model. In general, this ‘plug-in’ approach has been performed with the emphasis on testing for an improvement in in-sample fit (e.g. Day and Lewis, 1992; Blair *et al.*, 2001). However, some authors have also evaluated the resulting forecasts using post-sample data (e.g. Claessen and Mittnik, 2002; Giot, 2005; Donaldson and Kamstra, 2005).

3.3. Forecasting VaR using Implied Volatility

If implied volatility forecasts the future volatility well, as suggested by the literature review in Section 3.2, it is reasonable to think that implied volatility will be useful in estimating future quantiles of the returns distribution. As we explain in Section 3.5, by making a distributional assumption, it is straightforward to construct a VaR estimator from implied volatility. Giot (2005) predicts VaR using implied volatility with a skewed Student- t distributional assumption and concludes that VaR estimates relying exclusively on lagged implied volatility perform as well as VaR estimates based on the volatility forecasts of GARCH family models. On the other hand, some studies find that implied volatility is not an effective indicator for VaR estimation. In VaR estimation for currency exchange, Chong (2004) finds that time series models perform better than the model based on implied volatility. Christoffersen and Mazzotta (2005) assess the quality of different volatility forecasts, such as

implied volatility, historical volatility and volatilities from GARCH models. They find that implied volatility provides the most accurate volatility forecasts for all currency exchange rates and forecast horizons considered. Yet, in the application of the volatility forecasts to density estimation, they conclude that the density and interval forecasts based on implied volatility do not capture the tail behaviour of the distribution. However, the assumption of Gaussian tail shape might be a weakness in the studies of Chong (2004) and Christoffersen and Mazzotta (2005). In this paper, we make no distributional assumptions in our investigation of the usefulness of implied volatility for quantile estimation.

3.4. Combining VaR Forecasts from Time Series Models and Implied Volatility

If there is support for forecasting quantiles using implied volatility and also evidence in favour of time series methods for quantile prediction, there is some appeal in producing a quantile forecast from the combination of one quantile forecast based on implied volatility and another constructed from historical returns. There are very few studies that have looked at the combination of quantile forecasts. Granger (1989) and Granger *et al.* (1989) introduce the idea of using quantile regression to combine quantile forecasts. Granger *et al.* (1989) apply the idea to quantile forecasts produced using time series methods for two monthly economic time series. Using simulated data, Taylor and Bunn (1998) assess the usefulness of different restrictions on the parameters of the quantile regression combination. Giacomini and Komunjer (2005) describe how encompassing tests can be performed for two quantile predictors using the quantile regression combining framework. They apply their proposal to the S&P500 VaR estimates based on two time series volatility forecasting methods.

In Section 3, we have briefly reviewed comparative empirical forecasting studies that have considered implied volatility, as well as the literature on empirical studies of VaR estimation. We have also highlighted the appeal of combining volatility or quantile forecasts. Although volatility forecasts from time series models have been combined with implied

volatility, we are not aware of any studies that have directly estimated VaR using a combination of quantile predictions based on implied volatility with forecasts from a time series model. Our proposal is to do just this. We consider combinations of two attractive individual VaR approaches, namely the CAViaR models and a simple VaR estimator based on implied volatility, which we call the implied quantile (IQ) and present in Section 3.5. We describe the different combining methods in Section 4.

3.5. Implied Quantile (IQ)

We construct an implied quantile (IQ) estimator for period t as the product of implied volatility recorded in the previous period, $\sigma_{t-1}^{implied}$, and the quantile, $Q^{Emp}(\theta)$, of an empirical distribution, which we construct as the distribution of the in-sample values of y_t , defined in Section 2, standardised by implied volatility. The IQ estimator can be expressed in the following form:

$$Q_t^{IQ}(\theta) = Q^{Emp}(\theta) \sigma_{t-1}^{implied} \quad (2)$$

In basing quantile estimation on implied volatility, the IQ approach captures the market's expectation of future risk. Another advantage is that the method does not assume a particular distribution for the asset returns, and it involves no parameter estimation. We also considered the use of a Gaussian assumption, but the post-sample forecasting results were comfortably superior for the empirical distribution. It is interesting to note that this simple approach to capturing an 'implied quantile' assumes returns standardised with implied volatility are i.i.d. By contrast, the CAViaR models allow for the shape of the conditional distribution to be time-varying, as well as the volatility. The IQ method is similar to the filtered historical simulation method of Barone-Adesi *et al.* (1998), but different in that we use implied volatility instead of the variance estimated from a GARCH model.

4. Combining Implied Quantile and CAViaR

4.1. Simple Average Combining (*SimpAvg*)

The simplest and most widely used forecast combining method is to take the simple arithmetic mean of the individual forecasts. We consider the simple average of the quantile forecasts from the IQ method and one CAViaR model, as in expression (3). When presenting our empirical results, we refer to this method as ‘SimpAvg’ combining.

$$Q_t(\theta) = \frac{1}{2} Q_t^{IQ}(\theta) + \frac{1}{2} Q_t^{CAViaR}(\theta) \quad (3)$$

4.2. Unrestricted Linear Combination (*LinearComb*)

A traditional approach to combining is to compute linear combinations of forecasts. Granger *et al.* (1989) apply this combining method to quantile forecasts using quantile regression to optimise the parameters. We also use this method to combine the CAViaR and IQ forecasts. The resultant quantile forecast is of the following form:

$$Q_t(\theta) = \gamma_1 + \gamma_2 Q_t^{IQ}(\theta) + \gamma_3 Q_t^{CAViaR}(\theta) , \quad (4)$$

where the γ_i are parameters estimated using the quantile regression minimisation of expression (1). As expression (4) is linear, the minimisation can be performed efficiently using linear programming (see Koenker and Bassett, 1978). We refer to this method as ‘LinearComb’.

4.3. Weighted Averaged Combining (*WtdAvg*)

Granger (1989) suggests that if the quantile forecasts are unbiased, a weighted average combination could be used. This amounts to the linear combination model of expression (4) with no constant term and combining weights restricted to sum to one. This constrained form of the linear combination is often used for forecasting the mean of a series. In that context, Clemen (1986) advocates the use of the weighted average even if the

forecasts are biased, arguing that gains in efficiency can be made at the cost of some bias. Bunn (1989) strengthens the case for a weighted average combination by explaining that it can be more robust than an unconstrained model. With these benefits in mind, we included in our study the weighted average of the individual forecasts from the CAViaR models and the IQ. The resultant quantile forecast is of the following form:

$$Q_t(\theta) = \omega Q_t^{IQ}(\theta) + (1 - \omega) Q_t^{CAViaR}(\theta) , \quad (5)$$

where ω is a combining weight constrained to be between zero and one, which is estimated using quantile regression. As with the ‘LinearComb’ method, linear programming can be used to optimise the combining parameter. We refer to this method as ‘WtdAvg’ combining. As Taylor and Bunn (1998) point out, one benefit of the weighted average method, as in expression (5) above, is that the value of the weight, ω , indicates the relative explanatory powers of the two quantile predictors.

4.4. Weighted Average Combining Optimised using Exponential Weighting (WtdAvgExp)

This method is similar to the weighted average combination method of Section 4.3, but different in that it uses exponentially weighted quantile regression (EWQR) (see Taylor, 2008b) for the optimisation of the combining weight ω . The intuition of the EWQR method is that it gives more weight to more recent observations in the quantile regression optimisation. Boudoukh *et al.* (1998) assert that, as series of financial returns exhibit time-varying and cyclical volatility, an exponential weighting approach presents a reasonable trade-off between statistical precision and adaptiveness to recent news. The EWQR minimization has the following form:

$$\min_{\omega} \left[\sum_{t|y_t \geq Q_t(\theta)} \lambda^{T-t} \theta |y_t - Q_t(\theta)| + \sum_{t|y_t < Q_t(\theta)} \lambda^{T-t} (1 - \theta) |y_t - Q_t(\theta)| \right],$$

where $Q_t(\theta)$ is a weighted average combination of the form of expression (5). A lower value of the decay parameter, λ , implies faster exponential decay, and hence more weight is given

to the recent observations and less historical information is captured. In Section 6, we describe how we optimised the value of λ in our empirical study. In the discussion of our empirical results, we refer to this method as ‘WtdAvgExp’.

5. Plugging the Implied Quantile into CAViaR (PlugIn)

In Section 3.2, we discussed how implied volatility has been merged with GARCH models by simply plugging the implied volatility into the statistical models as an additional regressor. In this section, we present the analogous idea for quantile estimation, which involves the IQ predictor being plugged into the CAViaR models from Section 2. The resultant ‘PlugIn’ CAViaR models are presented in expressions (6) to (10).

Symmetric Absolute Value PlugIn:

$$Q_t(\theta) = \beta_1 + \beta_2 Q_{t-1}(\theta) + \beta_3 |y_{t-1}| + \beta_{IQ} Q_t^{IQ}(\theta) \quad (6)$$

Asymmetric Slope PlugIn:

$$Q_t(\theta) = \beta_1 + \beta_2 Q_{t-1}(\theta) + \beta_3 |y_{t-1}| I(y_{t-1} \geq 0) + \beta_4 |y_{t-1}| I(y_{t-1} < 0) + \beta_{IQ} Q_t^{IQ}(\theta) \quad (7)$$

Indirect GARCH(1,1) PlugIn:

$$Q_t(\theta) = (1 - 2I(\theta < 0.5)) \left(\beta_1 + \beta_2 Q_{t-1}(\theta)^2 + \beta_3 y_{t-1}^2 + \beta_{IQ} Q_t^{IQ}(\theta)^2 \right)^{\frac{1}{2}} \quad (8)$$

AR(1)-GARCH(1,1) PlugIn:

$$Q_t(\theta) = \alpha y_{t-1} + (1 - 2I(\theta < 0.5)) \left(\beta_1 + \beta_2 (Q_{t-1}(\theta) - \alpha y_{t-2})^2 + \beta_3 (y_{t-1} - \alpha y_{t-2})^2 + \beta_{IQ} Q_t^{IQ}(\theta)^2 \right)^{\frac{1}{2}} \quad (9)$$

Adaptive PlugIn:

$$Q_t(\theta) = \beta_1 + \beta_2 Q_{t-1}(\theta) + \beta_3 \left\{ [1 + \exp(G[y_{t-1} - Q_{t-1}(\theta)])]^{-1} - \theta \right\} + \beta_{IQ} Q_t^{IQ}(\theta) \quad (10)$$

These models enable an understanding of the importance of implied volatility by inspecting the coefficient, β_{IQ} , of the implied quantile predictor for different θ quantiles.

We optimise the models using the standard quantile regression minimisation of expression

(1), which is used to optimise the standard CAViaR models. We are not aware of any previous research that has considered the ‘CAViaR PlugIn’ models presented in this section.

6. Empirical Study

In this section, we compare the accuracy of the VaR estimation from the methods presented in Sections 4 and 5 with that of the standard CAViaR models and the IQ estimator. Our study used daily log returns for the S&P500 and the DAX30 stock indices, and their respective implied volatility indices, the VIX and the VDAX. The VIX is derived from S&P500 index call and put options of a wide range of strike prices that are further weighted to represent a hypothetical at-the-money option with a constant maturity of 22 trading days (30 calendar days) to expiry. The VDAX is constructed by call and put DAX index options of eight different strike prices that are further linearly interpolated to a remaining life of 45 calendar days. We divided the VIX and VDAX by $\sqrt{252}$ to derive daily implied volatility.

We opted to use 2,500 periods to estimate method parameters, and 500 periods for post-sample evaluation of day-ahead quantile estimates. An alternative to the use of a fixed estimation sample would be to use a rolling window approach that would involve repeated re-estimation of the various method parameters. Our focus on day-ahead estimation is consistent with the holding period considered for internal risk control by most financial firms. We first considered the period of 3,000 days that started on the start date of the VDAX. This period consisted of 1 January 1992 to 2 July 2003. For this set of data, the 500-day post-sample period fell on a rather highly volatile period for both stock indices. More recent periods had experienced substantially lower volatility. In view of this, we reran all of our analysis for a second sample that consisted of the most recent 3,000 days at the time of our data collection. This second sample consisted of 31 March 1995 to 29 September 2006. Figures 1 and 2 present plots of the two stock indices and their implied volatilities for a period spanning both samples. In the remainder of the paper, we refer to the earlier sample as

Period 1 and its volatile post-sample period as Post-sample 1. The later sample is referred to as Period 2, and its relatively tranquil post-sample period is referred to as Post-sample 2. We report the results for both samples, which enables us to check the consistency of our results under challenging and stable market conditions.

----- Figures 1 and 2 -----

For both indices, we subtracted from each return, r_t , the mean, μ , of the 2,500 in-sample returns. The quantile estimation methods were applied to the resulting residuals, $y_t = r_t - \mu$. The confidence level of VaR is typically chosen to be 1% or 5%. Our study evaluates forecast accuracy for the 1%, 5%, 95%, and 99% conditional quantiles. For a trader who is long in the index, the left quantile (e.g. 1% and 5%) is relevant because the trading losses occur on the left side of the returns density. However, for a trader who is short in the index, the right quantile (e.g. 95% and 99%) is more relevant because trading losses occur on the right side of the returns density.

6.1. Implementation of the VaR Estimation Methods

We implemented the following VaR estimation approaches: the five individual CAViaR models presented in Section 2; the IQ method of Section 3.5; for each of the five individual CAViaR models, the four combining methods presented in Section 4; and the IQ PlugIn CAViaR models of Section 5. In a combination, three or more methods could certainly be considered. However, we feel that a reasonable choice in this initial study is a combination of one time series method and one based on implied volatility. The post-sample quantile estimation results were extremely similar for the indirect GARCH CAViaR and indirect AR-GARCH CAViaR models, and so, for simplicity, in the remainder of the paper, we present and discuss only the results for the indirect GARCH CAViaR model. This result is consistent with that of Taylor (2008a), but contrasts with that of Kuester *et al.* (2006) who

found that the indirect AR-GARCH CAViaR model outperformed the simpler indirect GARCH CAViaR model.

To optimise the standard and IQ PlugIn CAViaR models of Sections 2 and 5, linear programming cannot be used to perform the quantile regression minimisation due to the nonlinearity of the problem. Instead, we performed the optimisation using both the approach proposed by Engle and Manganelli (2004) in their published paper and the approach that they used in the 1999 draft of the paper. The 1999 version of the paper used the differential evolutionary genetic algorithm by Storn and Price (1997), while the published version used a quasi-Newton optimisation algorithm. Our optimisation proceeded by first generating 10^5 vectors of parameters from a uniform random number generator between 0 and 1, or between -1 and 0, depending on the appropriate sign of the parameter. For each of the vectors, the quantile regression summation (QRSum) of expression (1) is evaluated. Then two approaches were used for further optimisation. In the first, the 10 vectors that produced the lowest values for the QRSum out of the 10^5 vectors of parameters were used as the best initial values in a quasi-Newton optimisation routine. In the second approach, the 200 vectors that produce the lowest values for the QRSum 10^5 vectors of parameters are used as the population in the genetic algorithm, which involved 2,000 generations and, following Engle and Manganelli, a mutation parameter of 0.8 and crossover parameter of 0.5. Out of the two further optimisation methods performed, the vector producing the lowest QRSum was chosen as the final parameter vector. In this study, all the computations were performed in Matlab 7.5.0.

To optimise the parameters of the LinearComb and WtdAvg combining methods, we expressed the quantile regression minimisation as a linear programme and applied the Nelder-Mead Simplex algorithm. We used a similar approach for the WtdAvgExp combining method after first optimising the EWQR decay parameter λ . To do this, we considered a grid of values for λ between 0.97 and 1, with a step size of 0.001. To find the optimal λ value, we performed EWQR on the in-sample data with the last 500 observations excluded. The

optimal value of λ was chosen as the value that led to the lowest QRSum calculated for the last 500 observations of the in-sample data. We performed the optimisation separately for each value of θ (i.e. for each different quantile).

6.2. In-sample Results for the WtdAvg Combining Method

In this section, we report the in-sample estimated values of the weight parameter ω in the WtdAvg combining method of expression (5) in Section 4.3. (For conciseness, we restrict attention here to the WtdAvg method, rather than consider the weights in each of the combining methods.) The value of the weight is constrained to lie between zero and one, and thus expresses the explanatory power of the IQ method in relation to the CAViaR model. As described in Section 4.3, the weight is estimated by quantile regression. Table 1 examines how the parameter changes for a range of different values of θ for the S&P500 and DAX30 series, respectively. Although the focus of our quantile forecasting empirical study in Section 6.3 is estimation of the 1%, 5%, 95% and 99% quantiles, we also consider in this section the 10% and 90% quantiles. Due to the large overlap between the in-sample data of Periods 1 and 2, the estimated weights were quite similar for the two periods. In view of this, we present only the in-sample results for the more recent set of data, Period 2.

----- Table 1 -----

Table 1 shows that, in contrast to the adaptive CAViaR model, the other CAViaR models do not have useful explanatory power when combined with the IQ method. The values of the combining weights in Table 1 for the S&P500 indicate that the explanatory power of the IQ method tends to increase, relative to that of the CAViaR models, as the focus moves from the extreme upper tail of the distribution towards the lower tail. Intuitively, it would seem that the variation in the lower tail of the distribution is better captured by the variation in the implied volatility and the empirical distribution of standardised residuals,

than by the autoregressive quantile model estimated by quantile regression. There is no such pattern evident in Table 1 for the DAX30. In fact, for the most extreme quantiles that we consider (the 1% and 99% quantiles), the IQ method has a smaller weighting in the lower tail than in the upper tail of the distribution.

A statistical test could perhaps be performed for the combining weight to test whether it is significantly different from zero or from one. However, the standard error and distribution for the test is not straightforward because the weight is constrained to fall between zero and one. Encompassing tests could be used for the LinearComb approach, but implementing such a test is not straightforward in the weighted average case due to the bounds on the value of the weight. This is presumably one reason for why it is not standard practice in the combining literature to carry out tests of the combining weight.

6.3. In-sample Results for the PlugIn Method

Similarly to Section 6.2, we also investigated the in-sample estimated values of the coefficient β_{IQ} in the PlugIn method of expressions (6) to (10) in Section 5. Using the approach of Engle and Manganelli (2004), we performed significance tests of the coefficients. In Table 2, we present estimated values of the coefficient β_{IQ} , along with the results of the significance test of the hypothesis $\beta_{IQ}=0$. Table 2 shows that, for approximately half of the cases, the coefficient β_{IQ} is significantly different from zero. This indicates that, for these values of θ , implied volatility tends to have incremental information in the PlugIn method.

----- Table 2 -----

6.4. Post-sample Forecasting Results

To assess the post-sample predictive performance of the VaR methods, we used the hit percentage and dynamic quantile (DQ) test statistic, which are two measures employed by

Engle and Manganelli (2004). The hit percentage assesses the percentage of observations falling below the VaR estimator. The ideal value is θ for estimation of the θ quantile. We examined significant difference from this ideal using a test based on the binomial distribution. The Engle and Manganelli DQ test for conditional coverage evaluates whether the dynamic sequence of the hit variable is distributed i.i.d. Bernoulli with probability θ , and is independent of the conditional quantile estimator, $Q_t(\theta)$. This test uses a regression framework to test whether the variable H_t , defined in Section 2, has zero unconditional and conditional expectations. As in the empirical studies of Engle and Manganelli (2004) and Huang *et al.* (2010), we included four lags of the variable H_t in the test's regression to deliver a DQ test statistic, which, under the null hypothesis of perfect conditional coverage, is distributed χ^2 with six degrees of freedom.

----- Tables 3 to 5 -----

We have four post-sample periods in our study. In Tables 3 and 4, for the S&P500, we present the results for Post-sample 1 and Post-sample 2, respectively. In each table, for each θ quantile, we report the hit percentage and the p-value of the DQ test statistic. Table 5 summarises the results of Tables 3 and 4 by presenting the number of occurrences of significance for each method applied to each of the two post-sample periods. Smaller values in Table 5 are better. Table 5 also presents a summary of the corresponding results for the DAX30. To help explain Table 5, let us focus on the first seven rows of values in the table. These seven rows summarise Table 3. In these rows, for each θ quantile and each method, the value shown is the number of entries in Table 3 that were significant at the 5% level for the hit percentage and for the DQ test. Let us consider the interpretation of a value of 1 for the SimpAvg method in the hit percentage column corresponding to the $\theta=99\%$ quantile for Post-Sample 1 for the S&P500. This means that, for one out of the four 99% CAViaR models

considered in Table 3, the hit percentage for the SimpAvg method was significant at the 5% level. The final seven rows of the table average the values in the rows above.

Using Table 5, let us briefly compare the results for the two different post sample periods, Post-samples 1 and 2. For the S&P500, the CAViaR method adapted better to the challenging trading conditions in Post-sample 1 than the IQ method, whereas the IQ method performed better than the CAViaR method in Post-sample 2. For the DAX30, it is not clear which individual method performed better. However, regardless of whether the data series is the S&P500 or the DAX30, or whether the post-sample period is volatile or tranquil, the results for the combining method would seem to be competitive with those of the two individual methods. Turning to the PlugIn method, the results show that it performed reasonably well in Post-sample 1, but it was less competitive for Post-sample 2.

Looking at the average of the hit percentage results in the bottom seven rows of Table 5, it is impressive to see that all the combining methods and the PlugIn method either match or outperform the individual CAViaR and IQ methods for each of the four quantiles. In the fifth column of values, we can see that the average values for the combining methods are better than those for the IQ method and the individual CAViaR methods. Of the seven methods, the LinearComb method produced the best hit percentage results, followed by the SimpAvg and WtdAvgExp combining methods.

Turning to the summary of the DQ test results in the bottom seven rows of Table 5, although there is no method that performs the best across all four quantiles, the four combining methods do not perform poorly, relative to the individual methods, for any of the four. Interestingly, the PlugIn method is not as competitive as the combining methods. In the final column of DQ test results, we see that overall the four combining methods outperform both the two individual methods and the PlugIn method. This final column shows the average values for the combining methods comparing favourably with those for the CAViaR method

and the IQ method. Of the combining methods, the results for the LinearComb and SimpAvg methods are the best.

To summarise, in terms of both the hit percentage and DQ test, the results of the combining methods were better than the two individual methods. The PlugIn method was better than the two individual methods in terms of the hit percentage, but less convincing in terms of the DQ test. Among the combining methods, our study suggests that the LinearComb and SimpAvg methods are the best. As for the reason why the LinearComb method performs better than the WtdAvg method, we would suggest that it may be because the inclusion of the intercept term in the model, and the lack of restrictions on the values of the weights, allows the individual quantile estimators to be debiased. This has certainly been the main argument in favour of the LinearComb method in the context of forecasting the level (or mean) of a time series (see Granger and Ramanathan, 1984). It is worth noting that the WtdAvgExp offered slight improvement over the simpler WtdAvg method in terms of the hit percentage, suggesting that there may be benefit in trying to optimise the combining weight by giving more weight to the more recent observations.

7. Summary and Concluding Comments

In this paper, we have introduced methods for VaR estimation that aim to synthesise the time series information supplied by CAViaR models with the information in implied volatility. The methods include the PlugIn approach that inserts, as an additional regressor in a CAViaR model, an implied quantile estimator based on implied volatility. The other methods that we considered involved linear combinations of the implied quantile and CAViaR model estimators, with parameters estimated using quantile regression. The derived values of the combining weight provided interesting evidence to suggest that the explanatory power of the implied quantile increases as one moves further into the left tail, and decreases as one goes further into the right tail with the S&P500, whereas there was no clear pattern in

the DAX30. Our post-sample forecasting results show that an unrestricted linear combination has the potential to outperform the individual methods. We also obtained encouraging results for the simple average and exponentially weighted average combining methods. Considering the relative scarcity of research on combining methods for estimating VaR, it is hoped that this study will motivate further combining research in this field.

References

- Amendola A, Storti G. 2008. A GMM procedure for combining volatility forecasts. *Computational Statistics and Data Analysis* **52** : 3047-3060.
- Bao Y, Lee T-, Saltoglu B. 2006. Evaluating predictive performance of value-at-risk models in emerging markets: A reality check. *Journal of Forecasting* **25** : 101-128.
- Barone-Adesi G, Bourgoin F, Giannopoulos K. 1998. Don't look back. *Risk* **11** : 100-104.
- Bates JM, Granger CWJ. 1969. The combination of forecasts. *Operations Research Quarterly* **20** : 451-468.
- Blair BJ, Poon SH, Taylor SJ. 2001. Forecasting S&P 100 volatility: The incremental information content of implied volatilities and high-frequency index returns. *Journal of Econometrics* **105** : 5-26.
- Boudoukh J, Richardson M, Whitelaw RF. 1998. The best of both worlds. *Risk* **11** : 64-67.
- Bunn DW. 1989. Forecasting with more than one model. *Journal of Forecasting* **8** : 161-166.
- Chong J. 2004. Value at risk from econometric models and implied from currency options. *Journal of Forecasting* **23** : 603-620.
- Christoffersen PF, Mazzotta S. 2005. The accuracy of density forecasts from foreign exchange options. *Journal of Financial Econometrics* **3** : 578-605.
- Claessen H, Mittnik S. 2002. Forecasting stock market volatility and the informational efficiency of the DAX-index options market. *The European Journal of Finance* **8** : 302-321.
- Clemen RT. 1986. Linear constraints and the efficiency of combined forecasts. *Journal of Forecasting* **5** : 31-38.

- Corredor P, Santamaría R. 2004. Forecasting volatility in the spanish option market. *Applied Financial Economics* **14** : 1-11.
- Day TE, Lewis CM. 1992. Stock market volatility and the information content of stock index options. *Journal of Econometrics* **52** : 267-287.
- Doidge C, Wei JZ. 1998. Volatility forecasting and the efficiency of the toronto 35 index options market. *Canadian Journal of Administrative Sciences* **15** : 28–38.
- Donaldson RG, Kamstra MJ. 2005. Volatility forecasts, trading volume, and the ARCH versus option-implied volatility trade-off. *The Journal of Financial Research* **28** : 519-538.
- Engle RF, Manganelli S. 2004. CAViaR: Conditional autoregressive value at risk by regression quantiles. *Journal of Business and Economic Statistics* **22** : 367-382.
- Giacomini R, Komunjer I. 2005. Evaluation and combination of conditional quantile forecasts. *Journal of Business and Economic Statistics* **23** : 416-431.
- Giot P. 2005. Implied volatility indexes and daily value at risk models. *Journal of Derivatives* **12** : 54-64.
- Giot P, Laurent S. 2007. The information content of implied volatility in light of the jump/continuous decomposition of realized volatility. *Journal of Futures Markets* **27** : 337-359.
- Granger CWJ. 1989. Invited review: Combining forecasts–20 years later. *Journal of Forecasting* **8** : 167-173.
- Granger CWJ, Ramanathan R. 1984. Improved methods of combining forecasts. *Journal of Forecasting* **3** : 197-204.
- Granger CWJ, White H, Kamstra MJ. 1989. Interval forecasting: An analysis based upon ARCH-quantile estimators. *Journal of Econometrics* **40** : 87-96.
- Huang D, Yu B, Lu Z, Fabozzi FJ, Focardi S, Fukushima M. 2010. Index-Exciting CAViaR: A New Empirical Time-Varying Risk Model. *Studies in Nonlinear Dynamics and Econometrics* **14** : art. no. 1.
- Koenker R, Bassett G. 1978. Regression quantiles. *Econometrica* **46** : 33-50.
- Kroner KF, Kneafsey KP, Claessens S. 1995. Forecasting volatility in commodity markets. *Journal of Forecasting* **14** : 77-95.

- Kuester K, Mittnik S, Paolella MS. 2006. Value-at-risk prediction: A comparison of alternative strategies. *Journal of Financial Econometrics* **4** : 53-89.
- Manganelli S, Engle RF. 2004. A Comparison of Value-at-Risk Models in Finance. Szegö G. ed. *Risk measures for the 21st century*. Wiley & Sons: Chichester, UK; 123-143.
- Mittnik S, Paolella MS. 2000. Conditional density and value-at-risk prediction of Asian currency exchange rates. *Journal of Forecasting* **19** : 313-333.
- Noh J, Kim TH. 2006. Forecasting volatility of futures market: The S&P 500 and FTSE 100 futures using high frequency returns and implied volatility. *Applied Economics* **38** : 395-413.
- Pong S, Shackleton MB, Taylor SJ, Xu X. 2004. Forecasting currency volatility: A comparison of implied volatilities and AR (FI) MA models. *Journal of Banking and Finance* **28** : 2541-2563.
- Poon SH, Granger CWJ. 2003. Forecasting volatility in financial markets: A review. *Journal of Economic Literature* **41** : 478-539.
- Storn R, Price K. 1997. Differential Evolution—A simple and efficient heuristic for global optimization over continuous spaces. *Journal of Global Optimization* **11** : 341-359.
- Szakmary A, Ors E, Kim JK, Davidson III WN. 2003. The predictive power of implied volatility: Evidence from 35 futures markets. *Journal of Banking and Finance* **27** : 2151-2175.
- Taylor JW. 2008a. Estimating value at risk and expected shortfall using expectiles. *Journal of Financial Econometrics* **6** : 231-252.
- Taylor JW. 2008b. Using exponentially weighted quantile regression to estimate value at risk and expected shortfall. *Journal of Financial Econometrics* **6** : 382-406.
- Taylor JW, Bunn DW. 1998. Combining forecast quantiles using quantile regression: Investigating the derived weights, estimator bias and imposing constraints. *Journal of Applied Statistics* **25** : 193-206.
- Taylor SJ, Xu X. 1997. The incremental volatility information in one million foreign exchange quotations. *Journal of Empirical Finance* **4** : 317-340.
- Yu PLH, Li WK, Jin S. 2010. On Some Models for Value-At-Risk. *Econometric Reviews* **29** : 622-641.

Table 1 For the in-sample data of Period 2 of the S&P500 and the DAX30, the WtdAvg combining weight on the IQ method for different values of θ . Bold indicates combining weight larger for the IQ method.

	θ	Sym Abs Value	Asym Slope	Ind Garch	AR-GARCH	Adapt
S&P500	1%	1.00	0.80	1.00	0.88	1.00
	5%	0.99	0.36	0.99	0.75	0.79
	10%	0.94	0.40	0.93	0.63	0.54
	90%	0.75	0.27	0.54	0.49	1.00
	95%	0.64	0.27	0.46	0.52	1.00
	99%	0.37	0.09	0.21	0.29	1.00
DAX30	1%	0.50	0.41	0.43	0.34	1.00
	5%	0.85	0.43	0.84	0.66	0.99
	10%	0.85	0.42	0.78	0.50	0.66
	90%	0.73	0.35	0.74	0.57	1.00
	95%	0.52	0.33	0.57	0.35	1.00
	99%	0.79	0.74	0.73	0.78	1.00

Table 2 For the in-sample data of Period 2 of the S&P500 and the DAX30, the PlugIn coefficient, β_{IQ} , of the IQ estimator for different values of θ .

	θ	Sym Abs Value	Asym Slope	Ind Garch	AR-GARCH	Adapt
S&P500	1%	1.85**	0.64**	1.35	1.29**	1.73**
	5%	1.50**	0.65**	1.25	1.31	0.45
	10%	1.76**	0.64**	1.30	1.30**	0.48
	90%	1.54	0.09*	0.56**	1.24	1.49**
	95%	1.48	0.04	0.53	0.59	1.53
	99%	1.61**	0.05*	0.25	0.19	1.55
DAX30	1%	0.05	0.16	0.39	0.32	0.97**
	5%	1.47**	0.44**	1.19*	1.14*	1.47*
	10%	1.73**	1.18**	1.23*	1.27**	1.66*
	90%	0.98	0.10*	0.53**	0.53*	1.05
	95%	0.18	0.12	0.45*	0.47	1.39
	99%	0.99**	0.35*	0.93**	0.93**	0.92

Note: Significance at 5% and 1% levels is indicated by * and **, respectively.

Table 3 Hit percentage and DQ test p-values for Post-sample 1 (volatile) of the S&P500.

		Hit %				DQ p-value			
		1%	5%	95%	99%	1%	5%	95%	99%
IQ		1.4	7.6**	93.0*	97.4**	0.85	0.17	0.02*	0.00**
Sym Abs Value CAViaR	CAViaR	0.4	4.4	95.4	99.0	0.91	0.02*	0.78	0.98
	SimpAvg	0.4	5.6	95.0	98.6	0.92	0.06	0.31	0.03*
	LinearComb	0.2	4.4	94.6	99.4	0.78	0.87	0.59	0.98
	WtdAvg	0.4	5.6	94.2	98.8	0.90	0.05*	0.43	0.99
	WtdAvgExp	0.6	5.6	94.2	99.0	0.99	0.05*	0.43	1.00
	PlugIn	1.0	6.4	94.4	99.2	0.93	0.21	0.71	0.97
	Asym Slope CAViaR	CAViaR	1.8	7.6**	95.4	99.4	0.01**	0.00**	0.82
SimpAvg		1.2	6.6	94.0	99.0	0.71	0.00**	0.43	1.00
LinearComb		1.2	5.2	94.8	99.4	0.75	0.23	0.78	0.96
WtdAvg		1.2	6.6	94.8	99.4	0.74	0.00**	0.78	0.96
WtdAvgExp		1.2	6.6	93.0*	98.4	0.65	0.00**	0.01*	0.00**
PlugIn		1.8	6.4	95.2	99.6	0.08	0.02*	0.68	0.90
Ind GARCH CAViaR		CAViaR	0.6	4.4	94.6	98.6	0.74	0.03*	0.23
	SimpAvg	0.8	5.4	94.6	98.0*	0.76	0.04*	0.18	0.00**
	LinearComb	1.2	4.0	95.4	99.2	0.93	0.83	0.76	1.00
	WtdAvg	1.4	5.8	94.6	98.6	0.85	0.06	0.19	0.03*
	WtdAvgExp	1.4	6.0	94.6	98.6	0.83	0.07	0.20	0.03*
	PlugIn	1.2	6.4	93.8	98.8	0.98	0.16	0.20	1.00
	Adaptive CAViaR	CAViaR	1.2	4.8	94.8	98.8	0.95	0.57	0.13
SimpAvg		1.2	5.2	93.8	99.0	0.98	0.36	0.08	0.00**
LinearComb		1.0	6.2	94.0	99.2	1.00	0.60	0.13	0.00**
WtdAvg		1.4	6.0	93.0*	98.0*	0.85	0.78	0.02*	0.01*
WtdAvgExp		1.4	6.0	93.0*	98.4	0.85	0.78	0.02*	0.02*
PlugIn		1.2	6.4	94.4	99.0	0.98	0.53	0.70	0.00**

Note: Larger values of the DQ test p-value are better. Significance at 5% and 1% levels is indicated by * and **, respectively.

Table 4 Hit percentage and DQ test p-values for Post-sample 2 (tranquil) of the S&P500.

		Hit %				DQ p-value			
		1%	5%	95%	99%	1%	5%	95%	99%
	IQ	1.0	5.0	95.2	99.2	1.00	0.08	0.00**	0.98
Sym Abs Value CAViaR	CAViaR	0.2	2.2**	96.4	99.4	0.75	0.01**	0.40	0.77
	SimpAvg	0.2	3.6	95.8	99.2	0.76	0.26	0.32	1.00
	LinearComb	0.2	6.2	92.6*	99.0	0.78	0.04*	0.01**	1.00
	WtdAvg	1.0	5.0	95.8	99.2	1.00	0.08	0.19	1.00
	WtdAvgExp	1.0	5.0	95.8	99.2	1.00	0.08	0.19	1.00
	PlugIn	0.8	8.2**	91.8**	97.8**	0.97	0.00**	0.00**	0.13
	Asym Slope CAViaR	CAViaR	0.2	1.6**	98.0**	100.0*	0.77	0.00**	0.07
SimpAvg		0.2	2.8*	96.2	99.6	0.77	0.14	0.02*	0.93
LinearComb		0.4	4.4	95.0	99.0	0.90	0.71	0.16	1.00
WtdAvg		0.4	2.2**	97.6**	99.8	0.90	0.03*	0.04*	0.78
WtdAvgExp		1.0	5.0	95.2	99.8	1.00	0.08	0.00**	0.78
PlugIn		1.8	6.8	96.0	99.0	0.01*	0.17	0.62	1.00
Ind GARCH CAViaR		CAViaR	0.2	2.0**	95.8	99.2	0.76	0.01**	0.66
	SimpAvg	0.2	3.4	95.8	99.2	0.77	0.27	0.32	1.00
	LinearComb	0.6	6.2	93.4	99.0	0.98	0.04*	0.08	1.00
	WtdAvg	1.0	5.0	95.8	99.2	1.00	0.08	0.36	1.00
	WtdAvgExp	1.0	5.0	95.8	99.0	1.00	0.08	0.36	1.00
	PlugIn	0.6	8.6**	93.2	99.0	0.95	0.00**	0.05	1.00
	Adaptive CAViaR	CAViaR	1.0	4.2	95.4	98.8	0.44	0.15	0.05*
SimpAvg		1.0	4.4	95.4	99.4	0.78	0.40	0.20	0.96
LinearComb		0.2	7.2*	92.0**	98.2	0.78	0.00**	0.00**	0.54
WtdAvg		1.0	4.8	95.2	99.2	1.00	0.07	0.00**	0.98
WtdAvgExp		1.0	4.8	95.2	99.2	1.00	0.09	0.00**	0.98
PlugIn		0.6	6.4	90.6**	97.6**	0.95	0.11	0.00**	0.03*

Note: Larger values of the DQ test p-value are better. Significance at 5% and 1% levels is indicated by * and **, respectively.

Table 5 Summary of results for each quantile. For each post-sample period and each method, the value shown is the number of CAViaR models for which the test null hypothesis was rejected. Smaller values are better. In the final seven rows, bold indicates the best performing method in each column.

		Hit(%)					DQ p-value				
		1%	5%	95%	99%	Mean	1%	5%	95%	99%	Mean
S&P500 Post-sample 1 (volatile) (Summary of Table 3)	IQ	0	4	4	4	3	0	0	4	4	2
	CAViaR	0	1	0	0	0.25	1	3	0	2	1.5
	SimpAvg	0	0	0	1	0.25	0	2	0	3	1.25
	LinearComb	0	0	0	0	0	0	0	0	1	0.25
	WtdAvg	0	0	1	1	0.5	0	2	1	2	1.25
	WtdAvgExp	0	0	2	0	0.5	0	2	2	3	1.75
	PlugIn	0	0	0	0	0	0	1	0	1	0.5
S&P500 Post-sample 2 (tranquil) (Summary of Table 4)	IQ	0	0	0	0	0	0	0	4	0	1
	CAViaR	0	3	1	1	1.25	0	3	1	1	1.25
	SimpAvg	0	1	0	0	0.25	0	0	1	0	0.25
	LinearComb	0	1	2	0	0.75	0	3	2	0	1.25
	WtdAvg	0	1	1	0	0.5	0	1	2	0	0.75
	WtdAvgExp	0	0	0	0	0	0	0	2	0	0.5
	PlugIn	0	2	2	2	1.5	1	2	2	1	1.5
DAX30 Post-sample 1 (volatile)	IQ	0	4	0	0	1	4	4	0	4	3
	CAViaR	0	2	3	1	1.5	1	3	0	1	1.25
	SimpAvg	0	3	0	1	1	2	0	0	2	1
	LinearComb	0	1	1	0	0.5	1	1	0	0	0.5
	WtdAvg	0	4	0	1	1.25	2	1	0	2	1.25
	WtdAvgExp	0	4	0	0	1	2	1	0	1	1
	PlugIn	0	0	1	0	0.25	2	3	1	2	2
DAX30 Post-sample 2 (tranquil)	IQ	0	0	0	0	0	4	0	0	0	1
	CAViaR	0	1	0	0	0.25	3	0	0	0	0.75
	SimpAvg	0	0	0	0	0	4	0	0	0	1
	LinearComb	0	0	0	0	0	2	0	0	3	1.25
	WtdAvg	0	0	0	0	0	4	0	0	0	1
	WtdAvgExp	0	0	0	0	0	4	0	0	0	1
	PlugIn	0	0	0	0	0	4	0	0	4	2
Mean	IQ	0	2	1	1	1	2	1	2	2	1.75
	CAViaR	0	1.75	1	0.5	0.81	1.25	2.25	0.25	1	1.19
	SimpAvg	0	1	0	0.5	0.38	1.5	0.5	0.25	1.25	0.88
	LinearComb	0	0.5	0.75	0	0.31	0.75	1	0.5	1	0.81
	WtdAvg	0	1.25	0.5	0.5	0.56	1.5	1	0.75	1	1.06
	WtdAvgExp	0	1	0.5	0	0.38	1.5	0.75	1	1	1.06
	PlugIn	0	0.5	0.75	0.5	0.44	1.75	1.5	0.75	2	1.50