

Advanced Philosophy of Physics: General Covariance and Background Independence

James Read¹

¹Faculty of Philosophy, University of Oxford, UK, OX2 6GG

MT20-W3

The plan

- W1: The philosophy of symmetries
- W2: The hole argument
- W3: General covariance and background independence
- W4: The dynamical approach to spacetime

What's special about general relativity?

Today

Preliminaries

General covariance

Diffeomorphism invariance

Absolute objects

Variational principles

Belot's approach

Fixed fields

Absolute fields

Conclusions

Today

Preliminaries

General covariance

Diffeomorphism invariance

Absolute objects

Variational principles

Belot's approach

Fixed fields

Absolute fields

Conclusions

Models

- ▶ The *kinematically possible models* (KPMs) of a given theory are picked out by tuples $\langle M, \Phi_1, \dots, \Phi_n \rangle$, with
 1. a manifold, M ;
 2. fields on M , Φ_1, \dots, Φ_n .

Models

- ▶ The *kinematically possible models* (KPMs) of a given theory are picked out by tuples $\langle M, \Phi_1, \dots, \Phi_n \rangle$, with
 1. a manifold, M ;
 2. fields on M , Φ_1, \dots, Φ_n .
- ▶ The *dynamically possible models* (DPMs) of a given theory are those KPMs in which the Φ_i satisfy certain dynamical equations.

Dynamical versus fixed fields

- ▶ *Dynamical fields* are not fixed identically in all KPMs, and may have their own associated dynamical equations.
 - ▶ E.g. g_{ab} in GR.

Dynamical versus fixed fields

- ▶ *Dynamical fields* are not fixed identically in all KPMs, and may have their own associated dynamical equations.
 - ▶ E.g. g_{ab} in GR.
- ▶ *Fixed fields* are fixed identically in all KPMs, and do not have their own associated dynamical equations.
 - ▶ E.g. η_{ab} in SR.

Three toy theories

Three toy theories

SR1:

- ▶ KPMs: $\langle M, \eta_{ab}, \varphi \rangle$.
- ▶ DPMs picked out by $\eta_{ab} \nabla^a \nabla^b \varphi = 0$.
- ▶ η_{ab} *fixed* identically in all KPMs.

Three toy theories

SR1:

- ▶ KPMs: $\langle M, \eta_{ab}, \varphi \rangle$.
- ▶ DPMs picked out by $\eta_{ab} \nabla^a \nabla^b \varphi = 0$.
- ▶ η_{ab} *fixed* identically in all KPMs.

SR2:

- ▶ KPMs: $\langle M, g_{ab}, \varphi \rangle$.
- ▶ DPMs picked out by $g_{ab} \nabla^a \nabla^b \varphi = 0$ and $R^a_{bcd} = 0$.
- ▶ Flatness of g_{ab} fixed *dynamically* by $R^a_{bcd} = 0$.

Three toy theories

SR1:

- ▶ KPMs: $\langle M, \eta_{ab}, \varphi \rangle$.
- ▶ DPMs picked out by $\eta_{ab} \nabla^a \nabla^b \varphi = 0$.
- ▶ η_{ab} *fixed* identically in all KPMs.

SR2:

- ▶ KPMs: $\langle M, g_{ab}, \varphi \rangle$.
- ▶ DPMs picked out by $g_{ab} \nabla^a \nabla^b \varphi = 0$ and $R^a_{bcd} = 0$.
- ▶ Flatness of g_{ab} fixed *dynamically* by $R^a_{bcd} = 0$.

GR1:

- ▶ KPMs: $\langle M, g_{ab}, \varphi \rangle$.
- ▶ DPMs picked out by $g_{ab} \nabla^a \nabla^b \varphi = 0$ and $G_{ab} = 8\pi T_{ab}$.

Today

Preliminaries

General covariance

Diffeomorphism invariance

Absolute objects

Variational principles

Belot's approach

Fixed fields

Absolute fields

Conclusions

Generalising the relativity principle

- ▶ One of Einstein's motivations in constructing general relativity was (as the name suggests!) to generalise the relativity principle: the laws of physics should take the same form not just in the inertial frames of reference, but in *all* frames of reference.

Generalising the relativity principle

- ▶ One of Einstein's motivations in constructing general relativity was (as the name suggests!) to generalise the relativity principle: the laws of physics should take the same form not just in the inertial frames of reference, but in *all* frames of reference.
- ▶ In this way, the need to anchor one's physics to the—somewhat etherial!—inertial frames would be eliminated.

Generalising the relativity principle

- ▶ One of Einstein's motivations in constructing general relativity was (as the name suggests!) to generalise the relativity principle: the laws of physics should take the same form not just in the inertial frames of reference, but in *all* frames of reference.
- ▶ In this way, the need to anchor one's physics to the—somewhat etherial!—inertial frames would be eliminated.

Definition

(General covariance) *A formulation of a theory is generally covariant iff the equations expressing its laws are written in a form that holds with respect to all members of a set of coordinate systems that are related by smooth but otherwise arbitrary transformations.*

Einstein's struggle with general covariance

- ▶ In 1913, Einstein and Grossmann developed a theory with all the essential components of general relativity.

Einstein's struggle with general covariance

- ▶ In 1913, Einstein and Grossmann developed a theory with all the essential components of general relativity.
- ▶ However, between 1913 and 1915, Einstein 'fell into the hole': he rejected all generally covariant theories, in light of the radical indeterminism which seemed to arise in light of the hole argument.

Einstein's struggle with general covariance

- ▶ In 1913, Einstein and Grossmann developed a theory with all the essential components of general relativity.
- ▶ However, between 1913 and 1915, Einstein 'fell into the hole': he rejected all generally covariant theories, in light of the radical indeterminism which seemed to arise in light of the hole argument.
- ▶ Einstein would then convince himself, via his 'point coincidence argument', that the hole argument was not problematic; he thus returned to general covariance and finalised general relativity.

Einstein's struggle with general covariance

- ▶ In 1913, Einstein and Grossmann developed a theory with all the essential components of general relativity.
- ▶ However, between 1913 and 1915, Einstein 'fell into the hole': he rejected all generally covariant theories, in light of the radical indeterminism which seemed to arise in light of the hole argument.
- ▶ Einstein would then convince himself, via his 'point coincidence argument', that the hole argument was not problematic; he thus returned to general covariance and finalised general relativity.
- ▶ The hole argument was, more or less, lost to history until it was revived in the latter half of the 20th Century by philosophers of physics.

The Kretschmann objection

- Things did not stay problem-free for general covariance for very long.

The Kretschmann objection

- ▶ Things did not stay problem-free for general covariance for very long.
- ▶ In 1917, Kretschmann argued to Einstein that *any* theory can be written in a generally covariant form.

The Kretschmann objection

- ▶ Things did not stay problem-free for general covariance for very long.
- ▶ In 1917, Kretschmann argued to Einstein that *any* theory can be written in a generally covariant form.

By means of a purely mathematical reformulation of the equations representing the theory, and with, at most, mathematical complications connected with that reformulation, any physical theory can be brought into agreement with any, arbitrary relativity postulate, even the most general one, and this without modifying any of its content that can be tested by observation.
(Kretschmann 1917, trans. Norton 1993, p. 818)

Modern endorsements

The Kretschmann point has become orthodoxy. For example, in 1983, Friedman wrote:

... the principle of general covariance has no physical content whatever: it specifies no particular physical theory; rather it merely expresses our commitment to a certain style of formulating physical theories. (Friedman 1983, p. 55)

Modern endorsements

The Kretschmann point has become orthodoxy. For example, in 1983, Friedman wrote:

... the principle of general covariance has no physical content whatever: it specifies no particular physical theory; rather it merely expresses our commitment to a certain style of formulating physical theories. (Friedman 1983, p. 55)

There are two different ways in which the Kretschmann point can be made, which I'll now go over.

Kretschmann objection: first presentation

The first way of making the Kretschmann point proceeds as follows:

Kretschmann objection: first presentation

The first way of making the Kretschmann point proceeds as follows:

1. Take the laws of some physical theory, written in a coordinate system.

Kretschmann objection: first presentation

The first way of making the Kretschmann point proceeds as follows:

1. Take the laws of some physical theory, written in a coordinate system.
2. Transform those laws into an arbitrary coordinate system.

Kretschmann objection: first presentation

The first way of making the Kretschmann point proceeds as follows:

1. Take the laws of some physical theory, written in a coordinate system.
2. Transform those laws into an arbitrary coordinate system.
3. The resulting equations might end up being more complex, but by construction they hold in all coordinate systems!

Kretschmann objection: first presentation

The first way of making the Kretschmann point proceeds as follows:

1. Take the laws of some physical theory, written in a coordinate system.
2. Transform those laws into an arbitrary coordinate system.
3. The resulting equations might end up being more complex, but by construction they hold in all coordinate systems!
4. So this is the generally covariant version of the theory under consideration.

Example: Klein-Gordon equation

First, we'll write the Klein-Gordon equation (a wave equation) in index notation. (This is just a presentational reformulation.)

$$-\frac{1}{c^2} \frac{\partial^2 \phi}{\partial t^2} + \frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} + \frac{\partial^2 \phi}{\partial z^2} = 0$$

$$\Rightarrow \left(\begin{array}{cccc} \frac{1}{c} \frac{\partial}{\partial t} & \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \end{array} \right) \left(\begin{array}{cccc} -1 & & & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{array} \right) \left(\begin{array}{c} \frac{1}{c} \frac{\partial}{\partial t} \\ \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \\ \frac{\partial}{\partial z} \end{array} \right) \phi = 0$$

$$\Rightarrow \eta_{\mu\nu} \partial^\mu \partial^\nu \phi = 0.$$

Example: Klein-Gordon equation

Then, we'll transform the Klein-Gordon equation to an arbitrary coordinate system:

$$\begin{aligned}\eta_{\mu\nu} \partial^\mu \partial^\nu \varphi &= 0 \\ \eta_{\mu\nu} \frac{\partial}{\partial x_\mu} \frac{\partial}{\partial x_\nu} \varphi &= 0 \\ \longrightarrow \eta_{\mu\nu} \frac{\partial x_{\mu'}}{\partial x_\mu} \frac{\partial}{\partial x_{\mu'}} \left(\frac{\partial x_{\nu'}}{\partial x_\nu} \frac{\partial}{\partial x_{\nu'}} \varphi \right) &= 0 \\ \eta_{\mu\nu} \frac{\partial x_{\mu'}}{\partial x_\mu} \left(\frac{\partial^2 x_{\nu'}}{\partial x_{\mu'} \partial x_\nu} \frac{\partial}{\partial x_{\nu'}} \varphi + \frac{\partial x_{\nu'}}{\partial x_\nu} \frac{\partial}{\partial x_{\mu'}} \frac{\partial}{\partial x_{\nu'}} \varphi \right) &= 0 \\ \eta_{\mu\nu} \frac{\partial^2 x_{\nu'}}{\partial x_\mu \partial x_\nu} \partial^{\nu'} \varphi + \eta_{\mu\nu} \frac{\partial x_{\mu'}}{\partial x_\mu} \frac{\partial x_{\nu'}}{\partial x_\nu} \partial^{\mu'} \partial^{\nu'} \varphi &= 0.\end{aligned}$$

Example: Klein-Gordon equation

Then, we'll transform the Klein-Gordon equation to an arbitrary coordinate system:

$$\begin{aligned}\eta_{\mu\nu} \partial^\mu \partial^\nu \varphi &= 0 \\ \eta_{\mu\nu} \frac{\partial}{\partial x_\mu} \frac{\partial}{\partial x_\nu} \varphi &= 0 \\ \longrightarrow \eta_{\mu\nu} \frac{\partial x_{\mu'}}{\partial x_\mu} \frac{\partial}{\partial x_{\mu'}} \left(\frac{\partial x_{\nu'}}{\partial x_\nu} \frac{\partial}{\partial x_{\nu'}} \varphi \right) &= 0 \\ \eta_{\mu\nu} \frac{\partial x_{\mu'}}{\partial x_\mu} \left(\frac{\partial^2 x_{\nu'}}{\partial x_{\mu'} \partial x_\nu} \frac{\partial}{\partial x_{\nu'}} \varphi + \frac{\partial x_{\nu'}}{\partial x_\nu} \frac{\partial}{\partial x_{\mu'}} \frac{\partial}{\partial x_{\nu'}} \varphi \right) &= 0 \\ \eta_{\mu\nu} \frac{\partial^2 x_{\nu'}}{\partial x_\mu \partial x_\nu} \partial^{\nu'} \varphi + \eta_{\mu\nu} \frac{\partial x_{\mu'}}{\partial x_\mu} \frac{\partial x_{\nu'}}{\partial x_\nu} \partial^{\mu'} \partial^{\nu'} \varphi &= 0.\end{aligned}$$

This is a generally covariant version of our theory! Note the extra term in the non-inertial frame (cf. fictitious forces).

Kretschmann objection: second presentation

The second way of making the Kretschmann point proceeds as follows:

Kretschmann objection: second presentation

The second way of making the Kretschmann point proceeds as follows:

1. Argue that all theories can be written in the coordinate-independent language of differential geometry.

Kretschmann objection: second presentation

The second way of making the Kretschmann point proceeds as follows:

1. Argue that all theories can be written in the coordinate-independent language of differential geometry.
(**Question:** Why think this? Is it obvious? See e.g. (Norton 1993) for discussion.)

Kretschmann objection: second presentation

The second way of making the Kretschmann point proceeds as follows:

1. Argue that all theories can be written in the coordinate-independent language of differential geometry.
(**Question:** Why think this? Is it obvious? See e.g. (Norton 1993) for discussion.)
2. Note that *a fortiori* the equations of the theory hold in all coordinate systems.

Kretschmann objection: second presentation

The second way of making the Kretschmann point proceeds as follows:

1. Argue that all theories can be written in the coordinate-independent language of differential geometry.
(**Question:** Why think this? Is it obvious? See e.g. (Norton 1993) for discussion.)
2. Note that *a fortiori* the equations of the theory hold in all coordinate systems.
3. So this is the generally covariant version of the theory under consideration.

Examples

- ▶ Coordinate-independent presentations of a special relativistic Klein-Gordon theory are **SR1** and **SR2**.

Examples

- ▶ Coordinate-independent presentations of a special relativistic Klein-Gordon theory are **SR1** and **SR2**.
- ▶ We saw a coordinate independent presentation of Newtonian mechanics last week:
 - ▶ KPMs: $\langle M, t_{ab}, h^{ab}, \nabla, \rho, \varphi \rangle$.
 - ▶ DPMs: $h^{ab} \nabla_a \nabla_b \varphi = 4\pi \rho$.

The upshot

It's not at all obvious that general covariance affords an adequate means of distinguishing general relativity from other spacetime theories.

Today

Preliminaries

General covariance

Diffeomorphism invariance

Absolute objects

Variational principles

Belot's approach

Fixed fields

Absolute fields

Conclusions

Diffeomorphism invariance

- Often, general covariance is conflated with diffeomorphism invariance.

Diffeomorphism invariance

- ▶ Often, general covariance is conflated with diffeomorphism invariance.
- ▶ Pooley (2017) provides different definitions of the two notions, such that—perhaps!—it is the latter notion which distinguishes general relativity from other theories.

Diffeomorphism invariance

- ▶ Often, general covariance is conflated with diffeomorphism invariance.
- ▶ Pooley (2017) provides different definitions of the two notions, such that—perhaps!—it is the latter notion which distinguishes general relativity from other theories.

Definition

(Diffeomorphism invariance) A theory \mathcal{T} is diffeomorphism invariant iff, if $\langle M, F_1, \dots, F_n, D_1, \dots, D_m \rangle$ is a DPM of \mathcal{T} (where the F_i are fixed fields and the D_i are dynamical fields), then so is $\langle M, F_1, \dots, F_n, d^*D_1, \dots, d^*D_m \rangle$, for all $d \in \text{Diff}(M)$.

Assessing diffeomorphism invariance

- ▶ General relativistic theories such as **GR1** are diffeomorphism invariant.

Assessing diffeomorphism invariance

- ▶ General relativistic theories such as **GR1** are diffeomorphism invariant.
- ▶ **SR1** fails to be diffeomorphism invariant, but **SR2** *is* diffeomorphism invariant.

Assessing diffeomorphism invariance

- ▶ General relativistic theories such as **GR1** are diffeomorphism invariant.
- ▶ **SR1** fails to be diffeomorphism invariant, but **SR2** *is* diffeomorphism invariant.
- ▶ Therefore, diffeomorphism invariance does not serve to distinguish special from general relativistic theories.

Background independence

Thought: If what distinguishes general relativistic theories isn't their general covariance, and it isn't their diffeomorphism invariance, it must be some other property. We'll label this *background independence*, as a placeholder.

Background independence

Thought: If what distinguishes general relativistic theories isn't their general covariance, and it isn't their diffeomorphism invariance, it must be some other property. We'll label this *background independence*, as a placeholder.

But what *is* background independence?

Today

Preliminaries

General covariance

Diffeomorphism invariance

Absolute objects

Variational principles

Belot's approach

Fixed fields

Absolute fields

Conclusions

Absolute objects

Anderson (1967) and Friedman (1983) both proposed that the background independence of general relativity consists in its lacking 'absolute objects'. They offered slightly different definitions of the notion; we'll use the following (which is closer to Friedman's):

Absolute objects

Anderson (1967) and Friedman (1983) both proposed that the background independence of general relativity consists in its lacking 'absolute objects'. They offered slightly different definitions of the notion; we'll use the following (which is closer to Friedman's):

Definition

***(Absolute object)** A geometrical object the same (up to isomorphism) in all DPMs of a theory.*

Absolute objects

Anderson (1967) and Friedman (1983) both proposed that the background independence of general relativity consists in its lacking 'absolute objects'. They offered slightly different definitions of the notion; we'll use the following (which is closer to Friedman's):

Definition

***(Absolute object)** A geometrical object the same (up to isomorphism) in all DPMs of a theory.*

Definition

***(Background independence, absolute objects)** A theory is background independent iff it has no absolute objects in its formulation.*

Problems for the absolute objects proposal

There are three kinds of problem case for the absolute object programme:

Problems for the absolute objects proposal

There are three kinds of problem case for the absolute object programme:

1. Cases in which structure that intuitively should count as background is not classified as absolute.
2. Cases in which structure that intuitively should not count as background is classified as absolute.
3. The observation that general relativity itself seems to have absolute objects.

Type I problems: Torretti constant curvature

- ▶ Torretti considers a modified Newtonian theory, in which each model's spatial metric has constant curvature, but different models have different values of that curvature.

Type I problems: Torretti constant curvature

- ▶ Torretti considers a modified Newtonian theory, in which each model's spatial metric has constant curvature, but different models have different values of that curvature.
- ▶ Because the spatial metric in every model has constant curvature, “such a theory has something rather like an absolute object in it” (Pitts 2006, p. 363).

Type I problems: Torretti constant curvature

- ▶ Torretti considers a modified Newtonian theory, in which each model's spatial metric has constant curvature, but different models have different values of that curvature.
- ▶ Because the spatial metric in every model has constant curvature, “such a theory has something rather like an absolute object in it” (Pitts 2006, p. 363).
- ▶ Nevertheless, the failure of the metrics to be locally diffeomorphically equivalent for distinct curvature values entails that the metric tensor does not satisfy Anderson's definition of an absolute object.

Responses to type I problems

- ▶ Pitts (2006, pp. 16-17) argues that Torretti's example does not succeed, for if one decomposes the spatial metric into a conformal spatial metric density and a scalar density, then the former *is* an absolute object.

Responses to type I problems

- ▶ Pitts (2006, pp. 16-17) argues that Torretti's example does not succeed, for if one decomposes the spatial metric into a conformal spatial metric density and a scalar density, then the former *is* an absolute object.
- ▶ More generally, Torretti's intuition about background structure would seem to over-generate, for almost all theories have some models in which a given piece of structure does not vary with respect to some given (possibly temporal) parameter. (Consider, e.g., a $3 + 1$ decomposition of general relativity with constant curvature of the induced metric on the spatial slices.)

Type II problems: Jones-Geroch dust

- Consider a general relativistic theory in which the metric field is coupled to matter characterised only by a 4-velocity field U^a and a mass density.

Type II problems: Jones-Geroch dust

- ▶ Consider a general relativistic theory in which the metric field is coupled to matter characterised only by a 4-velocity field U^a and a mass density.
- ▶ Then, as Friedman states, “since any two timelike, nowhere-vanishing vector fields defined on a relativistic space-time are d -equivalent, it follows that any such vector field counts as an absolute object ... and this is surely counter-intuitive” (Friedman 1983, p. 59).

Type III problems: GR's absolute object

In GR, we can always decompose the metric as

$$g_{\mu\nu} = \hat{g}_{\mu\nu} \sqrt{-g}^{2/n},$$

where $\hat{g}_{\mu\nu}$ is a conformal metric density, and $\sqrt{-g}$ is the square root of the (negative of the) determinant of the metric.

Type III problems: GR's absolute object

In GR, we can always decompose the metric as

$$g_{\mu\nu} = \hat{g}_{\mu\nu} \sqrt{-g}^{2/n},$$

where $\hat{g}_{\mu\nu}$ is a conformal metric density, and $\sqrt{-g}$ is the square root of the (negative of the) determinant of the metric.

GR has an absolute object! This absolute object $[\sqrt{-g}]$ is a scalar density of nonzero weight, because every neighbourhood in every model spacetime admits coordinates (at least locally) in which the component of the scalar density has a value of -1 . (Pitts 2006, p. 366)

Type III problems: GR's absolute object

In GR, we can always decompose the metric as

$$g_{\mu\nu} = \hat{g}_{\mu\nu} \sqrt{-g}^{2/n},$$

where $\hat{g}_{\mu\nu}$ is a conformal metric density, and $\sqrt{-g}$ is the square root of the (negative of the) determinant of the metric.

GR has an absolute object! This absolute object $[\sqrt{-g}]$ is a scalar density of nonzero weight, because every neighbourhood in every model spacetime admits coordinates (at least locally) in which the component of the scalar density has a value of -1 . (Pitts 2006, p. 366)

In any neighbourhood of the manifold in any model of GR, one can find a coordinate system such that the object $\sqrt{-g}$ takes the value -1 . Thus, this is a geometrical object which is the same (up to isomorphism) in all DPMs of the theory.

Today

Preliminaries

General covariance

Diffeomorphism invariance

Absolute objects

Variational principles

Belot's approach

Fixed fields

Absolute fields

Conclusions

Variational principles

Pooley (2017) proposes a different definition of background independence, in terms of variational principles:

Definition

(Background independence, variational) A theory is background independent iff its solution space is determined by a generally-covariant action, (i) all of whose dependent variables are subject to Hamilton's principle, and (ii) all of whose dependent variables represent physical fields.

Assessing the variational principle definition

- General relativity satisfies this definition: the field equations can be derived from the *Einstein-Hilbert action*:

$$S_{\text{EH}} = \frac{1}{16\pi} \int_M d^4x \sqrt{-g} R$$

Assessing the variational principle definition

- General relativity satisfies this definition: the field equations can be derived from the *Einstein-Hilbert action*:

$$S_{\text{EH}} = \frac{1}{16\pi} \int_M d^4x \sqrt{-g} R$$

- **SR1** violates the definition: η_{ab} is not subject to Hamilton's principle.

Assessing the variational principle definition

- ▶ General relativity satisfies this definition: the field equations can be derived from the *Einstein-Hilbert action*:

$$S_{\text{EH}} = \frac{1}{16\pi} \int_M d^4x \sqrt{-g} R$$

- ▶ **SR1** violates the definition: η_{ab} is not subject to Hamilton's principle.
- ▶ **SR2**...?

Variational principles

Rosen action for **SR2**:

$$S_{\text{SR2}} = \int d^4x \sqrt{-g} \left(\Theta^{abcd} R_{abcd} + g^{ab} \nabla_a \varphi \nabla_b \varphi \right)$$

Variational principles

Rosen action for **SR2**:

$$S_{\text{SR2}} = \int d^4x \sqrt{-g} \left(\Theta^{abcd} R_{abcd} + g^{ab} \nabla_a \varphi \nabla_b \varphi \right)$$

- All dependent variables subject to Hamilton's principle.

Variational principles

Rosen action for **SR2**:

$$S_{\text{SR2}} = \int d^4x \sqrt{-g} \left(\Theta^{abcd} R_{abcd} + g^{ab} \nabla_a \varphi \nabla_b \varphi \right)$$

- ▶ All dependent variables subject to Hamilton's principle.
- ▶ But **SR2** still not background independent, if Θ^{abcd} regarded as 'unphysical'.

Variational principles

Further worries:

Variational principles

Further worries:

- ▶ Parameterised versions of special relativity, written in terms of *clock fields* X^a . ($\eta_{\mu\nu} := \eta_{ab} \partial_\mu X^a \partial_\nu X^b$.)

Variational principles

Further worries:

- ▶ Parameterised versions of special relativity, written in terms of *clock fields* X^a . ($\eta_{\mu\nu} := \eta_{ab} \partial_\mu X^a \partial_\nu X^b$.)
- ▶ This definition assumes that every background independent theory admits of a Lagrangian formulation.

Today

Preliminaries

General covariance

Diffeomorphism invariance

Absolute objects

Variational principles

Belot's approach

Fixed fields

Absolute fields

Conclusions

Belot's account

- ▶ KPMs of a theory given by $\langle M, O_1, \dots, O_n \rangle$.

Belot's account

- ▶ KPMs of a theory given by $\langle M, O_1, \dots, O_n \rangle$.
- ▶ Identify *geometrical structure*; label by O^G , e.g. $\langle M, O^G, \dots, O_{n-1} \rangle$.

Belot's account

- ▶ KPMs of a theory given by $\langle M, O_1, \dots, O_n \rangle$.
- ▶ Identify *geometrical structure*; label by O^G , e.g. $\langle M, O^G, \dots, O_{n-1} \rangle$.
- ▶ *Geometrical degrees of freedom* are degrees of freedom needed to parameterise variation of O^G across DPMs.

Belot's account

- ▶ KPMs of a theory given by $\langle M, O_1, \dots, O_n \rangle$.
- ▶ Identify *geometrical structure*; label by O^G , e.g. $\langle M, O^G, \dots, O_{n-1} \rangle$.
- ▶ *Geometrical degrees of freedom* are degrees of freedom needed to parameterise variation of O^G across DPMs.
- ▶ *Physical degrees of freedom* are degrees of freedom needed to parameterise variation of $\langle O^G, \dots, O_{n-1} \rangle$ across DPMs (modulo gauge equivalence).

Belot's account

Definition

(Full background dependence, Belot) *A field theory is fully background dependent if it has no geometrical degrees of freedom: every solution is assigned the same spacetime geometry as every other solution.*

Definition

(Full background independence, Belot) *A field theory is fully background independent if all of its physical degrees of freedom correspond to geometrical degrees of freedom.*

Belot's account

- In **SR1** and **SR2**, the metric field is fixed (up to isomorphism) in all DPMs.

Belot's account

- ▶ In **SR1** and **SR2**, the metric field is fixed (up to isomorphism) in all DPMs.
- ▶ So there are no geometrical degrees of freedom.

Belot's account

- ▶ In **SR1** and **SR2**, the metric field is fixed (up to isomorphism) in all DPMs.
- ▶ So there are no geometrical degrees of freedom.
- ▶ So these theories are background dependent, for Belot.

Belot's account

- *Prima facie*, general relativity is background independent: each configuration of the matter fields looks to correspond to a unique configuration of the metric field ('geometry'), via the Einstein equations,

$$G_{ab} = 8\pi T_{ab}.$$

Belot's account

- *Prima facie*, general relativity is background independent: each configuration of the matter fields looks to correspond to a unique configuration of the metric field ('geometry'), via the Einstein equations,

$$G_{ab} = 8\pi T_{ab}.$$

- But: If $\langle M, g_{ab}, F_{ab} \rangle$ is a model of Einstein-Maxwell theory, then so is $\langle M, g_{ab}, {}^*F_{ab} \rangle$. Different matter, same geometry.

Belot's account

Another worry:

Belot's account

Another worry:

- ▶ Consider a theory with two 'geometrical' fields, g_1 and g_2 , and one non-'geometrical' field, φ .

Belot's account

Another worry:

- ▶ Consider a theory with two 'geometrical' fields, g_1 and g_2 , and one non-'geometrical' field, φ .
- ▶ KPMs $\langle M, g_1, g_2, \varphi \rangle$.

Belot's account

Another worry:

- ▶ Consider a theory with two 'geometrical' fields, g_1 and g_2 , and one non-'geometrical' field, φ .
- ▶ KPMs $\langle M, g_1, g_2, \varphi \rangle$.
- ▶ Let g_1 be a fixed field; let g_2 co-vary dynamically with φ .

Belot's account

Another worry:

- ▶ Consider a theory with two 'geometrical' fields, g_1 and g_2 , and one non-'geometrical' field, φ .
- ▶ KPMs $\langle M, g_1, g_2, \varphi \rangle$.
- ▶ Let g_1 be a fixed field; let g_2 co-vary dynamically with φ .
- ▶ For Belot, this theory is background independent, in spite of it having background structure g_1 .

Belot's account

Another worry:

- ▶ Consider a theory with two 'geometrical' fields, g_1 and g_2 , and one non-'geometrical' field, φ .
- ▶ KPMs $\langle M, g_1, g_2, \varphi \rangle$.
- ▶ Let g_1 be a fixed field; let g_2 co-vary dynamically with φ .
- ▶ For Belot, this theory is background independent, in spite of it having background structure g_1 .

Proposed fix: *Each* geometrical field must differ between *each* non-gauge-equivalent model.

Today

Preliminaries

General covariance

Diffeomorphism invariance

Absolute objects

Variational principles

Belot's approach

Fixed fields

Absolute fields

Conclusions

Fixed fields

Definition

***(Background independence, fixed fields)** A theory is background independent iff it has no formulation which features fixed fields.*

Fixed fields

- ▶ Doesn't this render **SR2** background independent?

Fixed fields

- ▶ Doesn't this render **SR2** background independent?
- ▶ Not if we say this is another formulation of **SR1**:

Fixed fields

- ▶ Doesn't this render **SR2** background independent?
- ▶ Not if we say this is another formulation of **SR1**:

Definition

(Equivalent theory formulations) Two theories, \mathcal{T}_1 and \mathcal{T}_2 , are equivalent formulations of the same theory when:

1. KPMs of \mathcal{T}_1 and \mathcal{T}_2 involve the same types of geometric object.
2. DPMs of \mathcal{T}_1 and \mathcal{T}_2 are isomorphic, up to classes of diffeomorphism-related models.

Fixed fields

- ▶ Doesn't this render **SR2** background independent?
- ▶ Not if we say this is another formulation of **SR1**:

Definition

(Equivalent theory formulations) Two theories, \mathcal{T}_1 and \mathcal{T}_2 , are equivalent formulations of the same theory when:

1. KPMs of \mathcal{T}_1 and \mathcal{T}_2 involve the same types of geometric object.
2. DPMs of \mathcal{T}_1 and \mathcal{T}_2 are isomorphic, up to classes of diffeomorphism-related models.

(One could, alternatively, appeal to one of the other notions of theoretical equivalence presented in (Weatherall 2019ab).)

Fixed fields

- ▶ Doesn't this render **SR2** background independent?
- ▶ Not if we say this is another formulation of **SR1**:

Definition

(Equivalent theory formulations) Two theories, \mathcal{T}_1 and \mathcal{T}_2 , are equivalent formulations of the same theory when:

1. *KPMs of \mathcal{T}_1 and \mathcal{T}_2 involve the same types of geometric object.*
2. *DPMs of \mathcal{T}_1 and \mathcal{T}_2 are isomorphic, up to classes of diffeomorphism-related models.*

(One could, alternatively, appeal to one of the other notions of theoretical equivalence presented in (Weatherall 2019ab).)

Potential problem: how do we know that the theory in question has no formulation featuring fixed fields?

Today

Preliminaries

General covariance

Diffeomorphism invariance

Absolute objects

Variational principles

Belot's approach

Fixed fields

Absolute fields

Conclusions

Absolute fields

Definition

(*Absolute field*) *A geometrical object specified in the KPMs of a theory, which is fixed (up to isomorphism) in all DPMs of that theory.*

Absolute fields

Definition

*(**Absolute field**) A geometrical object specified in the KPMs of a theory, which is fixed (up to isomorphism) in all DPMs of that theory.*

Definition

*(**Background independence, absolute fields**) A theory is background independent iff it has no absolute fields.*

Assessing absolute fields

- ▶ The idea would be that e.g. $\sqrt{-g}$ in GR doesn't qualify as an absolute field, because it's not directly written in the KPMs of the theory.

Assessing absolute fields

- ▶ The idea would be that e.g. $\sqrt{-g}$ in GR doesn't qualify as an absolute field, because it's not directly written in the KPMs of the theory.
- ▶ On the other hand, η_{ab} in **SR1**, and g_{ab} in **SR2**, do qualify as absolute fields.

Assessing absolute fields

- ▶ The idea would be that e.g. $\sqrt{-g}$ in GR doesn't qualify as an absolute field, because it's not directly written in the KPMs of the theory.
- ▶ On the other hand, η_{ab} in **SR1**, and g_{ab} in **SR2**, do qualify as absolute fields.
- ▶ Worry: Doesn't this make background independence too dependent upon (syntactic?) choices about the objects which constitute the KPMs of any given theory?

Today

Preliminaries

General covariance

Diffeomorphism invariance

Absolute objects

Variational principles

Belot's approach

Fixed fields

Absolute fields

Conclusions

Summary

- ▶ Since its inception, physicists and philosophers have debated what's special about general relativity.

Summary

- ▶ Since its inception, physicists and philosophers have debated what's special about general relativity.
- ▶ This began as a debate regarding 'substantive general covariance', and has evolved into a debate regarding 'background independence'.

Summary

- ▶ Since its inception, physicists and philosophers have debated what's special about general relativity.
- ▶ This began as a debate regarding 'substantive general covariance', and has evolved into a debate regarding 'background independence'.
- ▶ There are many interesting and illuminating ways of defining background independence—although arguably none without its problems.

Questions

1. Which of these proposals is the most promising?
2. Should we accept a plurality of definitions?
3. At what point does the task of trying to identify 'what's special about GR' become a futile one?

References



Gordon Belot, “Background-Independence”, *General Relativity and Gravitation* 43, pp. 2865-2884, 2011.



Michael Friedman, *Foundations of Spacetime Theories*, Princeton: Princeton University Press, 1983.



David Malament, *Topics in the Foundations of General Relativity and Newtonian Gravitation Theory*, Chicago: Chicago University Press, 2012.



Oliver Pooley, “Background Independence, Diffeomorphism Invariance, and the Meaning of Coordinates”, in D. Lehmkuhl, G. Schieman and E. Scholz (eds.), *Towards a Theory of Spacetime Theories*, Birkhäuser, 2017.



J. Brian Pitts, “Absolute Objects and Counterexamples: Jones-Geroch Dust, Torretti Constant Curvature, Tetrad-Spinor, and Scalar Density”, *Studies in History and Philosophy of Modern Physics* 37(2), pp. 347-371, 2006.



James Read, *Background Independence in Classical and Quantum Gravity*, book manuscript, 2020.



James Owen Weatherall, “Part 1: Theoretical Equivalence in Physics”, *Philosophy Compass* 14, e12592, 2019.



James Owen Weatherall, “Part 2: Theoretical Equivalence in Physics”, *Philosophy Compass* 14, e12591, 2019.