

# IPP-QM-2: Density operators and entanglement

James Read<sup>1</sup>

<sup>1</sup>Faculty of Philosophy, University of Oxford, UK, OX2 6GG

MT24

# The course

1. Basic quantum formalism
2. Density operators and entanglement
3. Decoherence
4. The measurement problem
5. Dynamical collapse theories
6. Bohmian mechanics
7. Everettian structure
8. Everettian probability
9. EPR and Bell's theorem
10. The Bell-CHSH inequalities and possible responses
11. Contextuality
12. The PBR theorem
13. Quantum logic
14. Pragmatism and QBism
15. Relational quantum mechanics
16. Wavefunction realism

# Today

Born rule recap

Density operators

Entanglement

More on density operators

# Today

Born rule recap

Density operators

Entanglement

More on density operators

# Motivating the Born rule

- ▶ Suppose we have  $\hat{1} = \sum_i |\phi_i\rangle \langle \phi_i|$ , where  $\{|\phi_i\rangle\}$  is the set of eigenstates of some physical quantity of interest.

# Motivating the Born rule

- ▶ Suppose we have  $\hat{1} = \sum_i |\phi_i\rangle \langle \phi_i|$ , where  $\{|\phi_i\rangle\}$  is the set of eigenstates of some physical quantity of interest.
- ▶ Associate each projector  $|\phi_i\rangle \langle \phi_i|$  with some outcome  $i$  of measurement, this outcome registering the response that the system being measured has the  $i$ th eigenvalue of the property in question.

# Motivating the Born rule

- ▶ Suppose we have  $\hat{1} = \sum_i |\phi_i\rangle \langle \phi_i|$ , where  $\{|\phi_i\rangle\}$  is the set of eigenstates of some physical quantity of interest.
- ▶ Associate each projector  $|\phi_i\rangle \langle \phi_i|$  with some outcome  $i$  of measurement, this outcome registering the response that the system being measured has the  $i$ th eigenvalue of the property in question.
- ▶ Let  $|\psi\rangle$  be the state of the system prior to measurement.

# Motivating the Born rule

- ▶ Notice that

$$\langle \psi | \hat{\mathbf{1}} | \psi \rangle = \langle \psi | \left( \sum_i |\phi_i\rangle \langle \phi_i| \right) | \psi \rangle = \mathbf{1},$$

since  $|\psi\rangle$  is normalised.

# Motivating the Born rule

- ▶ Notice that

$$\langle \psi | \hat{\mathbf{1}} | \psi \rangle = \langle \psi | \left( \sum_i |\phi_i\rangle \langle \phi_i| \right) | \psi \rangle = 1,$$

since  $|\psi\rangle$  is normalised.

- ▶ Writing this out, we have

$$\langle \psi | \phi_1 \rangle \langle \phi_1 | \psi \rangle + \dots + \langle \psi | \phi_n \rangle \langle \phi_n | \psi \rangle = 1,$$

i.e.

$$|\langle \psi | \phi_1 \rangle|^2 + \dots + |\langle \psi | \phi_n \rangle|^2 = 1.$$

# Motivating the Born rule

- ▶ Notice that

$$\langle \psi | \hat{1} | \psi \rangle = \langle \psi | \left( \sum_i |\phi_i\rangle \langle \phi_i| \right) | \psi \rangle = 1,$$

since  $|\psi\rangle$  is normalised.

- ▶ Writing this out, we have

$$\langle \psi | \phi_1 \rangle \langle \phi_1 | \psi \rangle + \dots + \langle \psi | \phi_n \rangle \langle \phi_n | \psi \rangle = 1,$$

i.e.

$$|\langle \psi | \phi_1 \rangle|^2 + \dots + |\langle \psi | \phi_n \rangle|^2 = 1.$$

- ▶ Hence if  $|\psi\rangle$  is normalised, then we can interpret the quantities  $|\langle \psi | \phi_i \rangle|^2$  as probabilities, as they are positive real numbers  $\leq 1$  which sum to 1.

# The Born rule

**Born rule:** For a measurement of some physical quantity represented by the self-adjoint operator  $\hat{A}$ , specified by a resolution of the identity in terms of the projectors  $\hat{P}(a_i)$  onto subspaces of  $\mathcal{H}$  corresponding to particular eigenvalues  $a_i$  of  $\hat{A}$ , the probability  $\text{Pr}(a_i)$  of getting outcome  $i$ , corresponding to  $\hat{P}(a_i)$  is given by  $\langle \psi | \hat{P}(a_i) | \psi \rangle$ , for a system in the state  $|\psi\rangle$ .

# The Born rule

**Born rule:** For a measurement of some physical quantity represented by the self-adjoint operator  $\hat{A}$ , specified by a resolution of the identity in terms of the projectors  $\hat{P}(a_i)$  onto subspaces of  $\mathcal{H}$  corresponding to particular eigenvalues  $a_i$  of  $\hat{A}$ , the probability  $\text{Pr}(a_i)$  of getting outcome  $i$ , corresponding to  $\hat{P}(a_i)$  is given by  $\langle \psi | \hat{P}(a_i) | \psi \rangle$ , for a system in the state  $|\psi\rangle$ .

Note that at this point this is an *operational prescription!* Also, note that the projection postulate—closely associated with ‘wavefunction collapse’—is an optional extra here...

# Today

Born rule recap

**Density operators**

Entanglement

More on density operators

# Trace of an operator

- ▶ Let's introduce the *trace* of an operator  $\hat{A}$ , denoted  $\text{Tr}(\hat{A})$ .

# Trace of an operator

- ▶ Let's introduce the *trace* of an operator  $\hat{A}$ , denoted  $\text{Tr}(\hat{A})$ .
- ▶ This is defined as the sum of the diagonal elements of  $\hat{A}$  (in any basis).

# Trace of an operator

- ▶ Let's introduce the *trace* of an operator  $\hat{A}$ , denoted  $\text{Tr}(\hat{A})$ .
- ▶ This is defined as the sum of the diagonal elements of  $\hat{A}$  (in any basis).
- ▶ So, if we choose a basis  $\{|\psi_i\rangle\}$  in the Hilbert space, we have

$$\text{Tr}(\hat{A}) = \sum_i \langle \psi_i | \hat{A} | \psi_i \rangle .$$

# Trace of an operator

- ▶ Let's introduce the *trace* of an operator  $\hat{A}$ , denoted  $\text{Tr}(\hat{A})$ .
- ▶ This is defined as the sum of the diagonal elements of  $\hat{A}$  (in any basis).
- ▶ So, if we choose a basis  $\{|\psi_i\rangle\}$  in the Hilbert space, we have

$$\text{Tr}(\hat{A}) = \sum_i \langle \psi_i | \hat{A} | \psi_i \rangle .$$

**Exercise:** Check that this definition is indeed independent of the basis chosen.

# Trace of an operator

- ▶ Let's introduce the *trace* of an operator  $\hat{A}$ , denoted  $\text{Tr}(\hat{A})$ .
- ▶ This is defined as the sum of the diagonal elements of  $\hat{A}$  (in any basis).
- ▶ So, if we choose a basis  $\{|\psi_i\rangle\}$  in the Hilbert space, we have

$$\text{Tr}(\hat{A}) = \sum_i \langle \psi_i | \hat{A} | \psi_i \rangle .$$

**Exercise:** Check that this definition is indeed independent of the basis chosen.

- ▶ Two other properties of the trace:
  1. Linearity:  $\text{Tr}(\alpha\hat{A} + \beta\hat{B}) = \alpha\text{Tr}(\hat{A}) + \beta\text{Tr}(\hat{B})$ .
  2. Cyclicity:  $\text{Tr}(\hat{A}\hat{B}) = \text{Tr}(\hat{B}\hat{A})$ .

# Density operators

- ▶ Define a *density operator* (or *density matrix*, if one is working in a particular basis) to be a self-adjoint positive operator  $\hat{\rho}$  with  $\text{Tr}(\hat{\rho}) = 1$ . (An operator's being positive means that its eigenvalues are all positive.)

# Density operators

- ▶ Define a *density operator* (or *density matrix*, if one is working in a particular basis) to be a self-adjoint positive operator  $\hat{\rho}$  with  $\text{Tr}(\hat{\rho}) = 1$ . (An operator's being positive means that its eigenvalues are all positive.)
- ▶ A special case is  $\hat{\rho} = |\psi\rangle\langle\psi|$  for some vector state  $|\psi\rangle$ .

# Density operators

- ▶ Define a *density operator* (or *density matrix*, if one is working in a particular basis) to be a self-adjoint positive operator  $\hat{\rho}$  with  $\text{Tr}(\hat{\rho}) = 1$ . (An operator's being positive means that its eigenvalues are all positive.)
- ▶ A special case is  $\hat{\rho} = |\psi\rangle\langle\psi|$  for some vector state  $|\psi\rangle$ .
- ▶ In this way, any vector state can be represented by a density operator.

# Pure and mixed states

- ▶ A system is in a *pure state* when its density operator is given by the projector  $|\psi\rangle\langle\psi|$  onto the 1D subspace spanned by the vector state  $|\psi\rangle$ .

# Pure and mixed states

- ▶ A system is in a *pure state* when its density operator is given by the projector  $|\psi\rangle\langle\psi|$  onto the 1D subspace spanned by the vector state  $|\psi\rangle$ .
- ▶ A system is said to be in a *mixed state* when the density operator associated with it is *not* given by a 1D projector; that is, when it is given by some weighted sum of projectors  $\hat{P}_i$  (in general these projectors needn't be onto orthogonal subspaces),

$$\hat{\rho} = \sum_i \lambda_i \hat{P}_i,$$

where more than one of the  $\lambda_i$  is non-zero.

# Pure and mixed states

- ▶ A system is in a *pure state* when its density operator is given by the projector  $|\psi\rangle\langle\psi|$  onto the 1D subspace spanned by the vector state  $|\psi\rangle$ .
- ▶ A system is said to be in a *mixed state* when the density operator associated with it is *not* given by a 1D projector; that is, when it is given by some weighted sum of projectors  $\hat{P}_i$  (in general these projectors needn't be onto orthogonal subspaces),

$$\hat{\rho} = \sum_i \lambda_i \hat{P}_i,$$

where more than one of the  $\lambda_i$  is non-zero.

- ▶ Note that whenever we write a vector state  $|\psi\rangle$  in the form  $|\psi\rangle\langle\psi|$ , the (redundant) overall phase cancels out, so in a sense this representation is *more accurate* physically than the one given by vectors in Hilbert space.

# Pure and mixed states

- ▶ A system is in a *pure state* when its density operator is given by the projector  $|\psi\rangle\langle\psi|$  onto the 1D subspace spanned by the vector state  $|\psi\rangle$ .
- ▶ A system is said to be in a *mixed state* when the density operator associated with it is *not* given by a 1D projector; that is, when it is given by some weighted sum of projectors  $\hat{P}_i$  (in general these projectors needn't be onto orthogonal subspaces),

$$\hat{\rho} = \sum_i \lambda_i \hat{P}_i,$$

where more than one of the  $\lambda_i$  is non-zero.

- ▶ Note that whenever we write a vector state  $|\psi\rangle$  in the form  $|\psi\rangle\langle\psi|$ , the (redundant) overall phase cancels out, so in a sense this representation is *more accurate* physically than the one given by vectors in Hilbert space.
- ▶ **Exercise:** Show that  $\hat{\rho}$  is pure iff  $\hat{\rho}^2 = \hat{\rho}$ , iff  $\det(\hat{\rho}) = 0$ .

# Proper and improper mixtures

- ▶ When a system is *really* in some pure state, but we don't know which one, then it is said to be *properly mixed*, or part of a *proper mixture*.

# Proper and improper mixtures

- ▶ When a system is *really* in some pure state, but we don't know which one, then it is said to be *properly mixed*, or part of a *proper mixture*.
- ▶ If a range of options is some set of (potentially non-orthogonal) states  $\{|\psi_i\rangle\}$ , and the probability of being in the  $i$ th state is  $\text{Pr}(i)$ , then the density operator describing the mixed state is

$$\hat{\rho} = \sum_i \text{Pr}(i) |\psi_i\rangle \langle \psi_i|;$$

since the system really is in one of the  $|\psi_i\rangle$ , we say that the density operator in this case has an *ignorance interpretation*.

# Proper and improper mixtures

- ▶ When a system is *really* in some pure state, but we don't know which one, then it is said to be *properly mixed*, or part of a *proper mixture*.
- ▶ If a range of options is some set of (potentially non-orthogonal) states  $\{|\psi_i\rangle\}$ , and the probability of being in the  $i$ th state is  $\text{Pr}(i)$ , then the density operator describing the mixed state is

$$\hat{\rho} = \sum_i \text{Pr}(i) |\psi_i\rangle \langle \psi_i|;$$

since the system really is in one of the  $|\psi_i\rangle$ , we say that the density operator in this case has an *ignorance interpretation*.

- ▶ We'll come in due course to examples where systems do not have pure states at all, only mixed ones (i.e., a non-pure density operator is the only way in which they can be described) and where an ignorance interpretation is not possible. These systems are called *improper mixtures*.

# Trace form of the Born rule

Density matrices and the trace operation give us a nice way of writing the Born rule:

# Trace form of the Born rule

Density matrices and the trace operation give us a nice way of writing the Born rule:

**Born rule:** For a measurement specified by a resolution of the identity  $\hat{1} = \sum_i \hat{P}_i$ , the probability of getting outcome  $i$ , associated with positive operator  $\hat{P}_i$ , for a system with density operator  $\hat{\rho}$ , is

$$\Pr(i) = \text{Tr}(\hat{\rho}\hat{P}_i).$$

# Trace form of the Born rule

Density matrices and the trace operation give us a nice way of writing the Born rule:

**Born rule:** For a measurement specified by a resolution of the identity  $\hat{1} = \sum_i \hat{P}_i$ , the probability of getting outcome  $i$ , associated with positive operator  $\hat{P}_i$ , for a system with density operator  $\hat{\rho}$ , is

$$\text{Pr}(i) = \text{Tr}(\hat{\rho}\hat{P}_i).$$

It's easy to see that this recovers the previous version of the Born rule. Consider some basis  $\{|\phi_k\rangle\}$ , let  $\hat{P}_i$  be the projector  $|\phi_i\rangle\langle\phi_i|$ , and let  $\hat{\rho}$  be associated with a system in a pure state  $|\psi\rangle$ , so that  $\hat{\rho} = |\psi\rangle\langle\psi|$ . Then:

$$\begin{aligned}\text{Tr}(\hat{\rho}\hat{P}_i) &= \sum_k \langle\phi_k|\hat{\rho}|\phi_i\rangle\langle\phi_i|\phi_k\rangle = \sum_k \langle\phi_k|\hat{\rho}|\phi_i\rangle\delta_{ij} \\ &= \langle\phi_i|\psi\rangle\langle\psi|\phi_i\rangle = |\langle\phi_i|\psi\rangle|^2\end{aligned}$$

# Today

Born rule recap

Density operators

**Entanglement**

More on density operators

# Product systems

- ▶ Suppose we have two quantum systems  $A$  and  $B$  with respective Hilbert spaces  $\mathcal{H}_A$  and  $\mathcal{H}_B$ . We want to consider how to combine these descriptions into a single description for the combined system,  $A \times B$ .

# Product systems

- ▶ Suppose we have two quantum systems  $A$  and  $B$  with respective Hilbert spaces  $\mathcal{H}_A$  and  $\mathcal{H}_B$ . We want to consider how to combine these descriptions into a single description for the combined system,  $A \times B$ .
- ▶ That combined system is still a quantum system, so we represent it with the ‘tensor product Hilbert space’,  $\mathcal{H}_A \otimes \mathcal{H}_B$ .

# Product systems

- ▶ Suppose we have two quantum systems  $A$  and  $B$  with respective Hilbert spaces  $\mathcal{H}_A$  and  $\mathcal{H}_B$ . We want to consider how to combine these descriptions into a single description for the combined system,  $A \times B$ .
- ▶ That combined system is still a quantum system, so we represent it with the ‘tensor product Hilbert space’,  $\mathcal{H}_A \otimes \mathcal{H}_B$ .
- ▶ We want to understand how  $\mathcal{H}_A \otimes \mathcal{H}_B$  is built up from  $\mathcal{H}_A$  and  $\mathcal{H}_B$ .

# Product systems

- ▶ Recall that a basis of states  $\{|m_i\rangle\}$  for  $\mathcal{H}_A$  can be thought of as defining a maximally specific measurement  $M$  on  $A$ : each  $|m_i\rangle$  always gives some particular result  $m_i$  on measuring  $M$ .

# Product systems

- ▶ Recall that a basis of states  $\{|m_i\rangle\}$  for  $\mathcal{H}_A$  can be thought of as defining a maximally specific measurement  $M$  on  $A$ : each  $|m_i\rangle$  always gives some particular result  $m_i$  on measuring  $M$ .
- ▶ Suppose we pick some such  $M$ , and suppose similarly that we pick an  $N$  for  $B$ , defined by another basis  $\{|n_i\rangle\}$  for  $\mathcal{H}_B$ .

# Product systems

- ▶ Recall that a basis of states  $\{|m_i\rangle\}$  for  $\mathcal{H}_A$  can be thought of as defining a maximally specific measurement  $M$  on  $A$ : each  $|m_i\rangle$  always gives some particular result  $m_i$  on measuring  $M$ .
- ▶ Suppose we pick some such  $M$ , and suppose similarly that we pick an  $N$  for  $B$ , defined by another basis  $\{|n_j\rangle\}$  for  $\mathcal{H}_B$ .
- ▶ We now make the assumption that the pairs  $|m_i\rangle \otimes |n_j\rangle$  of vectors from these two bases form a basis for  $\mathcal{H}_A \otimes \mathcal{H}_B$ :  $|m_i\rangle \otimes |n_j\rangle$  gives  $m_i$  as a result of measuring  $M$  on  $A$  and  $n_j$  as a result of measuring  $N$  on  $B$ .

# Product systems

- ▶ Recall that a basis of states  $\{|m_i\rangle\}$  for  $\mathcal{H}_A$  can be thought of as defining a maximally specific measurement  $M$  on  $A$ : each  $|m_i\rangle$  always gives some particular result  $m_i$  on measuring  $M$ .
- ▶ Suppose we pick some such  $M$ , and suppose similarly that we pick an  $N$  for  $B$ , defined by another basis  $\{|n_j\rangle\}$  for  $\mathcal{H}_B$ .
- ▶ We now make the assumption that the pairs  $|m_i\rangle \otimes |n_j\rangle$  of vectors from these two bases form a basis for  $\mathcal{H}_A \otimes \mathcal{H}_B$ :  $|m_i\rangle \otimes |n_j\rangle$  gives  $m_i$  as a result of measuring  $M$  on  $A$  and  $n_j$  as a result of measuring  $N$  on  $B$ .
- ▶ An arbitrary state of  $\mathcal{H}_A \otimes \mathcal{H}_B$  can now be built up from these basis vectors—a completely general expression is

$$|\psi\rangle = \sum_i \sum_j \alpha_{ij} |m_i\rangle \otimes |n_j\rangle.$$

# Specifying the state of a combined system

- ▶ In classical physics, we can always give the state of a combined system just by giving the states of the component systems.

# Specifying the state of a combined system

- ▶ In classical physics, we can always give the state of a combined system just by giving the states of the component systems.
- ▶ But is the same true in quantum mechanics?

# Tensor product of states

- ▶ Suppose  $|\chi\rangle$  and  $|\phi\rangle$  are states of  $A$  and  $B$  respectively, given by

$$|\chi\rangle = \sum_i c_i |m_i\rangle, \quad |\phi\rangle = \sum_j d_j |n_j\rangle.$$

# Tensor product of states

- ▶ Suppose  $|\chi\rangle$  and  $|\phi\rangle$  are states of  $A$  and  $B$  respectively, given by

$$|\chi\rangle = \sum_i c_i |m_i\rangle, \quad |\phi\rangle = \sum_j d_j |n_j\rangle.$$

- ▶ Then we can define their tensor product  $|\chi\rangle \otimes |\phi\rangle$  by linearity:

$$\begin{aligned} |\chi\rangle \otimes |\phi\rangle &= \left( \sum_i c_i |m_i\rangle \right) \otimes \left( \sum_j d_j |n_j\rangle \right) \\ &=: \sum_i \sum_j c_i d_j |m_i\rangle \otimes |n_j\rangle \end{aligned}$$

# Entanglement

- ▶ But not all states can be written in this way! Compare  $\sum_i \sum_j c_i d_j |m_i\rangle \otimes |n_j\rangle$  with the more general expression  $\sum_i \sum_j \alpha_{ij} |m_i\rangle \otimes |n_j\rangle$  which we saw before.

# Entanglement

- ▶ But not all states can be written in this way! Compare  $\sum_i \sum_j c_i d_j |m_i\rangle \otimes |n_j\rangle$  with the more general expression  $\sum_i \sum_j \alpha_{ij} |m_i\rangle \otimes |n_j\rangle$  which we saw before.
- ▶ States which cannot be written in the product form are called *entangled*.

# Entanglement

- ▶ But not all states can be written in this way! Compare  $\sum_i \sum_j c_i d_j |m_i\rangle \otimes |n_j\rangle$  with the more general expression  $\sum_i \sum_j \alpha_{ij} |m_i\rangle \otimes |n_j\rangle$  which we saw before.
- ▶ States which cannot be written in the product form are called *entangled*.
- ▶ Entangled states can be found in even the smallest systems.

# Entanglement

- ▶ But not all states can be written in this way! Compare  $\sum_i \sum_j c_i d_j |m_i\rangle \otimes |n_j\rangle$  with the more general expression  $\sum_i \sum_j \alpha_{ij} |m_i\rangle \otimes |n_j\rangle$  which we saw before.
- ▶ States which cannot be written in the product form are called *entangled*.
- ▶ Entangled states can be found in even the smallest systems.
- ▶ For example, if we have two spin-half particles each of which has z-spin eigenstates  $|+\rangle$ ,  $|-\rangle$ , then the ‘singlet’ state

$$\frac{1}{\sqrt{2}} (|+\rangle \otimes |-\rangle - |-\rangle \otimes |+\rangle)$$

is entangled.

# Entanglement

- ▶ But not all states can be written in this way! Compare  $\sum_i \sum_j c_i d_j |m_i\rangle \otimes |n_j\rangle$  with the more general expression  $\sum_i \sum_j \alpha_{ij} |m_i\rangle \otimes |n_j\rangle$  which we saw before.
- ▶ States which cannot be written in the product form are called *entangled*.
- ▶ Entangled states can be found in even the smallest systems.
- ▶ For example, if we have two spin-half particles each of which has z-spin eigenstates  $|+\rangle$ ,  $|-\rangle$ , then the ‘singlet’ state

$$\frac{1}{\sqrt{2}} (|+\rangle \otimes |-\rangle - |-\rangle \otimes |+\rangle)$$

is entangled. (**Exercise:** Convince yourself of this.)

# Entanglement

- ▶ But not all states can be written in this way! Compare  $\sum_i \sum_j c_i d_j |m_i\rangle \otimes |n_j\rangle$  with the more general expression  $\sum_i \sum_j \alpha_{ij} |m_i\rangle \otimes |n_j\rangle$  which we saw before.

- ▶ States which cannot be written in the product form are called *entangled*.

- ▶ Entangled states can be found in even the smallest systems.

- ▶ For example, if we have two spin-half particles each of which has z-spin eigenstates  $|+\rangle$ ,  $|-\rangle$ , then the ‘singlet’ state

$$\frac{1}{\sqrt{2}} (|+\rangle \otimes |-\rangle - |-\rangle \otimes |+\rangle)$$

is entangled. (**Exercise:** Convince yourself of this.)

- ▶ What’s the spin state of system 1 or system 2 in the above joint ‘singlet’ states? Answer: *undefined!*

# Intermezzo: tensor product of operators

To get a grip on the meaning of ' $\otimes$ ':

# Intermezzo: tensor product of operators

To get a grip on the meaning of ' $\otimes$ ':

- ▶ If  $\hat{A}$  acts on  $\mathcal{H}_1$  and  $\hat{B}$  acts on  $\mathcal{H}_2$  ...

# Intermezzo: tensor product of operators

To get a grip on the meaning of ' $\otimes$ ':

- ▶ If  $\hat{A}$  acts on  $\mathcal{H}_1$  and  $\hat{B}$  acts on  $\mathcal{H}_2$  ...
- ▶ ... then  $\hat{A} \otimes \hat{B}$  acts (linearly) on  $\mathcal{H}_1 \otimes \mathcal{H}_2$ :

# Intermezzo: tensor product of operators

To get a grip on the meaning of ' $\otimes$ ':

- ▶ If  $\hat{A}$  acts on  $\mathcal{H}_1$  and  $\hat{B}$  acts on  $\mathcal{H}_2$  ...
- ▶ ... then  $\hat{A} \otimes \hat{B}$  acts (linearly) on  $\mathcal{H}_1 \otimes \mathcal{H}_2$ :

$$\begin{aligned}(\hat{A} \otimes \hat{B}) |\psi\rangle_{12} &= (\hat{A} \otimes \hat{B}) \left( \sum_{ij} \alpha_{ij} |\phi_i\rangle_1 \otimes |\chi_j\rangle_2 \right) \\ &= \sum_{ij} \alpha_{ij} \hat{A} |\phi_i\rangle_1 \otimes \hat{B} |\chi_j\rangle_2.\end{aligned}$$

# Intermezzo: tensor product of operators

To get a grip on the meaning of ' $\otimes$ ':

- ▶ If  $\hat{A}$  acts on  $\mathcal{H}_1$  and  $\hat{B}$  acts on  $\mathcal{H}_2$  ...
- ▶ ... then  $\hat{A} \otimes \hat{B}$  acts (linearly) on  $\mathcal{H}_1 \otimes \mathcal{H}_2$ :

$$\begin{aligned}(\hat{A} \otimes \hat{B}) |\psi\rangle_{12} &= (\hat{A} \otimes \hat{B}) \left( \sum_{ij} \alpha_{ij} |\phi_i\rangle_1 \otimes |\chi_j\rangle_2 \right) \\ &= \sum_{ij} \alpha_{ij} \hat{A} |\phi_i\rangle_1 \otimes \hat{B} |\chi_j\rangle_2.\end{aligned}$$

Note: It is always possible to write any linear operator  $\hat{O}_{12}$  acting on  $\mathcal{H}_1 \otimes \mathcal{H}_2$  in the form

$$\hat{O}_{12} = \sum_{kl} c_{kl} \hat{A}_k \otimes \hat{B}_l.$$

# A qualitatively new phenomenon

- ▶ If there's no way to write the state  $|\Psi\rangle_{12}$  as a product of a state in  $\mathcal{H}_1$  and a state in  $\mathcal{H}_2$  then we have a *qualitatively new phenomenon*.

# A qualitatively new phenomenon

- ▶ If there's no way to write the state  $|\Psi\rangle_{12}$  as a product of a state in  $\mathcal{H}_1$  and a state in  $\mathcal{H}_2$  then we have a *qualitatively new phenomenon*.
- ▶ *The individual systems **do not have states of their own**—only the global system does.*

# A qualitatively new phenomenon

- ▶ If there's no way to write the state  $|\Psi\rangle_{12}$  as a product of a state in  $\mathcal{H}_1$  and a state in  $\mathcal{H}_2$  then we have a *qualitatively new phenomenon*.
- ▶ *The individual systems **do not have states of their own**—only the global system does.*

*Entanglement shows that there are global properties of a joint system which are not reducible to properties of subsystems!*

# A qualitatively new phenomenon

- ▶ If there's no way to write the state  $|\Psi\rangle_{12}$  as a product of a state in  $\mathcal{H}_1$  and a state in  $\mathcal{H}_2$  then we have a *qualitatively new phenomenon*.
- ▶ *The individual systems **do not have states of their own**—only the global system does.*

*Entanglement shows that there are global properties of a joint system which are not reducible to properties of subsystems!*

As we'll see in Lecture 9, this is closely related to 'non-separability'.

# Entanglement and mixed states

- ▶ Alice and Bob are going to do experiments involving measuring the spin of a single electron.

# Entanglement and mixed states

- ▶ Alice and Bob are going to do experiments involving measuring the spin of a single electron.
- ▶ Alice asks Bob to prepare an electron in an x-spin superposition state (specifically, the state  $|\psi\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle + |\downarrow_x\rangle)$ ).

# Entanglement and mixed states

- ▶ Alice and Bob are going to do experiments involving measuring the spin of a single electron.
- ▶ Alice asks Bob to prepare an electron in an x-spin superposition state (specifically, the state  $|\psi\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle + |\downarrow_x\rangle)$ ).
- ▶ Bob is lazy, and just prepared the electron in a probabilistic state (so it has x-spin or down definitely, but he makes sure there's a 50/50 chance of each).

# Entanglement and mixed states

- ▶ Alice and Bob are going to do experiments involving measuring the spin of a single electron.
- ▶ Alice asks Bob to prepare an electron in an x-spin superposition state (specifically, the state  $|\psi\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle + |\downarrow_x\rangle)$ ).
- ▶ Bob is lazy, and just prepared the electron in a probabilistic state (so it has x-spin up or down definitely, but he makes sure there's a 50/50 chance of each).
- ▶ Alice knows Bob well, and is suspicious. She doesn't have an interferometer to hand. What could she do to check up on Bob?

## The example continued

- ▶ Suppose Alice can perform linear transformations on the state that effectively rotate it: spin up states in the x direction become spin up states in the z direction (which are superpositions of x-spin states), etc.:

$$|\uparrow_x\rangle \rightarrow |\uparrow_z\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle + |\downarrow_x\rangle),$$

$$|\downarrow_x\rangle \rightarrow |\downarrow_z\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle - |\downarrow_x\rangle).$$

## The example continued

- ▶ Suppose Alice can perform linear transformations on the state that effectively rotate it: spin up states in the x direction become spin up states in the z direction (which are superpositions of x-spin states), etc.:

$$|\uparrow_x\rangle \rightarrow |\uparrow_z\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle + |\downarrow_x\rangle),$$

$$|\downarrow_x\rangle \rightarrow |\downarrow_z\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle - |\downarrow_x\rangle).$$

- ▶ What happens if Alice performs this transition on her desired superposition state? She gets a definite  $|\uparrow_x\rangle$  state. (Do the calculation!)

## The example continued

- ▶ Suppose Alice can perform linear transformations on the state that effectively rotate it: spin up states in the x direction become spin up states in the z direction (which are superpositions of x-spin states), etc.:

$$|\uparrow_x\rangle \rightarrow |\uparrow_z\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle + |\downarrow_x\rangle),$$
$$|\downarrow_x\rangle \rightarrow |\downarrow_z\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle - |\downarrow_x\rangle).$$

- ▶ What happens if Alice performs this transition on her desired superposition state? She gets a definite  $|\uparrow_x\rangle$  state. (Do the calculation!)
- ▶ What happens when she performs this transformation on one of Bob's fake superpositions/really definite states? She gets a superposition state in the x basis.

# The example continued

- ▶ So Alice can use the relationship between bases to uncover Bob's play!

# The example continued

- ▶ So Alice can use the relationship between bases to uncover Bob's ploy!
- ▶ If she performs her transformation but continues to get a 50/50 mix of up and down results, she'll know they were never really superposition states!

# The example continued

- ▶ So Alice can use the relationship between bases to uncover Bob's ploy!
- ▶ If she performs her transformation but continues to get a 50/50 mix of up and down results, she'll know they were never really superposition states!
- ▶ How could Bob circumvent this problem?

# The example continued

- ▶ Bob could agree in advance with Alice to entangle the electrons with another electron in the singlet state:

$$|\psi\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle_1 |\downarrow_x\rangle_2 - |\downarrow_x\rangle_1 |\uparrow_x\rangle_2)$$

## The example continued

- ▶ Bob could agree in advance with Alice to entangle the electrons with another electron in the singlet state:

$$|\psi\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle_1 |\downarrow_x\rangle_2 - |\downarrow_x\rangle_1 |\uparrow_x\rangle_2)$$

- ▶ But recall that the singlet state is spherically symmetric:

$$|\psi\rangle = \frac{1}{\sqrt{2}} (|\uparrow_z\rangle_1 |\downarrow_z\rangle_2 - |\downarrow_z\rangle_1 |\uparrow_z\rangle_2)$$

## The example continued

- ▶ Bob could agree in advance with Alice to entangle the electrons with another electron in the singlet state:

$$|\psi\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle_1 |\downarrow_x\rangle_2 - |\downarrow_x\rangle_1 |\uparrow_x\rangle_2)$$

- ▶ But recall that the singlet state is spherically symmetric:

$$|\psi\rangle = \frac{1}{\sqrt{2}} (|\uparrow_z\rangle_1 |\downarrow_z\rangle_2 - |\downarrow_z\rangle_1 |\uparrow_z\rangle_2)$$

- ▶ Now if Alice tries to perform her rotations, she won't get a definite spin state, so she won't expect to see a difference.

# Moral

*Entanglement can make probabilistic mixtures indistinguishable from superpositions (if we're not doing interference experiments)!*

# Today

Born rule recap

Density operators

Entanglement

**More on density operators**

# Back to density operators

A density matrix with vanishing off-diagonals can represent:

# Back to density operators

A density matrix with vanishing off-diagonals can represent:

1. A true probabilistic mixture of states (a 'proper mixture'), or

# Back to density operators

A density matrix with vanishing off-diagonals can represent:

1. A true probabilistic mixture of states (a 'proper mixture'), or
2. A superposition of states which are not interfering (an 'improper mixture').

# Back to density operators

A density matrix with vanishing off-diagonals can represent:

1. A true probabilistic mixture of states (a ‘proper mixture’), or
2. A superposition of states which are not interfering (an ‘improper mixture’).

We’ve seen that if the quantum state of some (sub)system can be written only using density operators (not vectors), it is ‘impure’, otherwise it is ‘pure’.

# Entanglement and density operators

When we have entanglement, there will be no vector state for subsystems, but it turns out that one *can* assign them *density operators*.

# Entanglement and density operators

When we have entanglement, there will be no vector state for subsystems, but it turns out that one *can* assign them *density operators*.

- ▶ These are called the *reduced states* (*reduced density operators*) of a subsystem.

# Entanglement and density operators

When we have entanglement, there will be no vector state for subsystems, but it turns out that one *can* assign them *density operators*.

- ▶ These are called the *reduced states* (*reduced density operators*) of a subsystem.
- ▶ When we have entanglement these reduced states will be *mixed* (by definition).

# Entanglement and density operators

When we have entanglement, there will be no vector state for subsystems, but it turns out that one *can* assign them *density operators*.

- ▶ These are called the *reduced states (reduced density operators)* of a subsystem.
- ▶ When we have entanglement these reduced states will be *mixed* (by definition).
- ▶ But these are not mixtures which can be given an *ignorance interpretation*: we cannot think of these mixtures as telling us that there is some underlying pure state of which we are ignorant.

# Reduced states

- ▶ Take a composite of two systems, with Hilbert spaces  $\mathcal{H}_1$  and  $\mathcal{H}_2$ , say in a state

$$|\Psi\rangle = \sum_i \alpha_{ij} |\psi_i\rangle \otimes |\phi_j\rangle.$$

# Reduced states

- ▶ Take a composite of two systems, with Hilbert spaces  $\mathcal{H}_1$  and  $\mathcal{H}_2$ , say in a state

$$|\Psi\rangle = \sum_i \alpha_{ij} |\psi_i\rangle \otimes |\phi_j\rangle.$$

- ▶ This state, as we know, defines expectation values for all the observables of the composite system  $\mathcal{H}_1 \otimes \mathcal{H}_2$ .

# Reduced states

- ▶ Take a composite of two systems, with Hilbert spaces  $\mathcal{H}_1$  and  $\mathcal{H}_2$ , say in a state

$$|\Psi\rangle = \sum_i \alpha_{ij} |\psi_i\rangle \otimes |\phi_j\rangle.$$

- ▶ This state, as we know, defines expectation values for all the observables of the composite system  $\mathcal{H}_1 \otimes \mathcal{H}_2$ .
- ▶ Special cases of these observables are ones of the form  $\hat{A} \otimes \hat{1}$  and  $\hat{1} \otimes \hat{B}$ , where  $\hat{A}$  is an observable on  $\mathcal{H}_1$  and  $\hat{B}$  is an observable on  $\mathcal{H}_2$ , and  $\hat{1}$  denotes the identity operator on either space.

# Reduced states

- ▶ Take a composite of two systems, with Hilbert spaces  $\mathcal{H}_1$  and  $\mathcal{H}_2$ , say in a state

$$|\Psi\rangle = \sum_i \alpha_{ij} |\psi_i\rangle \otimes |\phi_j\rangle.$$

- ▶ This state, as we know, defines expectation values for all the observables of the composite system  $\mathcal{H}_1 \otimes \mathcal{H}_2$ .
- ▶ Special cases of these observables are ones of the form  $\hat{A} \otimes \hat{1}$  and  $\hat{1} \otimes \hat{B}$ , where  $\hat{A}$  is an observable on  $\mathcal{H}_1$  and  $\hat{B}$  is an observable on  $\mathcal{H}_2$ , and  $\hat{1}$  denotes the identity operator on either space.
- ▶ The state  $|\Psi\rangle$  on  $\mathcal{H}_1 \otimes \mathcal{H}_2$  defines two (possibly impure) states on  $\mathcal{H}_1$  and  $\mathcal{H}_2$ , respectively.

# The mathematics of reduced states

- ▶ We shall now discuss how to calculate the (density operators corresponding to) reduced states. This is done by means of the *partial trace*.

# The mathematics of reduced states

- ▶ We shall now discuss how to calculate the (density operators corresponding to) reduced states. This is done by means of the *partial trace*.
- ▶ Recall that the trace of an operator is given by

$$\text{Tr}(\hat{A}) = \sum_i \langle \psi_i | \hat{A} | \psi_i \rangle ,$$

where  $\{|\psi_i\rangle\}$  is any orthonormal basis in the Hilbert space on which  $\hat{A}$  acts.

# The mathematics of reduced states

- ▶ We shall now discuss how to calculate the (density operators corresponding to) reduced states. This is done by means of the *partial trace*.
- ▶ Recall that the trace of an operator is given by

$$\text{Tr}(\hat{A}) = \sum_i \langle \psi_i | \hat{A} | \psi_i \rangle ,$$

where  $\{|\psi_i\rangle\}$  is any orthonormal basis in the Hilbert space on which  $\hat{A}$  acts.

- ▶ Applied to the space  $\mathcal{H}_1 \otimes \mathcal{H}_2$ , the basis  $\{|\phi_i\rangle \otimes |\psi_j\rangle\}$  and the pure state  $|\Psi\rangle$ , this yields for any observable  $\hat{C}$  on  $\mathcal{H}_1 \otimes \mathcal{H}_2$ :

$$\text{Tr} \left( |\Psi\rangle \langle \Psi| \hat{C} \right) = \sum_{i,j} \langle \phi_i | \otimes \langle \psi_j | \left[ |\Psi\rangle \langle \Psi| \hat{C} \right] | \phi_i\rangle \otimes | \psi_j\rangle$$

## Obtaining reduced states

If, say,  $\hat{C}$  has the form  $\hat{A} \otimes \hat{1}$ , then this simplifies to

$$\begin{aligned}\text{Tr}(|\Psi\rangle\langle\Psi|\hat{A}\otimes\hat{1}) &= \sum_{i,j} \langle\phi_i|\otimes\langle\psi_j| \left[ |\Psi\rangle\langle\Psi|\hat{A}\otimes\hat{1} \right] |\phi_i\rangle\otimes|\psi_j\rangle \\ &= \sum_i \langle\phi_i| \left[ \sum_j \langle\psi_j|\Psi\rangle\langle\Psi|\psi_j\rangle \right] \hat{A}|\phi_j\rangle\end{aligned}$$

## Obtaining reduced states

If, say,  $\hat{C}$  has the form  $\hat{A} \otimes \hat{1}$ , then this simplifies to

$$\begin{aligned}\text{Tr}(|\Psi\rangle\langle\Psi|\hat{A} \otimes \hat{1}) &= \sum_{i,j} \langle\phi_i|\otimes\langle\psi_j| \left[ |\Psi\rangle\langle\Psi|\hat{A} \otimes \hat{1} \right] |\phi_i\rangle\otimes|\psi_j\rangle \\ &= \sum_i \langle\phi_i| \left[ \sum_j \langle\psi_j|\Psi\rangle\langle\Psi|\psi_j\rangle \right] \hat{A}|\phi_i\rangle\end{aligned}$$

And defining

$$\hat{\rho}_1 := \sum_j \langle\psi_j|\Psi\rangle\langle\Psi|\psi_j\rangle,$$

we can rewrite this as

$$\text{Tr}(|\Psi\rangle\langle\Psi|\hat{A} \otimes \hat{1}) = \text{Tr}(\hat{\rho}_1 \hat{A}),$$

where the trace on the RHS is the trace in  $\mathcal{H}_1$ .

# Obtaining reduced states

- ▶ We have thus found the density operator describing the *reduced state* of  $\mathcal{H}_1$ .

# Obtaining reduced states

- ▶ We have thus found the density operator describing the *reduced state* of  $\mathcal{H}_1$ .
- ▶ This yields the expectation values for any observable on  $\mathcal{H}_1$  when the state of the total system is  $|\Psi\rangle$ —for recall the affinity between our previous expression and how we stated the Born rule.

# Obtaining reduced states

- ▶ We have thus found the density operator describing the *reduced state* of  $\mathcal{H}_1$ .
- ▶ This yields the expectation values for any observable on  $\mathcal{H}_1$  when the state of the total system is  $|\Psi\rangle$ —for recall the affinity between our previous expression and how we stated the Born rule.
- ▶ This operation is known as the *partial trace* over  $\mathcal{H}_2$ .

$\hat{\rho}_1$  is an operator!

Notice that, despite the notation,  $\langle \psi_j | \Psi \rangle \langle \Psi | \psi_j \rangle$  is not a complex number but an operator on  $\mathcal{H}_1$ , since

$$|\Psi\rangle = \sum_{m,n} \alpha_{mn} |\phi_m\rangle \otimes |\psi_n\rangle.$$

$\hat{\rho}_1$  is an operator!

Notice that, despite the notation,  $\langle \psi_j | \Psi \rangle \langle \Psi | \psi_j \rangle$  is not a complex number but an operator on  $\mathcal{H}_1$ , since

$$|\Psi\rangle = \sum_{m,n} \alpha_{mn} |\phi_m\rangle \otimes |\psi_n\rangle.$$

In fact, we can write explicitly:

$$\begin{aligned} \hat{\rho}_1 &= \sum_j \langle \psi_j | \left( \sum_{m,n} \alpha_{mn} |\phi_m\rangle \otimes |\psi_n\rangle \right) \left( \sum_{p,q} \alpha_{pq}^* \langle \phi_q| \otimes \langle \psi_q| \right) | \psi_j \rangle \\ &= \sum_{j,m,p} \alpha_{mj} \alpha_{pj}^* |\phi_m\rangle \langle \phi_p|, \end{aligned}$$

using the fact that  $\langle \psi_i | \psi_j \rangle = \delta_{ij}$ .

# Matrix representation of a reduced density operator

In matrix representation, choosing  $\{|\phi_j\rangle\}$  as a basis, we have

$$\hat{\rho}_1 = \begin{pmatrix} \sum_j |\alpha_{1j}|^2 & \sum_j \alpha_{1j}\alpha_{2j}^* & \cdots & \sum_j \alpha_{1j}\alpha_{nj}^* \\ \sum_j \alpha_{2j}\alpha_{1j}^* & \sum_j |\alpha_{2j}|^2 & \cdots & \sum_j \alpha_{2j}\alpha_{nj}^* \\ \vdots & \vdots & \ddots & \vdots \\ \sum_j \alpha_{nj}\alpha_{1j}^* & \sum_j \alpha_{nj}\alpha_{2j}^* & \cdots & \sum_j |\alpha_{nj}|^2 \end{pmatrix},$$

and we have something similar for  $\hat{\rho}_2$  too.

# Relative states

- ▶ The expressions we found can be made to look simpler and can often be applied more directly if we write the state of the composite,

$$|\Psi\rangle = \sum_{i,j} \alpha_{ij} |\phi_i\rangle \otimes |\psi_j\rangle$$

in terms of *relative states*.

# Relative states

- ▶ The expressions we found can be made to look simpler and can often be applied more directly if we write the state of the composite,

$$|\Psi\rangle = \sum_{i,j} \alpha_{ij} |\phi_i\rangle \otimes |\psi_j\rangle$$

in terms of *relative states*.

- ▶ To do this, we first perform the sum over  $j$ :

$$|\Psi\rangle = \sum_i |\phi_i\rangle \otimes \left( \sum_j \alpha_{ij} |\psi_j\rangle \right).$$

## Relative states, continued

The state in round brackets is generally not normalised, so we set

$$\alpha_i := \sum_j |\alpha_{ij}|^2,$$

and write

$$\begin{aligned} |\Psi\rangle &= \sum_i \alpha_i |\phi_i\rangle \otimes \left( \frac{1}{\alpha_i} \sum_j \alpha_{ij} |\psi_j\rangle \right) \\ &=: \sum_i \alpha_i |\phi_i\rangle \otimes |\tilde{\psi}_i\rangle. \end{aligned}$$

The states  $|\tilde{\psi}_i\rangle$  are the *relative states* with respect to the  $|\phi_i\rangle$  in a given state  $|\Psi\rangle$ .

# Making use of relative states

If we calculate the reduced state  $\hat{\rho}_1$  by partially tracing over  $\mathcal{H}_2$  we now obtain, using this notation,

$$\begin{aligned}\hat{\rho}_1 &= \sum_j \langle \psi_j | \left( \sum_i \alpha_i |\phi_i\rangle \otimes |\tilde{\psi}_i\rangle \right) \left( \sum_k \langle \phi_k | \otimes \langle \tilde{\psi}_k | \right) | \psi_j \rangle \\ &= \sum_{i,j,k} \alpha_i \alpha_k \langle \psi_j | \tilde{\psi}_i \rangle \langle \tilde{\psi}_k | \psi_j \rangle |\phi_i\rangle \langle \phi_k| \\ &= \sum_{i,k} \alpha_i \alpha_k \langle \tilde{\psi}_k | \left( \sum_j |\psi_j\rangle \langle \psi_j| \right) | \tilde{\psi}_k \rangle |\phi_i\rangle \langle \phi_k| \\ &= \sum_{i,k} \alpha_i \alpha_k \langle \tilde{\psi}_k | \tilde{\psi}_i \rangle |\phi_i\rangle \langle \phi_k|,\end{aligned}$$

since  $\sum_j |\psi_j\rangle \langle \psi_j| = \hat{\mathbf{1}}$ .

# Another convenient representation of reduced density operators

Thus we can write  $\hat{\rho}_1$  in matrix form as:

$$\hat{\rho}_1 = \begin{pmatrix} \alpha_1^2 & \alpha_1\alpha_2 \langle \tilde{\psi}_2 | \tilde{\psi}_1 \rangle & \cdots & \alpha_1\alpha_n \langle \tilde{\psi}_n | \tilde{\psi}_1 \rangle \\ \alpha_2\alpha_1 \langle \tilde{\psi}_1 | \tilde{\psi}_2 \rangle & \alpha_2^2 & \cdots & \alpha_2\alpha_n \langle \tilde{\psi}_n | \tilde{\psi}_2 \rangle \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_n\alpha_1 \langle \tilde{\psi}_1 | \tilde{\psi}_n \rangle & \alpha_n\alpha_2 \langle \tilde{\psi}_2 | \tilde{\psi}_n \rangle & \cdots & \alpha_n^2 \end{pmatrix}$$

# Another convenient representation of reduced density operators

Thus we can write  $\hat{\rho}_1$  in matrix form as:

$$\hat{\rho}_1 = \begin{pmatrix} \alpha_1^2 & \alpha_1\alpha_2 \langle \tilde{\psi}_2 | \tilde{\psi}_1 \rangle & \cdots & \alpha_1\alpha_n \langle \tilde{\psi}_n | \tilde{\psi}_1 \rangle \\ \alpha_2\alpha_1 \langle \tilde{\psi}_1 | \tilde{\psi}_2 \rangle & \alpha_2^2 & \cdots & \alpha_2\alpha_n \langle \tilde{\psi}_n | \tilde{\psi}_2 \rangle \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_n\alpha_1 \langle \tilde{\psi}_1 | \tilde{\psi}_n \rangle & \alpha_n\alpha_2 \langle \tilde{\psi}_2 | \tilde{\psi}_n \rangle & \cdots & \alpha_n^2 \end{pmatrix}$$

Clearly, the off-diagonal elements are proportional to the inner products of the relative states.

# Another convenient representation of reduced density operators

Thus we can write  $\hat{\rho}_1$  in matrix form as:

$$\hat{\rho}_1 = \begin{pmatrix} \alpha_1^2 & \alpha_1\alpha_2 \langle \tilde{\psi}_2 | \tilde{\psi}_1 \rangle & \cdots & \alpha_1\alpha_n \langle \tilde{\psi}_n | \tilde{\psi}_1 \rangle \\ \alpha_2\alpha_1 \langle \tilde{\psi}_1 | \tilde{\psi}_2 \rangle & \alpha_2^2 & \cdots & \alpha_2\alpha_n \langle \tilde{\psi}_n | \tilde{\psi}_2 \rangle \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_n\alpha_1 \langle \tilde{\psi}_1 | \tilde{\psi}_n \rangle & \alpha_n\alpha_2 \langle \tilde{\psi}_2 | \tilde{\psi}_n \rangle & \cdots & \alpha_n^2 \end{pmatrix}$$

Clearly, the off-diagonal elements are proportional to the inner products of the relative states.

This formulation of the reduced density matrix will be very useful when we look at decoherence next time!

# Reduced states as mixed

- ▶ We know that any such state, say on  $\mathcal{H}_1$ , is represented in general by a density operator acting on  $\mathcal{H}_1$ .

# Reduced states as mixed

- ▶ We know that any such state, say on  $\mathcal{H}_1$ , is represented in general by a density operator acting on  $\mathcal{H}_1$ .
- ▶ That this state, call it  $\hat{\rho}_1$ , must in general be mixed can be shown by example.

## Reduced states as mixed

- ▶ We know that any such state, say on  $\mathcal{H}_1$ , is represented in general by a density operator acting on  $\mathcal{H}_1$ .
- ▶ That this state, call it  $\hat{\rho}_1$ , must in general be mixed can be shown by example.
- ▶ Consider for instance two electrons in the singlet state,

$$|\Psi\rangle = \frac{1}{\sqrt{2}} (|+\rangle |-\rangle - |-\rangle |+\rangle),$$

where  $|+\rangle$  and  $|-\rangle$  are the up and down spin states in (say) the  $x$  direction.

## Reduced states as mixed

- ▶ We know that any such state, say on  $\mathcal{H}_1$ , is represented in general by a density operator acting on  $\mathcal{H}_1$ .
- ▶ That this state, call it  $\hat{\rho}_1$ , must in general be mixed can be shown by example.
- ▶ Consider for instance two electrons in the singlet state,

$$|\Psi\rangle = \frac{1}{\sqrt{2}} (|+\rangle |-\rangle - |-\rangle |+\rangle),$$

where  $|+\rangle$  and  $|-\rangle$  are the up and down spin states in (say) the  $x$  direction.

- ▶ One can compute

$$\hat{\rho}_1 = \hat{\rho}_2 = \frac{1}{2} \hat{1},$$

so the subsystems cannot be in pure states!

# Application to the measurement procedure

- ▶ Unitary measurement interactions *create entanglement*.

$$|\psi\rangle_S |r_0\rangle_A \mapsto \sum_i \beta_i |a_i\rangle_S |r_i\rangle_A.$$

# Application to the measurement procedure

- ▶ Unitary measurement interactions *create entanglement*.

$$|\psi\rangle_S |r_0\rangle_A \mapsto \sum_i \beta_i |a_i\rangle_S |r_i\rangle_A.$$

- ▶ But: calculate the reduced state for the system alone:

$$\text{Tr}_A (|\Psi'\rangle \langle\Psi'|) = \sum_i |\beta_i|^2 |a_i\rangle \langle a_i|.$$

# Application to the measurement procedure

- ▶ Unitary measurement interactions *create entanglement*.

$$|\psi\rangle_S |r_0\rangle_A \mapsto \sum_i \beta_i |a_i\rangle_S |r_i\rangle_A.$$

- ▶ But: calculate the reduced state for the system alone:

$$\text{Tr}_A (|\Psi'\rangle \langle\Psi'|) = \sum_i |\beta_i|^2 |a_i\rangle \langle a_i|.$$

**Exercise:** Check that this is an *improper* mixture!

# Application to the measurement procedure

- ▶ Unitary measurement interactions *create entanglement*.

$$|\psi\rangle_S |r_0\rangle_A \mapsto \sum_i \beta_i |a_i\rangle_S |r_i\rangle_A.$$

- ▶ But: calculate the reduced state for the system alone:

$$\text{Tr}_A (|\Psi'\rangle \langle\Psi'|) = \sum_i |\beta_i|^2 |a_i\rangle \langle a_i|.$$

**Exercise:** Check that this is an *improper* mixture!

- ▶ Again, we'll see lots more about this next time, when we look at decoherence.

# Summary

Today, I:

# Summary

Today, I:

1. Reminded you of some facts about the Born rule.

# Summary

Today, I:

1. Reminded you of some facts about the Born rule.
2. Introduced density operators.

# Summary

Today, I:

1. Reminded you of some facts about the Born rule.
2. Introduced density operators.
3. Introduced quantum entanglement as a qualitatively new phenomenon.

# Summary

Today, I:

1. Reminded you of some facts about the Born rule.
2. Introduced density operators.
3. Introduced quantum entanglement as a qualitatively new phenomenon.
4. Showed how one can associate reduced density matrices with subsystems even in the presence of quantum entanglement.

# References

-  Guido Bacciagaluppi, “Density Operators in Quantum Mechanics”, 1998.