

# IPP-QM-1: Basic quantum formalism

James Read<sup>1</sup>

<sup>1</sup>Faculty of Philosophy, University of Oxford, UK, OX2 6GG

MT24

# The course

1. Basic quantum formalism
2. Density operators and entanglement
3. Decoherence
4. The measurement problem
5. Dynamical collapse theories
6. Bohmian mechanics
7. Everettian structure
8. Everettian probability
9. EPR and Bell's theorem
10. The Bell-CHSH inequalities and possible responses
11. Contextuality
12. The PBR theorem
13. Quantum logic
14. Pragmatism and QBism
15. Relational quantum mechanics
16. Wavefunction realism

## Books etc.

- ▶ David Albert, *Quantum Mechanics and Experience*, Boston: Harvard University Press, 1994.
- ▶ Tim Maudlin, *Philosophy of Physics Volume II: Quantum Mechanics*, Princeton: Princeton University Press, 2019.
- ▶ Tim Maudlin, *Quantum Non-Locality and Relativity*, third edition, Oxford: Wiley-Blackwell, 2011.
- ▶ David Wallace, *The Emergent Multiverse*, Oxford: Oxford University Press, 2012.

## Books etc.

- ▶ David Albert, *Quantum Mechanics and Experience*, Boston: Harvard University Press, 1994.
- ▶ Tim Maudlin, *Philosophy of Physics Volume II: Quantum Mechanics*, Princeton: Princeton University Press, 2019.
- ▶ Tim Maudlin, *Quantum Non-Locality and Relativity*, third edition, Oxford: Wiley-Blackwell, 2011.
- ▶ David Wallace, *The Emergent Multiverse*, Oxford: Oxford University Press, 2012.
  
- ▶ John S. Bell, “Against Measurement”, in *Speakable and Unsayable in Quantum Mechanics*, second edition, Cambridge: Cambridge University Press, 2004.
- ▶ David Wallace, “Philosophy of Quantum Mechanics”, in D. Rickles (ed.), *The Ashgate Companion to Contemporary Philosophy of Physics*, London: Routledge, 2008.

# Formalism

The first three lectures are going to be a lot of formalism...

# Formalism

The first three lectures are going to be a lot of formalism...

...but I'll still try to make it fun!

# Formalism

The first three lectures are going to be a lot of formalism...

...but I'll still try to make it fun!

Highly recommended are:

- ▶ David Wallace, “The Formalism of Quantum Mechanics”, 2005.
- ▶ David Wallace, “The Formalism of Quantum Mechanics II – Density Operators and Entanglement”, 2005.
- ▶ Guido Bacciagaluppi, “Density Operators in Quantum Mechanics”, 1998.
- ▶ Frederic P. Schuller, “Lectures on Quantum Theory”, 2015.

(All on Canvas.)

# Today

Quantum states

Hilbert spaces

More about Hilbert spaces

Linear operators

The physical significance of operators

Wavefunctions and infinite-dimensional Hilbert spaces

# Today

Quantum states

Hilbert spaces

More about Hilbert spaces

Linear operators

The physical significance of operators

Wavefunctions and infinite-dimensional Hilbert spaces

# States

- ▶ Suppose that we have some (quantum) system.

# States

- ▶ Suppose that we have some (quantum) system.
- ▶ Suppose that we have some measurable quantity  $M$  for that system.

# States

- ▶ Suppose that we have some (quantum) system.
- ▶ Suppose that we have some measurable quantity  $M$  for that system.
- ▶  $M$  is *maximally specific* iff whenever we know the value of  $M$  with certainty for the system, then we know everything else (which is knowable) about that system.

# States

- ▶ Suppose that we have some (quantum) system.
- ▶ Suppose that we have some measurable quantity  $M$  for that system.
- ▶  $M$  is *maximally specific* iff whenever we know the value of  $M$  with certainty for the system, then we know everything else (which is knowable) about that system.
  1. if the system is a one-dimensional spinless particle then *position* is maximally specific;
  2. if the system is the spin dofs of a neutron, then spin in the  $z$  direction is maximally specific.
  3. energy *isn't* always maximally specific, because two distinct states can have the same energy.

# States

- ▶ Suppose that we have some (quantum) system.
- ▶ Suppose that we have some measurable quantity  $M$  for that system.
- ▶  $M$  is *maximally specific* iff whenever we know the value of  $M$  with certainty for the system, then we know everything else (which is knowable) about that system.
  1. if the system is a one-dimensional spinless particle then *position* is maximally specific;
  2. if the system is the spin dofs of a neutron, then spin in the  $z$  direction is maximally specific.
  3. energy *isn't* always maximally specific, because two distinct states can have the same energy.
- ▶ Write the states of definite  $m$  as  $|m_1\rangle, |m_2\rangle, \dots, |m_n\rangle$ .

# States

- ▶ Suppose that we have some (quantum) system.
- ▶ Suppose that we have some measurable quantity  $M$  for that system.
- ▶  $M$  is *maximally specific* iff whenever we know the value of  $M$  with certainty for the system, then we know everything else (which is knowable) about that system.
  1. if the system is a one-dimensional spinless particle then *position* is maximally specific;
  2. if the system is the spin dofs of a neutron, then spin in the  $z$  direction is maximally specific.
  3. energy *isn't* always maximally specific, because two distinct states can have the same energy.
- ▶ Write the states of definite  $m$  as  $|m_1\rangle, |m_2\rangle, \dots, |m_n\rangle$ .
- ▶  $|m_i\rangle$  is a state such that, if we measure  $M$  when the system is in that state (assuming that we have an ideal measuring device), then we definitely get the result  $m_i$ .

# Superpositions

- ▶ In classical physics, this would be the whole story: quantities always have measurable values, so the system would be in one of the  $|m_i\rangle$ , and we just ask which.

# Superpositions

- ▶ In classical physics, this would be the whole story: quantities always have measurable values, so the system would be in one of the  $|m_i\rangle$ , and we just ask which.
- ▶ But quantum mechanics permits *superpositions*, e.g.

$$|\psi\rangle = \alpha_1 |m_1\rangle + \alpha_2 |m_2\rangle + \dots + \alpha_n |m_N\rangle, \quad \forall \alpha_j \in \mathbb{C}.$$

# Superpositions

- ▶ In classical physics, this would be the whole story: quantities always have measurable values, so the system would be in one of the  $|m_i\rangle$ , and we just ask which.
- ▶ But quantum mechanics permits *superpositions*, e.g.

$$|\psi\rangle = \alpha_1 |m_1\rangle + \alpha_2 |m_2\rangle + \dots + \alpha_n |m_N\rangle, \quad \forall \alpha_j \in \mathbb{C}.$$

- ▶ What these superposition states *mean*, or *represent*, is part of the great puzzle of quantum mechanics.

# Superpositions

- ▶ In classical physics, this would be the whole story: quantities always have measurable values, so the system would be in one of the  $|m_i\rangle$ , and we just ask which.
- ▶ But quantum mechanics permits *superpositions*, e.g.

$$|\psi\rangle = \alpha_1 |m_1\rangle + \alpha_2 |m_2\rangle + \dots + \alpha_n |m_N\rangle, \quad \forall \alpha_j \in \mathbb{C}.$$

- ▶ What these superposition states *mean*, or *represent*, is part of the great puzzle of quantum mechanics.
- ▶ But *operationally*, we can make sense of them via the *Born rule*: If  $|\psi\rangle$  is the state of the system, and we measure  $M$ , then the probability of getting result  $m_j$  is  $|\alpha_j|^2$ .

# Superpositions

- ▶ In classical physics, this would be the whole story: quantities always have measurable values, so the system would be in one of the  $|m_i\rangle$ , and we just ask which.
- ▶ But quantum mechanics permits *superpositions*, e.g.

$$|\psi\rangle = \alpha_1 |m_1\rangle + \alpha_2 |m_2\rangle + \dots + \alpha_n |m_N\rangle, \quad \forall \alpha_j \in \mathbb{C}.$$

- ▶ What these superposition states *mean*, or *represent*, is part of the great puzzle of quantum mechanics.
- ▶ But *operationally*, we can make sense of them via the *Born rule*: If  $|\psi\rangle$  is the state of the system, and we measure  $M$ , then the probability of getting result  $m_j$  is  $|\alpha_j|^2$ .
- ▶ Because probabilities sum to one, this implies that quantum states must be *normalised*,

$$\sum_{i=1}^N |\alpha_i|^2 = 1.$$

# Today

Quantum states

**Hilbert spaces**

More about Hilbert spaces

Linear operators

The physical significance of operators

Wavefunctions and infinite-dimensional Hilbert spaces

# Hilbert spaces

- ▶ Mathematically, to what kind of space do quantum states belong?

# Hilbert spaces

- ▶ Mathematically, to what kind of space do quantum states belong?
- ▶ For the quantum system under consideration, call its *Hilbert space* the space

$$\mathcal{H} := \left\{ \sum_i \alpha_i |m_i\rangle : \text{all complex numbers } \alpha_i \right\}.$$

# Hilbert spaces

- ▶ Mathematically, to what kind of space do quantum states belong?
- ▶ For the quantum system under consideration, call its *Hilbert space* the space

$$\mathcal{H} := \left\{ \sum_i \alpha_i |m_i\rangle : \text{all complex numbers } \alpha_i \right\}.$$

- ▶ Elements of a Hilbert space needn't be normalised (only the physical ones are), and are called *vectors*.

# Hilbert space inner product

- ▶ We can define an *inner product* on Hilbert space. Like all inner products, this eats two vectors and spits out a (here complex) number.

# Hilbert space inner product

- ▶ We can define an *inner product* on Hilbert space. Like all inner products, this eats two vectors and spits out a (here complex) number.
- ▶ (Actually, technically a Hilbert space is *defined* as having an inner product—I'll come back to this.)

# Hilbert space inner product

- ▶ We can define an *inner product* on Hilbert space. Like all inner products, this eats two vectors and spits out a (here complex) number.
- ▶ (Actually, technically a Hilbert space is *defined* as having an inner product—I'll come back to this.)
- ▶ If  $|\psi\rangle = \sum_i \alpha_i |m_i\rangle$  and  $|\phi\rangle = \sum_i \beta_i |m_i\rangle$  are vectors, then their inner product is defined as

$$\langle \psi | \phi \rangle := \sum_i \alpha_i^* \beta_i.$$

# Hilbert space inner product

- ▶ We can define an *inner product* on Hilbert space. Like all inner products, this eats two vectors and spits out a (here complex) number.
- ▶ (Actually, technically a Hilbert space is *defined* as having an inner product—I'll come back to this.)
- ▶ If  $|\psi\rangle = \sum_i \alpha_i |m_i\rangle$  and  $|\phi\rangle = \sum_i \beta_i |m_i\rangle$  are vectors, then their inner product is defined as

$$\langle \psi | \phi \rangle := \sum_i \alpha_i^* \beta_i.$$

- ▶ Properties of the inner product:
  1. *Linear*:  $\langle \phi | (\alpha |\psi_1\rangle + \beta |\psi_2\rangle) \rangle = \alpha \langle \phi | \psi_1 \rangle + \beta \langle \phi | \psi_2 \rangle$
  2. *Satisfies*:  $\langle \phi | \psi \rangle = (\langle \psi | \phi \rangle)^*$ .
  3. *Positive*: If  $|\psi\rangle \neq 0$  then  $\langle \psi | \psi \rangle \in \mathbb{R}^+$ .

# Hilbert space inner product

- ▶ We can define an *inner product* on Hilbert space. Like all inner products, this eats two vectors and spits out a (here complex) number.
- ▶ (Actually, technically a Hilbert space is *defined* as having an inner product—I'll come back to this.)
- ▶ If  $|\psi\rangle = \sum_i \alpha_i |m_i\rangle$  and  $|\phi\rangle = \sum_i \beta_i |m_i\rangle$  are vectors, then their inner product is defined as

$$\langle\psi|\phi\rangle := \sum_i \alpha_i^* \beta_i.$$

- ▶ Properties of the inner product:
  1. *Linear*:  $\langle\phi|(\alpha|\psi_1\rangle + \beta|\psi_2\rangle) = \alpha\langle\phi|\psi_1\rangle + \beta\langle\phi|\psi_2\rangle$
  2. Satisfies:  $\langle\phi|\psi\rangle = (\langle\psi|\phi\rangle)^*$ .
  3. *Positive*: If  $|\psi\rangle \neq 0$  then  $\langle\psi|\psi\rangle \in \mathbb{R}^+$ .
- ▶ We can write the normalisation condition on states as:  $|\psi\rangle$  is normalised iff  $\langle\psi|\psi\rangle = 1$ .

# Mathematical approaches to Hilbert space

- ▶ Here, I've presented the Hilbert space and the inner product by starting with a particular set of vectors and building the rest of the space from them.

# Mathematical approaches to Hilbert space

- ▶ Here, I've presented the Hilbert space and the inner product by starting with a particular set of vectors and building the rest of the space from them.
- ▶ In more mathematically formal literature, one defines Hilbert space in a more abstract way:

*A Hilbert space is a real or complex inner product space that is also a complete metric space with respect to the distance function induced by the inner product.*

# Mathematical approaches to Hilbert space

- ▶ Here, I've presented the Hilbert space and the inner product by starting with a particular set of vectors and building the rest of the space from them.
- ▶ In more mathematically formal literature, one defines Hilbert space in a more abstract way:

*A Hilbert space is a real or complex inner product space that is also a complete metric space with respect to the distance function induced by the inner product.*

- ▶ See e.g. the Schuller notes for more details.

# Bases for Hilbert space

- ▶ A *basis* for a Hilbert space  $\mathcal{H}$  is a set of vectors in  $\mathcal{H}$  s.t.:
  1. Any vector in  $\mathcal{H}$  can be written as a superposition of basis vectors;
  2. No basis vector is a superposition of other basis vectors.

# Bases for Hilbert space

- ▶ A *basis* for a Hilbert space  $\mathcal{H}$  is a set of vectors in  $\mathcal{H}$  s.t.:
  1. Any vector in  $\mathcal{H}$  can be written as a superposition of basis vectors;
  2. No basis vector is a superposition of other basis vectors.
- ▶ It is typical to insist that a basis be *orthonormal*, which means that it also satisfies:
  1. If  $|\psi\rangle$  and  $|\phi\rangle$  are (distinct) basis vectors, then  $\langle\psi|\phi\rangle = 0$ .
  2. If  $|\psi\rangle$  is a basis vector, then  $\langle\psi|\psi\rangle = 1$ .

# Bases for Hilbert space

- ▶ A *basis* for a Hilbert space  $\mathcal{H}$  is a set of vectors in  $\mathcal{H}$  s.t.:
  1. Any vector in  $\mathcal{H}$  can be written as a superposition of basis vectors;
  2. No basis vector is a superposition of other basis vectors.
- ▶ It is typical to insist that a basis be *orthonormal*, which means that it also satisfies:
  1. If  $|\psi\rangle$  and  $|\phi\rangle$  are (distinct) basis vectors, then  $\langle\psi|\phi\rangle = 0$ .
  2. If  $|\psi\rangle$  is a basis vector, then  $\langle\psi|\psi\rangle = 1$ .
- ▶ **Exercise:** Prove that any two bases for  $\mathcal{H}$  have the same number of vectors.

# Bases for Hilbert space

- ▶ A *basis* for a Hilbert space  $\mathcal{H}$  is a set of vectors in  $\mathcal{H}$  s.t.:
  1. Any vector in  $\mathcal{H}$  can be written as a superposition of basis vectors;
  2. No basis vector is a superposition of other basis vectors.
- ▶ It is typical to insist that a basis be *orthonormal*, which means that it also satisfies:
  1. If  $|\psi\rangle$  and  $|\phi\rangle$  are (distinct) basis vectors, then  $\langle\psi|\phi\rangle = 0$ .
  2. If  $|\psi\rangle$  is a basis vector, then  $\langle\psi|\psi\rangle = 1$ .
- ▶ **Exercise:** Prove that any two bases for  $\mathcal{H}$  have the same number of vectors.
- ▶ Let  $|\psi\rangle \in \mathcal{H}$  and let  $\{|i\rangle\}$  be any orthonormal basis. By definition, there will be complex numbers  $\{\alpha_i\}$  s.t.  $|\psi\rangle = \sum_i \alpha_i |i\rangle$ . Because the basis is orthonormal,  $\langle i|\psi\rangle = \alpha_i$ ; that is, we have the useful expansion

$$|\psi\rangle = \sum_i |i\rangle \langle i|\psi\rangle$$

# Bases and measurement

- ▶ Suppose  $K$  is *another* maximally specific measurement on the same system, and suppose that the states with definite values of  $K$  are  $|k_1\rangle, \dots, |k_N\rangle$ .

# Bases and measurement

- ▶ Suppose  $K$  is *another* maximally specific measurement on the same system, and suppose that the states with definite values of  $K$  are  $|k_1\rangle, \dots, |k_N\rangle$ .
- ▶ If  $|\psi\rangle$  is an arbitrary state, what's the probability of getting result  $k_j$  when we measure  $K$ ?

# Bases and measurement

- ▶ Suppose  $K$  is *another* maximally specific measurement on the same system, and suppose that the states with definite values of  $K$  are  $|k_1\rangle, \dots, |k_N\rangle$ .
- ▶ If  $|\psi\rangle$  is an arbitrary state, what's the probability of getting result  $k_i$  when we measure  $K$ ?
- ▶ It's a *postulate* of 'standard', textbook quantum mechanics that the probability is  $|\langle k_i|\psi\rangle|^2$ .

# Summary so far

- ▶ A physical system is represented by a Hilbert space.

# Summary so far

- ▶ A physical system is represented by a Hilbert space.
- ▶ States of that system are represented by normalised vectors in that Hilbert space.

# Summary so far

- ▶ A physical system is represented by a Hilbert space.
- ▶ States of that system are represented by normalised vectors in that Hilbert space.
- ▶ A maximally specific measurement is represented by a basis of the Hilbert space.

# Summary so far

- ▶ A physical system is represented by a Hilbert space.
- ▶ States of that system are represented by normalised vectors in that Hilbert space.
- ▶ A maximally specific measurement is represented by a basis of the Hilbert space.
- ▶ If  $K$  is a maximally specific measurement and  $|k_i\rangle$  has definite value  $k_i$  of  $K$ , then the probability is of getting  $k_i$  when we measure  $K$  on the state  $|\psi\rangle$  is  $|\langle k_i|\psi\rangle|^2$ .

# Today

Quantum states

Hilbert spaces

**More about Hilbert spaces**

Linear operators

The physical significance of operators

Wavefunctions and infinite-dimensional Hilbert spaces

# Subspaces

- ▶ Suppose we pick some subset  $\{|n_1\rangle, \dots, |n_P\rangle\}$  of a basis, and consider all those vectors which can be expressed as superpositions of the members of that subset.

# Subspaces

- ▶ Suppose we pick some subset  $\{|n_1\rangle, \dots, |n_P\rangle\}$  of a basis, and consider all those vectors which can be expressed as superpositions of the members of that subset.
- ▶ This set is a Hilbert space in its own right!

# Subspaces

- ▶ Suppose we pick some subset  $\{|n_1\rangle, \dots, |n_P\rangle\}$  of a basis, and consider all those vectors which can be expressed as superpositions of the members of that subset.
- ▶ This set is a Hilbert space in its own right!  
(**Exercise:** prove this! You just need to show that it satisfies the definition of a Hilbert space...)

# Subspaces

- ▶ Suppose we pick some subset  $\{|n_1\rangle, \dots, |n_P\rangle\}$  of a basis, and consider all those vectors which can be expressed as superpositions of the members of that subset.
- ▶ This set is a Hilbert space in its own right!  
(**Exercise:** prove this! You just need to show that it satisfies the definition of a Hilbert space...)
- ▶ It's a smaller Hilbert space, entirely contained within the original one.

# Subspaces

- ▶ Suppose we pick some subset  $\{|n_1\rangle, \dots, |n_P\rangle\}$  of a basis, and consider all those vectors which can be expressed as superpositions of the members of that subset.
- ▶ This set is a Hilbert space in its own right!  
(**Exercise:** prove this! You just need to show that it satisfies the definition of a Hilbert space...)
- ▶ It's a smaller Hilbert space, entirely contained within the original one.
- ▶ Such spaces are called *subspaces*.

# Global phase transformations

- ▶ The probability rule defined previously gives the same probabilities for all measurements if we carry out a *global phase transformation* of all states,

$$|\psi\rangle \rightarrow \exp(i\theta) |\psi\rangle .$$

# Global phase transformations

- ▶ The probability rule defined previously gives the same probabilities for all measurements if we carry out a *global phase transformation* of all states,

$$|\psi\rangle \rightarrow \exp(i\theta) |\psi\rangle .$$

- ▶ As such, it's standard to regard the physical system as unchanged by such transformations.

# Global phase transformations

- ▶ The probability rule defined previously gives the same probabilities for all measurements if we carry out a *global phase transformation* of all states,

$$|\psi\rangle \rightarrow \exp(i\theta) |\psi\rangle .$$

- ▶ As such, it's standard to regard the physical system as unchanged by such transformations.
- ▶ So strictly, it isn't states in Hilbert space which represent physical systems—it's equivalence classes of states related by global phase, sometimes called *rays*.

# Global phase transformations

- ▶ The probability rule defined previously gives the same probabilities for all measurements if we carry out a *global phase transformation* of all states,

$$|\psi\rangle \rightarrow \exp(i\theta) |\psi\rangle .$$

- ▶ As such, it's standard to regard the physical system as unchanged by such transformations.
- ▶ So strictly, it isn't states in Hilbert space which represent physical systems—it's equivalence classes of states related by global phase, sometimes called *rays*.
- ▶ Practically, however, it's easier to just carry on working with vectors in Hilbert space.

# On the reality of the global phase

- ▶ Denying the reality of the global phase is, to put it mildly, *entirely mainstream*.

# On the reality of the global phase

- ▶ Denying the reality of the global phase is, to put it mildly, *entirely mainstream*.
- ▶ A few philosophers have resisted this, though. Stepping back, it's related to Leibniz shifts, and the general literature on symmetry transformations in physics.

# On the reality of the global phase

- ▶ Denying the reality of the global phase is, to put it mildly, *entirely mainstream*.
- ▶ A few philosophers have resisted this, though. Stepping back, it's related to Leibniz shifts, and the general literature on symmetry transformations in physics.
- ▶ For defences of the mainstream, see (Wallace 2022) or (Gao 2024); I won't question it further in these lectures.

# Today

Quantum states

Hilbert spaces

More about Hilbert spaces

**Linear operators**

The physical significance of operators

Wavefunctions and infinite-dimensional Hilbert spaces

# Linear operators

- ▶ A *linear operator*  $\hat{A}$  is a map from Hilbert space to itself which has the linearity property:

$$\hat{A}(\alpha |\psi\rangle + \beta |\phi\rangle) = \alpha \hat{A}|\psi\rangle + \beta \hat{A}|\phi\rangle .$$

# Linear operators

- ▶ A *linear operator*  $\hat{A}$  is a map from Hilbert space to itself which has the linearity property:

$$\hat{A}(\alpha |\psi\rangle + \beta |\phi\rangle) = \alpha \hat{A}|\psi\rangle + \beta \hat{A}|\phi\rangle .$$

- ▶ Simple example:  $|\phi\rangle \langle \chi|$ , where  $|\psi\rangle$  and  $|\chi\rangle$  are arbitrary (normalised) vectors; this is defined as follows:  
 $(|\phi\rangle \langle \chi|) |\psi\rangle = |\phi\rangle \times \langle \chi|\psi\rangle$ .

# Projectors

- ▶ An important sub-class of linear operators are *one-dimensional projectors*, which can be written  $|\phi\rangle\langle\phi|$ , for arbitrary normalised  $|\phi\rangle$ .

# Projectors

- ▶ An important sub-class of linear operators are *one-dimensional projectors*, which can be written  $|\phi\rangle\langle\phi|$ , for arbitrary normalised  $|\phi\rangle$ .
- ▶ They're called 'projectors' because, when they act on a vector, they project out the  $|\phi\rangle$  component of the vector and throw the rest away.

# Projectors

- ▶ An important sub-class of linear operators are *one-dimensional projectors*, which can be written  $|\phi\rangle\langle\phi|$ , for arbitrary normalised  $|\phi\rangle$ .
- ▶ They're called 'projectors' because, when they act on a vector, they project out the  $|\phi\rangle$  component of the vector and throw the rest away.
- ▶ E.g., acting on  $|\psi\rangle = \alpha|\phi\rangle + \beta|\chi\rangle$  with  $|\phi\rangle\langle\phi|$  yields  $\alpha|\phi\rangle$  (where  $|\phi\rangle$  and  $|\psi\rangle$  are part of an orthonormal basis for the Hilbert space).

# Projectors

- ▶ An important sub-class of linear operators are *one-dimensional projectors*, which can be written  $|\phi\rangle\langle\phi|$ , for arbitrary normalised  $|\phi\rangle$ .
- ▶ They're called 'projectors' because, when they act on a vector, they project out the  $|\phi\rangle$  component of the vector and throw the rest away.
- ▶ E.g., acting on  $|\psi\rangle = \alpha|\phi\rangle + \beta|\chi\rangle$  with  $|\phi\rangle\langle\phi|$  yields  $\alpha|\phi\rangle$  (where  $|\phi\rangle$  and  $|\psi\rangle$  are part of an orthonormal basis for the Hilbert space).
- ▶ More general (i.e., higher-dimensional) projectors don't just project out one particular component of a vector: they project out all components in some particular subspace.

# Resolutions of the identity

- ▶ The limiting case of a projector is one which projects onto the entire Hilbert space—and so leaves the vector alone!

# Resolutions of the identity

- ▶ The limiting case of a projector is one which projects onto the entire Hilbert space—and so leaves the vector alone!
- ▶ The *identity operator* can be written in the same way as any other projector:

$$\hat{1} = \sum_i |n_i\rangle \langle n_i|,$$

where  $\{|n_i\rangle\}$  is any basis. An expression like this for  $\hat{1}$  is called a *resolution of the identity*.

# Resolutions of the identity

- ▶ By construction, for any vector  $\hat{1} |\psi\rangle = |\psi\rangle$ .

# Resolutions of the identity

- ▶ By construction, for any vector  $\hat{1} |\psi\rangle = |\psi\rangle$ .
- ▶ This means that we can insert resolutions of the identity into quantum expressions wherever we please, e.g.:
  - ▶  $|\psi\rangle = \hat{1} |\psi\rangle = \sum_i (|n_i\rangle \langle n_i|) |\psi\rangle = \sum_i |n_i\rangle \langle n_i|\psi\rangle$ .
  - ▶  $\hat{A} = \hat{1}\hat{A}\hat{1} = (\sum_i |n_i\rangle \langle n_i|) \hat{A} (\sum_j |n_j\rangle \langle n_j|) = \sum_{i,j} |n_i\rangle \langle n_j| \langle n_i| \hat{A} |n_j\rangle$ .

# Resolutions of the identity

- ▶ By construction, for any vector  $\hat{1} |\psi\rangle = |\psi\rangle$ .
- ▶ This means that we can insert resolutions of the identity into quantum expressions wherever we please, e.g.:
  - ▶  $|\psi\rangle = \hat{1} |\psi\rangle = \sum_i (|n_i\rangle \langle n_i|) |\psi\rangle = \sum_i |n_i\rangle \langle n_i|\psi\rangle$ .
  - ▶  $\hat{A} = \hat{1}\hat{A}\hat{1} = (\sum_i |n_i\rangle \langle n_i|) \hat{A} (\sum_j |n_j\rangle \langle n_j|) = \sum_{i,j} |n_i\rangle \langle n_j| \langle n_i| \hat{A} |n_j\rangle$ .
- ▶ The later of these is called a *matrix representation* of  $\hat{A}$ , and we can write  $A_{ij} := \langle n_i| \hat{A} |n_j\rangle$ .

# Resolutions of the identity

- ▶ By construction, for any vector  $\hat{1} |\psi\rangle = |\psi\rangle$ .
- ▶ This means that we can insert resolutions of the identity into quantum expressions wherever we please, e.g.:
  - ▶  $|\psi\rangle = \hat{1} |\psi\rangle = \sum_i (|n_i\rangle \langle n_i|) |\psi\rangle = \sum_i |n_i\rangle \langle n_i|\psi\rangle$ .
  - ▶  $\hat{A} = \hat{1}\hat{A}\hat{1} = (\sum_i |n_i\rangle \langle n_i|) \hat{A} (\sum_j |n_j\rangle \langle n_j|) = \sum_{i,j} |n_i\rangle \langle n_j| \langle n_i| \hat{A} |n_j\rangle$ .
- ▶ The later of these is called a *matrix representation* of  $\hat{A}$ , and we can write  $A_{ij} := \langle n_i| \hat{A} |n_j\rangle$ .
- ▶ This gives us the components  $A_{ij}$  of the matrix associated with the operator  $\hat{A}$  in the  $\{|n_i\rangle\}$  basis.

# Eigenvalues and eigenvectors

- ▶ Recall that an *eigenvector*  $\mathbf{v}$  of a matrix  $\mathbf{M}$  is a column vector satisfying  $\mathbf{M}\mathbf{v} = \lambda\mathbf{v}$ , and the number  $\lambda$  is called the *eigenvalue* associated with  $\mathbf{v}$ .

# Eigenvalues and eigenvectors

- ▶ Recall that an *eigenvector*  $\mathbf{v}$  of a matrix  $\mathbf{M}$  is a column vector satisfying  $\mathbf{M}\mathbf{v} = \lambda\mathbf{v}$ , and the number  $\lambda$  is called the *eigenvalue* associated with  $\mathbf{v}$ .
- ▶ This translates to Hilbert space in the obvious way:  $|\psi\rangle$  is an eigenvector of an operator  $\hat{M}$  iff for some  $\lambda \in \mathbb{C}$ ,  $\hat{M}|\psi\rangle = \lambda|\psi\rangle$ .

# Eigenvalues and eigenvectors

- ▶ Recall that an *eigenvector*  $\mathbf{v}$  of a matrix  $\mathbf{M}$  is a column vector satisfying  $\mathbf{M}\mathbf{v} = \lambda\mathbf{v}$ , and the number  $\lambda$  is called the *eigenvalue* associated with  $\mathbf{v}$ .
- ▶ This translates to Hilbert space in the obvious way:  $|\psi\rangle$  is an eigenvector of an operator  $\hat{M}$  iff for some  $\lambda \in \mathbb{C}$ ,  $\hat{M}|\psi\rangle = \lambda|\psi\rangle$ .
- ▶ If  $\hat{M}$  has more than one eigenvector with the same eigenvalue, it is called *degenerate*. The set of all eigenvalues of  $\hat{M}$  is called the *spectrum* of  $\hat{M}$ .

# Eigenvalues and eigenvectors

- ▶ Recall that an *eigenvector*  $\mathbf{v}$  of a matrix  $\mathbf{M}$  is a column vector satisfying  $\mathbf{M}\mathbf{v} = \lambda\mathbf{v}$ , and the number  $\lambda$  is called the *eigenvalue* associated with  $\mathbf{v}$ .
- ▶ This translates to Hilbert space in the obvious way:  $|\psi\rangle$  is an eigenvector of an operator  $\hat{M}$  iff for some  $\lambda \in \mathbb{C}$ ,  $\hat{M}|\psi\rangle = \lambda|\psi\rangle$ .
- ▶ If  $\hat{M}$  has more than one eigenvector with the same eigenvalue, it is called *degenerate*. The set of all eigenvalues of  $\hat{M}$  is called the *spectrum* of  $\hat{M}$ .
- ▶ An *eigensubspace* of an operator is the subspace defined by all those eigenvectors with some particular eigenvalue.

# Eigenvalues and eigenvectors

- ▶ Recall that an *eigenvector*  $\mathbf{v}$  of a matrix  $\mathbf{M}$  is a column vector satisfying  $\mathbf{M}\mathbf{v} = \lambda\mathbf{v}$ , and the number  $\lambda$  is called the *eigenvalue* associated with  $\mathbf{v}$ .
- ▶ This translates to Hilbert space in the obvious way:  $|\psi\rangle$  is an eigenvector of an operator  $\hat{M}$  iff for some  $\lambda \in \mathbb{C}$ ,  $\hat{M}|\psi\rangle = \lambda|\psi\rangle$ .
- ▶ If  $\hat{M}$  has more than one eigenvector with the same eigenvalue, it is called *degenerate*. The set of all eigenvalues of  $\hat{M}$  is called the *spectrum* of  $\hat{M}$ .
- ▶ An *eigensubspace* of an operator is the subspace defined by all those eigenvectors with some particular eigenvalue.
- ▶ (So, non-degenerate operators have only one-dimensional eigensubspaces.)

# Adjoint operators

- ▶ Recall that the adjoint  $\mathbf{A}^\dagger$  of a complex matrix is defined as the complex conjugate of its transpose:  $A_{ij}^\dagger := (A_{ji})^*$ .

# Adjoint operators

- ▶ Recall that the adjoint  $\mathbf{A}^\dagger$  of a complex matrix is defined as the complex conjugate of its transpose:  $A_{ij}^\dagger := (A_{ji})^*$ .
- ▶ We can define an abstract version of this:  $\hat{A}^\dagger$  is defined by  $\langle \psi | \hat{A}^\dagger | \chi \rangle := \left( \langle \chi | \hat{A} | \psi \rangle \right)^*$ .

# Adjoint operators

- ▶ Recall that the adjoint  $\mathbf{A}^\dagger$  of a complex matrix is defined as the complex conjugate of its transpose:  $A_{ij}^\dagger := (A_{ji})^*$ .
- ▶ We can define an abstract version of this:  $\hat{A}^\dagger$  is defined by  $\langle \psi | \hat{A}^\dagger | \chi \rangle := \left( \langle \chi | \hat{A} | \psi \rangle \right)^*$ .
- ▶ Operators which commute with their adjoints (i.e.  $\hat{A}\hat{A}^\dagger = \hat{A}^\dagger\hat{A}$ ) are called *normal*. These are significant because of the *spectral theorem*:

# Adjoint operators

- ▶ Recall that the adjoint  $\mathbf{A}^\dagger$  of a complex matrix is defined as the complex conjugate of its transpose:  $A_{ij}^\dagger := (A_{ji})^*$ .
- ▶ We can define an abstract version of this:  $\hat{A}^\dagger$  is defined by  $\langle \psi | \hat{A}^\dagger | \chi \rangle := \left( \langle \chi | \hat{A} | \psi \rangle \right)^*$ .
- ▶ Operators which commute with their adjoints (i.e.  $\hat{A}\hat{A}^\dagger = \hat{A}^\dagger\hat{A}$ ) are called *normal*. These are significant because of the *spectral theorem*:

**Spectral theorem:** If  $\hat{C}$  is a normal operator, then there is an orthonormal basis for the Hilbert space made up entirely of eigenvectors of  $\hat{C}$ .

# Adjoint operators

- ▶ Recall that the adjoint  $\mathbf{A}^\dagger$  of a complex matrix is defined as the complex conjugate of its transpose:  $A_{ij}^\dagger := (A_{ji})^*$ .
- ▶ We can define an abstract version of this:  $\hat{A}^\dagger$  is defined by  $\langle \psi | \hat{A}^\dagger | \chi \rangle := \left( \langle \chi | \hat{A} | \psi \rangle \right)^*$ .
- ▶ Operators which commute with their adjoints (i.e.  $\hat{A}\hat{A}^\dagger = \hat{A}^\dagger\hat{A}$ ) are called *normal*. These are significant because of the *spectral theorem*:  
**Spectral theorem:** If  $\hat{C}$  is a normal operator, then there is an orthonormal basis for the Hilbert space made up entirely of eigenvectors of  $\hat{C}$ .
- ▶ (For further details on the spectral theorem, see e.g. the Schuller notes.)

# Adjoint operators

- ▶ Recall that the adjoint  $\mathbf{A}^\dagger$  of a complex matrix is defined as the complex conjugate of its transpose:  $A_{ij}^\dagger := (A_{ji})^*$ .
- ▶ We can define an abstract version of this:  $\hat{A}^\dagger$  is defined by  $\langle \psi | \hat{A}^\dagger | \chi \rangle := (\langle \chi | \hat{A} | \psi \rangle)^*$ .
- ▶ Operators which commute with their adjoints (i.e.  $\hat{A}\hat{A}^\dagger = \hat{A}^\dagger\hat{A}$ ) are called *normal*. These are significant because of the *spectral theorem*:  
**Spectral theorem:** If  $\hat{C}$  is a normal operator, then there is an orthonormal basis for the Hilbert space made up entirely of eigenvectors of  $\hat{C}$ .
- ▶ (For further details on the spectral theorem, see e.g. the Schuller notes.)
- ▶ If  $\hat{C}$  is non-degenerate, then there's a *unique* such basis.

# Spectral resolutions

- ▶ Suppose  $\hat{C}$  is a normal operator, and let  $\{|c_i\rangle\}$  be an orthonormal basis comprised of its eigenvectors (so  $\hat{C}|c_i\rangle = c_i|c_i\rangle$ ).

# Spectral resolutions

- ▶ Suppose  $\hat{C}$  is a normal operator, and let  $\{|c_i\rangle\}$  be an orthonormal basis comprised of its eigenvectors (so  $\hat{C}|c_i\rangle = c_i|c_i\rangle$ ).
- ▶ In that case, the *spectral resolution* of  $\hat{C}$  is

$$\hat{C} = \sum_i c_i |c_i\rangle \langle c_i|.$$

# Spectral resolutions

- ▶ Suppose  $\hat{C}$  is a normal operator, and let  $\{|c_i\rangle\}$  be an orthonormal basis comprised of its eigenvectors (so  $\hat{C}|c_i\rangle = c_i|c_i\rangle$ ).
- ▶ In that case, the *spectral resolution* of  $\hat{C}$  is

$$\hat{C} = \sum_i c_i |c_i\rangle \langle c_i|.$$

- ▶ The function  $f(\hat{C})$  of a normal operator  $\hat{C}$  can be written

$$f(\hat{C}) = \sum_i f(c_i) |c_i\rangle \langle c_i|.$$

**(Exercise: Prove this!)**

# Spectral resolutions

- ▶ Suppose  $\hat{C}$  is a normal operator, and let  $\{|c_i\rangle\}$  be an orthonormal basis comprised of its eigenvectors (so  $\hat{C}|c_i\rangle = c_i|c_i\rangle$ ).
- ▶ In that case, the *spectral resolution* of  $\hat{C}$  is

$$\hat{C} = \sum_i c_i |c_i\rangle \langle c_i|.$$

- ▶ The function  $f(\hat{C})$  of a normal operator  $\hat{C}$  can be written

$$f(\hat{C}) = \sum_i f(c_i) |c_i\rangle \langle c_i|.$$

**(Exercise:** Prove this!)

- ▶ If  $\hat{A}$  and  $\hat{B}$  are normal operators which commute, then there exists a normal operator  $\hat{C}$  and functions  $f, g$  such that  $\hat{A} = f(\hat{C})$  and  $\hat{B} = g(\hat{C})$ .

# A zoo of operators

- ▶ A normal operator commutes with its adjoint:  $\hat{A}\hat{A}^\dagger = \hat{A}^\dagger\hat{A}$ .

# A zoo of operators

- ▶ A normal operator commutes with its adjoint:  $\hat{A}\hat{A}^\dagger = \hat{A}^\dagger\hat{A}$ .
- ▶ *Self-adjoint* operators are such that  $\hat{A} = \hat{A}^\dagger$ .

# A zoo of operators

- ▶ A normal operator commutes with its adjoint:  $\hat{A}\hat{A}^\dagger = \hat{A}^\dagger\hat{A}$ .
- ▶ *Self-adjoint* operators are such that  $\hat{A} = \hat{A}^\dagger$ .
- ▶ All self-adjoint operators are normal, but not *vice versa*.

# A zoo of operators

- ▶ A normal operator commutes with its adjoint:  $\hat{A}\hat{A}^\dagger = \hat{A}^\dagger\hat{A}$ .
- ▶ *Self-adjoint* operators are such that  $\hat{A} = \hat{A}^\dagger$ .
- ▶ All self-adjoint operators are normal, but not *vice versa*.
- ▶ (E.g., *unitary* operators—for which  $\hat{A}\hat{A}^\dagger = \hat{1}$ —are normal but not self-adjoint.)

# Today

Quantum states

Hilbert spaces

More about Hilbert spaces

Linear operators

**The physical significance of operators**

Wavefunctions and infinite-dimensional Hilbert spaces

# Self-adjoint operators and measurements

- ▶ Recall that a maximally specific measurement  $M$  is specified by an orthonormal basis  $\{|m_i\rangle\}$  of states with definite values of  $M$ , together with the value  $m_i$  of  $M$  for each state.

# Self-adjoint operators and measurements

- ▶ Recall that a maximally specific measurement  $M$  is specified by an orthonormal basis  $\{|m_i\rangle\}$  of states with definite values of  $M$ , together with the value  $m_i$  of  $M$  for each state.
- ▶ We can codify  $M$  into an operator by defining

$$\hat{M} = \sum_i m_i |m_i\rangle \langle m_i|.$$

# Self-adjoint operators and measurements

- ▶ Recall that a maximally specific measurement  $M$  is specified by an orthonormal basis  $\{|m_i\rangle\}$  of states with definite values of  $M$ , together with the value  $m_i$  of  $M$  for each state.
- ▶ We can codify  $M$  into an operator by defining

$$\hat{M} = \sum_i m_i |m_i\rangle \langle m_i|.$$

- ▶ This operator has the following two properties:

# Self-adjoint operators and measurements

- ▶ Recall that a maximally specific measurement  $M$  is specified by an orthonormal basis  $\{|m_i\rangle\}$  of states with definite values of  $M$ , together with the value  $m_i$  of  $M$  for each state.
- ▶ We can codify  $M$  into an operator by defining

$$\hat{M} = \sum_i m_i |m_i\rangle \langle m_i|.$$

- ▶ This operator has the following two properties:
  1.  $\hat{M}$  is *self-adjoint*:  $\hat{M}^\dagger = \hat{M}$ . (**Exercise:** Show that an operator is self-adjoint iff its eigenvalues are all real.)

# Self-adjoint operators and measurements

- ▶ Recall that a maximally specific measurement  $M$  is specified by an orthonormal basis  $\{|m_i\rangle\}$  of states with definite values of  $M$ , together with the value  $m_i$  of  $M$  for each state.
- ▶ We can codify  $M$  into an operator by defining

$$\hat{M} = \sum_i m_i |m_i\rangle \langle m_i|.$$

- ▶ This operator has the following two properties:
  1.  $\hat{M}$  is *self-adjoint*:  $\hat{M}^\dagger = \hat{M}$ . (**Exercise:** Show that an operator is self-adjoint iff its eigenvalues are all real.)
  2.  $\hat{M}$  is non-degenerate. (This follows from the definition of a maximally specific measurement.)

# Self-adjoint operators and measurements

- ▶ Recall that a maximally specific measurement  $M$  is specified by an orthonormal basis  $\{|m_i\rangle\}$  of states with definite values of  $M$ , together with the value  $m_i$  of  $M$  for each state.
- ▶ We can codify  $M$  into an operator by defining

$$\hat{M} = \sum_i m_i |m_i\rangle \langle m_i|.$$

- ▶ This operator has the following two properties:
  1.  $\hat{M}$  is *self-adjoint*:  $\hat{M}^\dagger = \hat{M}$ . (**Exercise:** Show that an operator is self-adjoint iff its eigenvalues are all real.)
  2.  $\hat{M}$  is non-degenerate. (This follows from the definition of a maximally specific measurement.)
- ▶ Conversely, the spectral theorem means that every non-degenerate self-adjoint operator determines a maximally specific measurement.

# The orthodoxy on self-adjoint operators

- ▶ It's for these reasons that textbook quantum mechanics orthodoxy has it that *self-adjoint* operators are associated with the measurement of physical quantities.

# The orthodoxy on self-adjoint operators

- ▶ It's for these reasons that textbook quantum mechanics orthodoxy has it that *self-adjoint* operators are associated with the measurement of physical quantities.
- ▶ However, for a recent and philosophically rich article arguing that other normal operators could serve just as well (despite having complex eigenvalues!), see (Roberts 2017).

# The projection postulate

- ▶ The reason that the operator representation of measurements is useful is the following theorem:

# The projection postulate

- ▶ The reason that the operator representation of measurements is useful is the following theorem:  
If  $\hat{X}$  is a self-adjoint operator, then the expected value of making the associated measurement on a state  $|\psi\rangle$  is  $\langle\psi|\hat{X}|\psi\rangle$ .  
**(Exercise: prove this.)**

# The projection postulate

- ▶ The reason that the operator representation of measurements is useful is the following theorem:  
If  $\hat{X}$  is a self-adjoint operator, then the expected value of making the associated measurement on a state  $|\psi\rangle$  is  $\langle\psi|\hat{X}|\psi\rangle$ .  
(**Exercise:** prove this.)
- ▶ The **projection postulate** is the postulate that after measurement a system will be found in one of the eigenstates of the quantity being measured.

# The projection postulate

- ▶ The reason that the operator representation of measurements is useful is the following theorem:  
If  $\hat{X}$  is a self-adjoint operator, then the expected value of making the associated measurement on a state  $|\psi\rangle$  is  $\langle\psi|\hat{X}|\psi\rangle$ .  
**(Exercise: prove this.)**
- ▶ The **projection postulate** is the postulate that after measurement a system will be found in one of the eigenstates of the quantity being measured.
- ▶ This has to be treated with some care, and is *wrong* in its simplest form: in general, measurements disrupt the system being measured.

# The projection postulate

- ▶ The reason that the operator representation of measurements is useful is the following theorem:  
If  $\hat{X}$  is a self-adjoint operator, then the expected value of making the associated measurement on a state  $|\psi\rangle$  is  $\langle\psi|\hat{X}|\psi\rangle$ .

(**Exercise:** prove this.)

- ▶ The **projection postulate** is the postulate that after measurement a system will be found in one of the eigenstates of the quantity being measured.
- ▶ This has to be treated with some care, and is *wrong* in its simplest form: in general, measurements disrupt the system being measured.
  - ▶ For example, a photon detector destroys a photon completely when measuring its location!

# Commuting measurements

- ▶ Suppose I know how to perform a measurement of  $\hat{A}$ .

# Commuting measurements

- ▶ Suppose I know how to perform a measurement of  $\hat{A}$ .
- ▶ I can immediately, and with no further work, perform a measurement of  $f(\hat{A})$ : all I do is measure  $\hat{A}$  and apply the function  $f$  to the result.

# Commuting measurements

- ▶ Suppose I know how to perform a measurement of  $\hat{A}$ .
- ▶ I can immediately, and with no further work, perform a measurement of  $f(\hat{A})$ : all I do is measure  $\hat{A}$  and apply the function  $f$  to the result.
- ▶ Now suppose that  $\hat{A}$  and  $\hat{B}$  are commuting self-adjoint operators. Then we already know that there is some  $\hat{C}$  s.t.  $\hat{A}$  and  $\hat{B}$  are functions of  $\hat{C}$ .

# Commuting measurements

- ▶ Suppose I know how to perform a measurement of  $\hat{A}$ .
- ▶ I can immediately, and with no further work, perform a measurement of  $f(\hat{A})$ : all I do is measure  $\hat{A}$  and apply the function  $f$  to the result.
- ▶ Now suppose that  $\hat{A}$  and  $\hat{B}$  are commuting self-adjoint operators. Then we already know that there is some  $\hat{C}$  s.t.  $\hat{A}$  and  $\hat{B}$  are functions of  $\hat{C}$ .
- ▶ So if we measure  $\hat{C}$  then we automatically measure  $\hat{A}$  and  $\hat{B}$ , just by applying the appropriate functions to the outcome.

# Commuting measurements

- ▶ Suppose I know how to perform a measurement of  $\hat{A}$ .
- ▶ I can immediately, and with no further work, perform a measurement of  $f(\hat{A})$ : all I do is measure  $\hat{A}$  and apply the function  $f$  to the result.
- ▶ Now suppose that  $\hat{A}$  and  $\hat{B}$  are commuting self-adjoint operators. Then we already know that there is some  $\hat{C}$  s.t.  $\hat{A}$  and  $\hat{B}$  are functions of  $\hat{C}$ .
- ▶ So if we measure  $\hat{C}$  then we automatically measure  $\hat{A}$  and  $\hat{B}$ , just by applying the appropriate functions to the outcome.
- ▶ We conclude that *commuting operators can be measured simultaneously*.

# Commuting measurements

- ▶ Suppose I know how to perform a measurement of  $\hat{A}$ .
- ▶ I can immediately, and with no further work, perform a measurement of  $f(\hat{A})$ : all I do is measure  $\hat{A}$  and apply the function  $f$  to the result.
- ▶ Now suppose that  $\hat{A}$  and  $\hat{B}$  are commuting self-adjoint operators. Then we already know that there is some  $\hat{C}$  s.t.  $\hat{A}$  and  $\hat{B}$  are functions of  $\hat{C}$ .
- ▶ So if we measure  $\hat{C}$  then we automatically measure  $\hat{A}$  and  $\hat{B}$ , just by applying the appropriate functions to the outcome.
- ▶ We conclude that *commuting operators can be measured simultaneously*.
- ▶ Conversely, if two operators don't commute then it's at best unclear what it would mean to measure them simultaneously.

# Unitary operators

- ▶ We have seen that self-adjoint operators can be defined as those satisfying  $\hat{A}^\dagger = \hat{A}$ , or equivalently those whose eigenvalues are real.

# Unitary operators

- ▶ We have seen that self-adjoint operators can be defined as those satisfying  $\hat{A}^\dagger = \hat{A}$ , or equivalently those whose eigenvalues are real.
- ▶ Similarly, *unitary operators* can be defined either as (i) operators satisfying  $\hat{A}^\dagger \hat{A} = \hat{1}$ , or (ii) operators whose eigenvalues all have modulus 1 (so of the form  $e^{i\theta}$ ).

# Unitary operators

- ▶ We have seen that self-adjoint operators can be defined as those satisfying  $\hat{A}^\dagger = \hat{A}$ , or equivalently those whose eigenvalues are real.
- ▶ Similarly, *unitary operators* can be defined either as (i) operators satisfying  $\hat{A}^\dagger \hat{A} = \hat{1}$ , or (ii) operators whose eigenvalues all have modulus 1 (so of the form  $e^{i\theta}$ ).
- ▶ The significance of unitary operators is that they map physical states to physical states, for if a map is to do this then it must preserve normalisation: if  $|\psi\rangle$  is a physical state, then  $|\psi'\rangle = \hat{U}|\psi\rangle$  is a physical state only if

$$1 = \langle \psi' | \psi' \rangle = \langle \psi | \hat{U}^\dagger \hat{U} | \psi \rangle,$$

which is true for all states only if  $\hat{U}^\dagger \hat{U} = \hat{1}$ .

# Significance of unitary operators

There are two important sorts of transformation described by unitary operators:

# Significance of unitary operators

There are two important sorts of transformation described by unitary operators:

1. *Time evolution*: It's a postulate of textbook quantum mechanics that states evolve linearly (in accordance with the Schrödinger equation, to be discussed later): so if  $|\psi\rangle$  evolves into  $|\psi'\rangle$  and  $|\phi\rangle$  evolves into  $|\phi'\rangle$ , then  $\alpha|\psi\rangle + \beta|\phi\rangle$  evolves into  $\alpha|\psi'\rangle + \beta|\phi'\rangle$ . This means that time evolution is described by unitary operators: for any times  $t, t_0$ , there must be a unitary operator  $\hat{U}(t, t_0)$  which evolves states at time  $t_0$  into states at time  $t$ :

$$|\psi(t)\rangle = \hat{U}(t, t_0) |\psi(t_0)\rangle.$$

# Significance of unitary operators

There are two important sorts of transformation described by unitary operators:

1. *Time evolution*: It's a postulate of textbook quantum mechanics that states evolve linearly (in accordance with the Schrödinger equation, to be discussed later): so if  $|\psi\rangle$  evolves into  $|\psi'\rangle$  and  $|\phi\rangle$  evolves into  $|\phi'\rangle$ , then  $\alpha|\psi\rangle + \beta|\phi\rangle$  evolves into  $\alpha|\psi'\rangle + \beta|\phi'\rangle$ . This means that time evolution is described by unitary operators: for any times  $t, t_0$ , there must be a unitary operator  $\hat{U}(t, t_0)$  which evolves states at time  $t_0$  into states at time  $t$ :

$$|\psi(t)\rangle = \hat{U}(t, t_0) |\psi(t_0)\rangle .$$

2. *Symmetry transformations*: Symmetries of quantum systems, such as rotations or translations, are described by (anti-)unitary operators. (See e.g. Weinberg 1995)

# The Schrödinger equation

Consider a very short time evolution:

$$|\psi(t_0 + \delta t)\rangle = \hat{U}(t_0 + \delta t, t_0) |\psi(t_0)\rangle.$$

# The Schrödinger equation

Consider a very short time evolution:

$$|\psi(t_0 + \delta t)\rangle = \hat{U}(t_0 + \delta t, t_0) |\psi(t_0)\rangle.$$

Then in the limit  $\delta t \rightarrow 0$ , we obtain

$$|\psi(t_0)\rangle + \frac{d}{dt} |\psi(t_0)\rangle \delta t = \left( \hat{1} + \frac{d}{dt} \hat{U}(t, t_0) \right) |\psi(t_0)\rangle.$$

# The Schrödinger equation

Consider a very short time evolution:

$$|\psi(t_0 + \delta t)\rangle = \hat{U}(t_0 + \delta t, t_0) |\psi(t_0)\rangle.$$

Then in the limit  $\delta t \rightarrow 0$ , we obtain

$$|\psi(t_0)\rangle + \frac{d}{dt} |\psi(t_0)\rangle \delta t = \left( \hat{1} + \frac{d}{dt} \hat{U}(t, t_0) \right) |\psi(t_0)\rangle.$$

Define the (self-adjoint) operator

$$\hat{H}(t_0) = i\hbar \frac{d}{dt} \Big|_{t_0} \hat{U}(t, t_0),$$

# The Schrödinger equation

Consider a very short time evolution:

$$|\psi(t_0 + \delta t)\rangle = \hat{U}(t_0 + \delta t, t_0) |\psi(t_0)\rangle.$$

Then in the limit  $\delta t \rightarrow 0$ , we obtain

$$|\psi(t_0)\rangle + \frac{d}{dt} |\psi(t_0)\rangle \delta t = \left( \hat{1} + \frac{d}{dt} \hat{U}(t, t_0) \right) |\psi(t_0)\rangle.$$

Define the (self-adjoint) operator

$$\hat{H}(t_0) = i\hbar \left. \frac{d}{dt} \right|_{t_0} \hat{U}(t, t_0),$$

We obtain:

$$\frac{d}{dt} |\psi(t)\rangle = -\frac{i}{\hbar} \hat{H}(t) |\psi(t)\rangle,$$

which is the time-dependent Schrödinger equation.

# Today

Quantum states

Hilbert spaces

More about Hilbert spaces

Linear operators

The physical significance of operators

**Wavefunctions and infinite-dimensional Hilbert spaces**

# Infinite-dimensional Hilbert spaces

- ▶ Recall that the dimension of a Hilbert space is the number of vectors in a basis for that space.

# Infinite-dimensional Hilbert spaces

- ▶ Recall that the dimension of a Hilbert space is the number of vectors in a basis for that space.
- ▶ In general, this could be infinite!

# Infinite-dimensional Hilbert spaces

- ▶ Recall that the dimension of a Hilbert space is the number of vectors in a basis for that space.
- ▶ In general, this could be infinite!
- ▶ Consider e.g., the energy levels of an atom—here, there must be an infinity of eigenstates.

# Infinite-dimensional Hilbert spaces

- ▶ Recall that the dimension of a Hilbert space is the number of vectors in a basis for that space.
- ▶ In general, this could be infinite!
- ▶ Consider e.g., the energy levels of an atom—here, there must be an infinity of eigenstates.
- ▶ To represent these systems, we need to make use of *infinite-dimensional* Hilbert spaces.

# Infinite-dimensional Hilbert spaces

- ▶ Recall that the dimension of a Hilbert space is the number of vectors in a basis for that space.
- ▶ In general, this could be infinite!
- ▶ Consider e.g., the energy levels of an atom—here, there must be an infinity of eigenstates.
- ▶ To represent these systems, we need to make use of *infinite-dimensional* Hilbert spaces.
- ▶ The only reason why we can often get away with using finite-dimensional Hilbert spaces is that we abstract away some of the system's degrees of freedom.

# Infinite-dimensional Hilbert spaces

- ▶ Recall that the dimension of a Hilbert space is the number of vectors in a basis for that space.
- ▶ In general, this could be infinite!
- ▶ Consider e.g., the energy levels of an atom—here, there must be an infinity of eigenstates.
- ▶ To represent these systems, we need to make use of *infinite-dimensional* Hilbert spaces.
- ▶ The only reason why we can often get away with using finite-dimensional Hilbert spaces is that we abstract away some of the system's degrees of freedom.
- ▶ Mathematically, infinite-dimensional Hilbert spaces are difficult to work with! We'll follow a 'physics approach' (i.e., generally ignore the problems).

# Infinite-dimensional Hilbert spaces

- ▶ Recall that the dimension of a Hilbert space is the number of vectors in a basis for that space.
- ▶ In general, this could be infinite!
- ▶ Consider e.g., the energy levels of an atom—here, there must be an infinity of eigenstates.
- ▶ To represent these systems, we need to make use of *infinite-dimensional* Hilbert spaces.
- ▶ The only reason why we can often get away with using finite-dimensional Hilbert spaces is that we abstract away some of the system's degrees of freedom.
- ▶ Mathematically, infinite-dimensional Hilbert spaces are difficult to work with! We'll follow a 'physics approach' (i.e., generally ignore the problems).
- ▶ (For more, see the seminal (von Neumann 1955).)

# Operators with continuous spectra

- ▶ Not all observables have discrete spectra—consider e.g. position or momentum.

# Operators with continuous spectra

- ▶ Not all observables have discrete spectra—consider e.g. position or momentum.
- ▶ We represent these observables with *continuous* self-adjoint operators: operators whose spectra include all real numbers in some range, or perhaps just all real numbers full stop.

# Operators with continuous spectra

- ▶ Not all observables have discrete spectra—consider e.g. position or momentum.
- ▶ We represent these observables with *continuous* self-adjoint operators: operators whose spectra include all real numbers in some range, or perhaps just all real numbers full stop.
- ▶ E.g., the spectral representation of the position operator is

$$\hat{X} = \int_{-\infty}^{+\infty} dx x |x\rangle \langle x|.$$

# Operators with continuous spectra

- ▶ Not all observables have discrete spectra—consider e.g. position or momentum.
- ▶ We represent these observables with *continuous* self-adjoint operators: operators whose spectra include all real numbers in some range, or perhaps just all real numbers full stop.
- ▶ E.g., the spectral representation of the position operator is

$$\hat{X} = \int_{-\infty}^{+\infty} dx x |x\rangle \langle x|.$$

- ▶ Then the probability rule for finding some quantum system in (for this case) some position interval is

$$\Pr(x_1 < x < x_2) = \int_{x_1}^{x_2} dx |\langle x|\psi\rangle|^2.$$

# Operators with continuous spectra

- ▶ Not all observables have discrete spectra—consider e.g. position or momentum.
- ▶ We represent these observables with *continuous* self-adjoint operators: operators whose spectra include all real numbers in some range, or perhaps just all real numbers full stop.
- ▶ E.g., the spectral representation of the position operator is

$$\hat{X} = \int_{-\infty}^{+\infty} dx x |x\rangle \langle x|.$$

- ▶ Then the probability rule for finding some quantum system in (for this case) some position interval is

$$\Pr(x_1 < x < x_2) = \int_{x_1}^{x_2} dx |\langle x|\psi\rangle|^2.$$

- ▶ The *wavefunction* is defined as  $\psi(x) := \langle x|\psi\rangle$ .

# Ontology of the wavefunction

- ▶ It's tempting to think of the wavefunction of a particle as a field on space.

# Ontology of the wavefunction

- ▶ It's tempting to think of the wavefunction of a particle as a field on space.
- ▶ After all, it's a function of three coordinates, labelled 'positions', and when it is concentrated in some region, the probability is very high that a measurement of position will give a result in that region.

# Ontology of the wavefunction

- ▶ It's tempting to think of the wavefunction of a particle as a field on space.
- ▶ After all, it's a function of three coordinates, labelled 'positions', and when it is concentrated in some region, the probability is very high that a measurement of position will give a result in that region.
- ▶ But this is a mistake! Consider the wavefunction for *two* particles: this must be a (complex-valued) function of *six* variables: three for the position of the first particle, and three for the position of the second.

# Ontology of the wavefunction

- ▶ It's tempting to think of the wavefunction of a particle as a field on space.
- ▶ After all, it's a function of three coordinates, labelled 'positions', and when it is concentrated in some region, the probability is very high that a measurement of position will give a result in that region.
- ▶ But this is a mistake! Consider the wavefunction for *two* particles: this must be a (complex-valued) function of *six* variables: three for the position of the first particle, and three for the position of the second.
- ▶ So, the wavefunction can't straightforwardly be thought of as living on space at all.

# Ontology of the wavefunction

- ▶ It's tempting to think of the wavefunction of a particle as a field on space.
- ▶ After all, it's a function of three coordinates, labelled 'positions', and when it is concentrated in some region, the probability is very high that a measurement of position will give a result in that region.
- ▶ But this is a mistake! Consider the wavefunction for *two* particles: this must be a (complex-valued) function of *six* variables: three for the position of the first particle, and three for the position of the second.
- ▶ So, the wavefunction can't straightforwardly be thought of as living on space at all.
- ▶ Rather, the wavefunction is a complex-valued field on *configuration space*, which is the space of all possible coordinates of all particles, with  $3N$  dimensions for an  $N$ -particle system.

# Summary for today

- ▶ Today, I've introduced:
  1. Quantum states
  2. Hilbert spaces
  3. Operators and their connections with measurements
  4. Wavefunctions

# Summary for today

- ▶ Today, I've introduced:
  1. Quantum states
  2. Hilbert spaces
  3. Operators and their connections with measurements
  4. Wavefunctions
- ▶ Tomorrow, we'll look at *density operators* and *entanglement*.

# References

-  Shan Gao, “Why the Global Phase is Not Real”, *Foundations of Physics* 54:19, 2024.
-  Bryan W. Roberts, “Unreal Observables”, *Philosophy of Science* 84, pp. 1265–74, 2017.
-  John von Neumann, *Mathematical Foundations of Quantum Mechanics*, Princeton, NJ: Princeton University Press, 1955.
-  David Wallace, “On the Reality of the Global Phase”, 2022.
-  Steven Weinberg, *The Quantum Theory of Fields*, volume 1, Cambridge: Cambridge University Press, 1995.