

# IPP-QM-11: Contextuality

James Read<sup>1</sup>

<sup>1</sup>Faculty of Philosophy, University of Oxford, UK, OX2 6GG

MT24

# The course

1. Basic quantum formalism
2. Density operators and entanglement
3. Decoherence
4. The measurement problem
5. Dynamical collapse theories
6. Bohmian mechanics
7. Everettian structure
8. Everettian probability
9. EPR and Bell's theorem
10. The Bell-CHSH inequalities and possible responses
11. Contextuality
12. The PBR theorem
13. Quantum logic
14. Pragmatism and QBism
15. Relational quantum mechanics
16. Wavefunction realism

# Today

## Contextuality

The Bell-Kochen-Specker theorem introduced

The Clifton-Stairs state-dependent proof

Proof of the Bell-Kochen-Specker theorem

The Klyachko-Can-Binicioglu-Shumovsky inequality

Coda: Gleason's theorem

# Today

## Contextuality

The Bell-Kochen-Specker theorem introduced

The Clifton-Stairs state-dependent proof

Proof of the Bell-Kochen-Specker theorem

The Klyachko-Can-Binicioglu-Shumovsky inequality

Coda: Gleason's theorem



# Contextuality

What does it mean for a physical theory to be *contextual*?

# Contextuality

What does it mean for a physical theory to be *contextual*?

- ▶ The outcome of a measurement is associated with the *whole experimental arrangement*.

# Contextuality

What does it mean for a physical theory to be *contextual*?


- ▶ The outcome of a measurement is associated with the *whole experimental arrangement*.
- ▶ Different experimental arrangements lead to *different* physics.

# Contextuality


What does it mean for a physical theory to be *contextual*?

- ▶ The outcome of a measurement is associated with the *whole experimental arrangement*.
- ▶ Different experimental arrangements lead to *different* physics.
- ▶ So measurements are not *simply* revealing properties of the system being measured.

# Contextuality reminder: Bohmian mechanics




# Contextuality reminder: Bohmian mechanics



- The Bohm trajectories are deterministic. The outcomes of a measurement are fixed in advance.

# Contextuality reminder: Bohmian mechanics



- ▶ The Bohm trajectories are deterministic. The outcomes of a measurement are fixed in advance.
- ▶ But the outcomes of e.g. Stern-Gerlach experiments also depend upon the context of *how* the measurement is performed.

# Note on terminology

There's some fiddly terminology here which is worth clarifying:



# Note on terminology

There's some fiddly terminology here which is worth clarifying:

**Outcome determinism:** The measurement reveals a pre-existing property of the system.

# Note on terminology

There's some fiddly terminology here which is worth clarifying:

**Outcome determinism:** The measurement reveals a pre-existing property of the system.

**Measurement non-contextuality:** The property does not depend upon how the measurement is performed.

# Note on terminology

There's some fiddly terminology here which is worth clarifying:

**Outcome determinism:** The measurement reveals a pre-existing property of the system.

**Measurement non-contextuality:** The property does not depend upon how the measurement is performed.

Historical notes on terminology:

# Note on terminology

There's some fiddly terminology here which is worth clarifying:

**Outcome determinism:** The measurement reveals a pre-existing property of the system.

**Measurement non-contextuality:** The property does not depend upon how the measurement is performed.

Historical notes on terminology:

- ▶ 'Outcome determinism' is sometimes known as 'value definiteness'.

# Note on terminology

There's some fiddly terminology here which is worth clarifying:

**Outcome determinism:** The measurement reveals a pre-existing property of the system.

**Measurement non-contextuality:** The property does not depend upon how the measurement is performed.

Historical notes on terminology:

- ▶ 'Outcome determinism' is sometimes known as 'value definiteness'.
- ▶ The conjunction of 'outcome determinism' with 'measurement non-contextuality' used to be called 'non-contextuality' or 'non-contextual value definiteness.'

# Note on terminology

There's some fiddly terminology here which is worth clarifying:

**Outcome determinism:** The measurement reveals a pre-existing property of the system.

**Measurement non-contextuality:** The property does not depend upon how the measurement is performed.

Historical notes on terminology:

- ▶ 'Outcome determinism' is sometimes known as 'value definiteness'.
- ▶ The conjunction of 'outcome determinism' with 'measurement non-contextuality' used to be called 'non-contextuality' or 'non-contextual value definiteness.'
- ▶ Now it is sometimes called **traditional non-contextuality** or 'Kochen-Specker non-contextuality'.

# Non-contextuality no-go theorems

- ▶ Traditionally non-contextual theories are possible in 2D Hilbert spaces (e.g. the ‘Kochen-Specker qbit’).

# Non-contextuality no-go theorems

- ▶ Traditionally non-contextual theories are possible in 2D Hilbert spaces (e.g. the 'Kochen-Specker qbit').
- ▶ They are provably not possible in any Hilbert space of larger dimension:



# Non-contextuality no-go theorems

- ▶ Traditionally non-contextual theories are possible in 2D Hilbert spaces (e.g. the ‘Kochen-Specker qbit’).
- ▶ They are provably not possible in any Hilbert space of larger dimension:
  - ▶ The Clifton-Stairs state-dependent proof.
  - ▶ The Bell-Kochen-Specker (BKS) state-independent proof.
  - ▶ The Klyachko-Can-Binicioglu-Shumovsky (KCBS) inequality.

# Today

Contextuality

The Bell-Kochen-Specker theorem introduced

The Clifton-Stairs state-dependent proof

Proof of the Bell-Kochen-Specker theorem

The Klyachko-Can-Binicioglu-Shumovsky inequality

Coda: Gleason's theorem

# The Bell-Kochen-Specker theorem

- ▶ The BKS theorem tells us that any hidden variable theory which assigns values to all properties represented by projectors must be *contextual*.

# The Bell-Kochen-Specker theorem

- ▶ The BKS theorem tells us that any hidden variable theory which assigns values to all properties represented by projectors must be *contextual*.
- ▶ Specifically, whether or not a system is found, on measurement, to possess a given property must depend upon which *other* properties are measured simultaneously.

# The Bell-Kochen-Specker theorem

- ▶ The BKS theorem tells us that any hidden variable theory which assigns values to all properties represented by projectors must be *contextual*.
- ▶ Specifically, whether or not a system is found, on measurement, to possess a given property must depend upon which *other* properties are measured simultaneously.
- ▶ As Wallace (2007, p. 51) writes, “Contextuality seems well-nigh inconsistent with the idea that systems determinately do or do not possess given properties and that measurements simply determine whether or not they do.”

# Today

Contextuality

The Bell-Kochen-Specker theorem introduced

**The Clifton-Stairs state-dependent proof**

Proof of the Bell-Kochen-Specker theorem

The Klyachko-Can-Binicioglu-Shumovsky inequality

Coda: Gleason's theorem

# The Clifton-Stairs state-dependent proof

Before getting to the full BKS contextuality theorem, we'll first look at a simpler, 'state-dependent' proof due to Clifton (1993) and Stairs (1992).

# The Clifton-Stairs proof

- In all diagrams to follow, let a circle be a projector onto a given state.



# The Clifton-Stairs proof

- ▶ In all diagrams to follow, let a circle be a projector onto a given state.
- ▶ Circles are connected by lines just in case their associated projectors are orthogonal.

# The Clifton-Stairs proof

- ▶ In all diagrams to follow, let a circle be a projector onto a given state.
- ▶ Circles are connected by lines just in case their associated projectors are orthogonal.
- ▶ The measurement scenario for the Clifton-Stairs proof is three mutually orthogonal projectors:

$$\begin{array}{ccc} \frac{1}{\sqrt{2}}(|1\rangle + |2\rangle) & \begin{array}{c} \text{---} \text{---} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \end{array} & \frac{1}{\sqrt{2}}(|1\rangle - |2\rangle) \\ & |3\rangle & \end{array}$$

$\epsilon_a(\phi_k|\lambda) \in \{0,1\}$

$\sum \epsilon_a(\phi_k|\lambda) = 1$

# The Clifton-Stairs proof

- ▶ In all diagrams to follow, let a circle be a projector onto a given state.
- ▶ Circles are connected by lines just in case their associated projectors are orthogonal.
- ▶ The measurement scenario for the Clifton-Stairs proof is three mutually orthogonal projectors:

$$\begin{array}{ccc} \frac{1}{\sqrt{2}}(|1\rangle + |2\rangle) & \begin{array}{c} \circ \text{---} \circ \\ \diagdown \quad \diagup \\ \circ \end{array} & \frac{1}{\sqrt{2}}(|1\rangle - |2\rangle) \\ & & \sum \epsilon_a(\phi_k|\lambda) = 1 \end{array}$$

$|3\rangle$

- ▶ Any triangle is a measurement scenario.

# The Clifton-Stairs proof

- ▶ In all diagrams to follow, let a circle be a projector onto a given state.
- ▶ Circles are connected by lines just in case their associated projectors are orthogonal.
- ▶ The measurement scenario for the Clifton-Stairs proof is three mutually orthogonal projectors:

$$\begin{array}{ccc} \frac{1}{\sqrt{2}}(|1\rangle + |2\rangle) & \begin{array}{c} \circ \text{---} \circ \\ \diagdown \quad \diagup \\ \circ \end{array} & \frac{1}{\sqrt{2}}(|1\rangle - |2\rangle) \\ & |3\rangle & \end{array}$$

$\epsilon_a(\phi_k|\lambda) \in \{0,1\}$   
 $\sum \epsilon_a(\phi_k|\lambda) = 1$

- ▶ Any triangle is a measurement scenario.
- ▶ We require that the measurement outcomes are 0 or 1, and that they sum to 1.

# The Clifton-Stairs proof

- ▶ In all diagrams to follow, let a circle be a projector onto a given state.
- ▶ Circles are connected by lines just in case their associated projectors are orthogonal.
- ▶ The measurement scenario for the Clifton-Stairs proof is three mutually orthogonal projectors:

$$\begin{array}{ccc} \frac{1}{\sqrt{2}}(|1\rangle + |2\rangle) & \begin{array}{c} \text{---} \text{---} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \end{array} & \frac{1}{\sqrt{2}}(|1\rangle - |2\rangle) \\ & & |3\rangle \end{array}$$


$\epsilon_a(\phi_k|\lambda) \in \{0,1\}$

$\sum \epsilon_a(\phi_k|\lambda) = 1$

- ▶ Any triangle is a measurement scenario.
- ▶ We require that the measurement outcomes are 0 or 1, and that they sum to 1.
- ▶ So for any triple of ontic states, one of those states is going to be 1 and the other two 0.

# The Clifton-Stairs proof

$$\frac{1}{\sqrt{2}}(|1\rangle + |2\rangle)$$



$$\frac{1}{\sqrt{2}}(|1\rangle - |2\rangle)$$

$$\epsilon_a(\phi_k|\lambda) \in \{0,1\}$$

$$\sum \epsilon_a(\phi_k|\lambda) = 1$$

# The Clifton-Stairs proof

- Now suppose that we have two measurement scenarios which contain the same projector.


# The Clifton-Stairs proof

- ▶ Now suppose that we have two measurement scenarios which contain the same projector.
- ▶ We link up the two triangles, and non-contextuality demands that if a circle is red in one measurement scenario then it had better be red in the other measurement scenario.



# The Clifton-Stairs proof

- Now suppose that we have two measurement scenarios which contain the same projector.
- We link up the two triangles, and non-contextuality demands that if a circle is red in one measurement scenario then it had better be red in the other measurement scenario.




$$\epsilon_a(\phi_k|\lambda) \in \{0,1\}$$

$$\sum \epsilon_a(\phi_k|\lambda) = 1$$

$$\epsilon_a(\phi_1|\lambda) = \epsilon_{a'}(\phi_1|\lambda)$$

# The Clifton-Stairs proof

So, naturally, the other circles are going to have to be blue:




$$\epsilon_a(\phi_k|\lambda) \in \{0,1\}$$

$$\sum \epsilon_a(\phi_k|\lambda) = 1$$

$$\epsilon_a(\phi_1|\lambda) = \epsilon_{a'}(\phi_1|\lambda)$$

# The Clifton-Stairs proof


Now consider a scenario like this:



One of the other two circles is going to have to be red...


# The Clifton-Stairs proof

E.g.:



# The Clifton-Stairs proof

Now consider this scenario:




Notice that the far-left and far-right states are *not orthogonal to each other*.

# The Clifton-Stairs proof

Suppose that we prepare the right-most state:

# The Clifton-Stairs proof

Suppose that we prepare the right-most state:



$$\int \epsilon_a(\Psi|\lambda) \mu_\Psi(\lambda) = 1 \quad |\Psi\rangle = \frac{1}{\sqrt{3}}(|1\rangle + |2\rangle + |3\rangle) \quad \epsilon_a(\phi_1|\lambda) = \epsilon_{a'}(\phi_1|\lambda)$$


# The Clifton-Stairs proof

- ▶ Since the leftmost state isn't orthogonal to the rightmost state, some of the ontic states which are prepared by the rightmost state will also be in the leftmost state.



# The Clifton-Stairs proof

- Since the leftmost state isn't orthogonal to the rightmost state, some of the ontic states which are prepared by the rightmost state will also be in the leftmost state.
- In that case, we have:




$$\int \epsilon_a(\Psi|\lambda) \mu_\Psi(\lambda) = 1$$

$$\int \epsilon_a(\Phi|\lambda) \mu_\Psi(\lambda) = \frac{1}{9}$$

$$|\Psi\rangle = \frac{1}{\sqrt{3}}(|1\rangle + |2\rangle + |3\rangle) \quad \epsilon_a(\phi_1|\lambda) = \epsilon_{a'}(\phi_1|\lambda)$$

$$\left| \langle \Psi | \frac{(|1\rangle - |2\rangle + |3\rangle)}{\sqrt{3}} \right|^2 = \frac{1}{9}$$

# The Clifton-Stairs proof




$$\int \epsilon_a(\Psi|\lambda) \mu_\Psi(\lambda) = 1$$

$$\int \epsilon_a(\Phi|\lambda) \mu_\Psi(\lambda) = \frac{1}{9}$$

$$|\Psi\rangle = \frac{1}{\sqrt{3}}(|1\rangle + |2\rangle + |3\rangle) \quad \epsilon_a(\phi_1|\lambda) = \epsilon_{a'}(\phi_1|\lambda)$$

$$\left| \langle \Psi | \frac{(|1\rangle - |2\rangle + |3\rangle)}{\sqrt{3}} \right|^2 = \frac{1}{9}$$

# The Clifton-Stairs proof




$$\int \epsilon_a(\Psi|\lambda) \mu_\Psi(\lambda) = 1$$

$$\int \epsilon_a(\Phi|\lambda) \mu_\Psi(\lambda) = \frac{1}{9}$$

$$|\Psi\rangle = \frac{1}{\sqrt{3}}(|1\rangle + |2\rangle + |3\rangle) \quad \epsilon_a(\phi_1|\lambda) = \epsilon_{a'}(\phi_1|\lambda)$$

$$\left| \langle \Psi | \frac{(|1\rangle - |2\rangle + |3\rangle)}{\sqrt{3}} \right|^2 = \frac{1}{9}$$

# The Clifton-Stairs proof




$$\int \epsilon_a(\Psi|\lambda) \mu_\Psi(\lambda) = 1$$

$$\int \epsilon_a(\Phi|\lambda) \mu_\Psi(\lambda) = \frac{1}{9}$$

$$|\Psi\rangle = \frac{1}{\sqrt{3}}(|1\rangle + |2\rangle + |3\rangle) \quad \epsilon_a(\phi_1|\lambda) = \epsilon_{a'}(\phi_1|\lambda)$$

$$\left| \langle \Psi | \frac{(|1\rangle - |2\rangle + |3\rangle)}{\sqrt{3}} \right|^2 = \frac{1}{9}$$

# The Clifton-Stairs proof



$$\int \epsilon_a(\Psi|\lambda) \mu_\Psi(\lambda) = 1$$

$$\int \epsilon_a(\Phi|\lambda) \mu_\Psi(\lambda) = \frac{1}{9}$$

$$|\Psi\rangle = \frac{1}{\sqrt{3}}(|1\rangle + |2\rangle + |3\rangle) \quad \epsilon_a(\phi_1|\lambda) = \epsilon_{a'}(\phi_1|\lambda)$$

$$\left| \langle \Psi | \frac{(|1\rangle - |2\rangle + |3\rangle)}{\sqrt{3}} \right|^2 = \frac{1}{9}$$

# State-dependent vs state-independent proofs

- ▶ So contradiction, the model fails.

# State-dependent vs state-independent proofs

- ▶ So contradiction, the model fails.
- ▶ The trick in completing quantum non-contextuality proofs like this lies in finding the right combination of states.

# State-dependent vs state-independent proofs

- ▶ So contradiction, the model fails.
- ▶ The trick in completing quantum non-contextuality proofs like this lies in finding the right combination of states.
- ▶ It really does bear stressing that the Clifton-Stairs proof is *state-dependent*:



# State-dependent vs state-independent proofs

- ▶ So contradiction, the model fails.
- ▶ The trick in completing quantum non-contextuality proofs like this lies in finding the right combination of states.
- ▶ It really does bear stressing that the Clifton-Stairs proof is *state-dependent*:
  - ▶ For any given quantum state, there are combinations of observables that can't all be traditionally non-contextual.

# State-dependent vs state-independent proofs

- ▶ So contradiction, the model fails.
- ▶ The trick in completing quantum non-contextuality proofs like this lies in finding the right combination of states.
- ▶ It really does bear stressing that the Clifton-Stairs proof is *state-dependent*:
  - ▶ For any given quantum state, there are combinations of observables that can't all be traditionally non-contextual.
  - ▶ Different quantum states require different observables.

# State-dependent vs state-independent proofs

- ▶ So contradiction, the model fails.
- ▶ The trick in completing quantum non-contextuality proofs like this lies in finding the right combination of states.
- ▶ It really does bear stressing that the Clifton-Stairs proof is *state-dependent*:
  - ▶ For any given quantum state, there are combinations of observables that can't all be traditionally non-contextual.
  - ▶ Different quantum states require different observables.
- ▶ One might desire a proof which doesn't mention states at all, but which just has to do with different combinations of projectors.

# State-dependent vs state-independent proofs

- ▶ So contradiction, the model fails.
- ▶ The trick in completing quantum non-contextuality proofs like this lies in finding the right combination of states.
- ▶ It really does bear stressing that the Clifton-Stairs proof is *state-dependent*:
  - ▶ For any given quantum state, there are combinations of observables that can't all be traditionally non-contextual.
  - ▶ Different quantum states require different observables.
- ▶ One might desire a proof which doesn't mention states at all, but which just has to do with different combinations of projectors.
- ▶ BKS proofs meet this desideratum:

# State-dependent vs state-independent proofs

- ▶ So contradiction, the model fails.
- ▶ The trick in completing quantum non-contextuality proofs like this lies in finding the right combination of states.
- ▶ It really does bear stressing that the Clifton-Stairs proof is *state-dependent*:
  - ▶ For any given quantum state, there are combinations of observables that can't all be traditionally non-contextual.
  - ▶ Different quantum states require different observables.
- ▶ One might desire a proof which doesn't mention states at all, but which just has to do with different combinations of projectors.
- ▶ BKS proofs meet this desideratum:
  - ▶ Contradiction arises for any quantum state.

# State-dependent vs state-independent proofs

- ▶ So contradiction, the model fails.
- ▶ The trick in completing quantum non-contextuality proofs like this lies in finding the right combination of states.
- ▶ It really does bear stressing that the Clifton-Stairs proof is *state-dependent*:
  - ▶ For any given quantum state, there are combinations of observables that can't all be traditionally non-contextual.
  - ▶ Different quantum states require different observables.
- ▶ One might desire a proof which doesn't mention states at all, but which just has to do with different combinations of projectors.
- ▶ BKS proofs meet this desideratum:
  - ▶ Contradiction arises for any quantum state.
  - ▶ Contradiction arises for any ontic state.

# Today

Contextuality

The Bell-Kochen-Specker theorem introduced


The Clifton-Stairs state-dependent proof

**Proof of the Bell-Kochen-Specker theorem**

The Klyachko-Can-Binicioglu-Shumovsky inequality


Coda: Gleason's theorem

# The BKS proof






# The BKS proof




# The BKS proof




(Note: Stairs was working after Kochen & Specker, and noticed that one could use the mini-diagrams for a state-dependent proof.)


# The BKS proof




# The BKS proof



# The BKS proof




# The BKS proof




(At least one of the blobs for each of the pairs at the bottom has to be red given that we have a blue at the top.)

# The BKS proof




# The BKS proof




(This follows because we've seen from Clifton-Stairs that we can't have red on both sides of one of the little constructions.)




## The BKS proof



## The BKS proof



# The BKS proof



Contradiction, because we've already proven that if we've got one of the mini-diagrams then we can't have red on both sides!

# On the BKS theorem

- ▶ The proof here is straightforward.

# On the BKS theorem

- ▶ The proof here is straightforward.
- ▶ Obviously, the ingenuity is in designing this enormous set of states (117 projectors!).

# On the BKS theorem

- ▶ The proof here is straightforward.
- ▶ Obviously, the ingenuity is in designing this enormous set of states (117 projectors!).
- ▶ Bell (1966) actually did something similar to Kochen & Specker (1967) one year earlier than them, but nobody realised until afterwards (hence now the ‘BKS theorem’).

# Later history of BKS-type results

- ▶ Since 1967, other sets of uncolourable directions have been discovered with fewer vectors.

# Later history of BKS-type results


- ▶ Since 1967, other sets of uncolourable directions have been discovered with fewer vectors.
- ▶ E.g., Peres (1991) found a set of 33 with cubic symmetry.



# Later history of BKS-type results

- ▶ Since 1967, other sets of uncolourable directions have been discovered with fewer vectors.
- ▶ E.g., Peres (1991) found a set of 33 with cubic symmetry.
- ▶ Penrose pointed out that Peres' set of 33 directions can be described as follows: take a cube and superimpose it with its 90-degree rotations about two perpendicular lines connecting its centre to the midpoints of an edge. Peres' directions point to the vertices and the centres of the faces and edges of the resulting set of three interpenetrating cubes. (Obviously...)

# The Penrose cube



(Escher, *Waterfall*, 1961)

# Today

Contextuality

The Bell-Kochen-Specker theorem introduced

The Clifton-Stairs state-dependent proof

Proof of the Bell-Kochen-Specker theorem

The Klyachko-Can-Binicioglu-Shumovsky inequality

Coda: Gleason's theorem

# The Klyachko-Can-Binicioglu-Shumovsky inequality

- ▶ The BKS theorem proves that non-contextual hidden variable theories can't exist.

# The Klyachko-Can-Binicioglu-Shumovsky inequality

- ▶ The BKS theorem proves that non-contextual hidden variable theories can't exist.
- ▶ But if we want to do an experimental *test*, we need to have an inequality.


# The Klyachko-Can-Binicioglu-Shumovsky inequality

- ▶ The BKS theorem proves that non-contextual hidden variable theories can't exist.
- ▶ But if we want to do an experimental *test*, we need to have an inequality.
- ▶ One can derive an inequality for non-contextual hidden variable theories in the following way.

# The Klyachko-Can-Binicioglu-Shumovsky inequality


- ▶ The BKS theorem proves that non-contextual hidden variable theories can't exist.
- ▶ But if we want to do an experimental *test*, we need to have an inequality.
- ▶ One can derive an inequality for non-contextual hidden variable theories in the following way.
- ▶ I'll first present this, before discussing its significance.

# The KCBS inequality







# The KCBS inequality




# The KCBS inequality




# The KCBS inequality




# The KCBS inequality




# The KCBS inequality




# The KCBS inequality



# The KCBS inequality




# The KCBS inequality






# The KCBS inequality



# The KCBS inequality



Main point: at least one of the outer points in the star must be red in a configuration like this.

# The KCBS inequality


- Take any quantum state.

# The KCBS inequality

- ▶ Take any quantum state.
- ▶ From the five measurement bases (and measure it on that quantum state),  $\Pr(\text{Outer} = R) \geq \frac{1}{5}$ .

# The KCBS inequality


Now here comes quantum theory:



For any state you like, you can create this diagram around it.

# The KCBS inequality

$$P(\text{Outer} = R) \geq \frac{1}{5}$$



$$|l_a\rangle = \sin\theta\cos\phi_a|1\rangle + \sin\theta\sin\phi_a|2\rangle + \cos\theta|3\rangle$$

$$|l_1\rangle, |l_5\rangle, |l_{1,5}\rangle$$

$$\cos\theta = \frac{1}{\sqrt[4]{5}}$$

$$\phi_a = \frac{4}{5}\pi a$$

$$|\langle 3|l_a\rangle|^2 = \cos^2\theta = \frac{1}{\sqrt{5}}$$

$$|\langle 3|l_{1,5}\rangle|^2 = 1 - \frac{2}{\sqrt{5}} = 0.1056... < \frac{1}{5}$$

# Experimental violation of the KCBS inequality

- ▶ According to quantum theory, for any measurement basis you pick,  $\Pr(\text{Outer} = R) < \frac{1}{5}$ .

# Experimental violation of the KCBS inequality

- ▶ According to quantum theory, for any measurement basis you pick,  $\Pr(\text{Outer} = R) < \frac{1}{5}$ .
- ▶ So non-contextual hidden variable theories are ruled out on the basis of experiments!



# Experimental violation of the KCBS inequality

- ▶ According to quantum theory, for any measurement basis you pick,  $\Pr(\text{Outer} = R) < \frac{1}{5}$ .
- ▶ So non-contextual hidden variable theories are ruled out on the basis of experiments!
- ▶ But the KCBS inequality invites some foundational questions:

# Experimental violation of the KCBS inequality

- ▶ According to quantum theory, for any measurement basis you pick,  $\Pr(\text{Outer} = R) < \frac{1}{5}$ .
- ▶ So non-contextual hidden variable theories are ruled out on the basis of experiments!
- ▶ But the KCBS inequality invites some foundational questions:
  1. How does the KCBS inequality (related to the BKS theorem) compare with the CHSH inequalities (related to Bell's theorem)?

# Experimental violation of the KCBS inequality

- ▶ According to quantum theory, for any measurement basis you pick,  $\Pr(\text{Outer} = R) < \frac{1}{5}$ .
- ▶ So non-contextual hidden variable theories are ruled out on the basis of experiments!
- ▶ But the KCBS inequality invites some foundational questions:
  1. How does the KCBS inequality (related to the BKS theorem) compare with the CHSH inequalities (related to Bell's theorem)?
  2. Since the BKS theorem is a logical impossibility proof, isn't the KCBS inequality somewhat unnecessary?

# Experimental violation of the KCBS inequality

- ▶ According to quantum theory, for any measurement basis you pick,  $\Pr(\text{Outer} = R) < \frac{1}{5}$ .
- ▶ So non-contextual hidden variable theories are ruled out on the basis of experiments!
- ▶ But the KCBS inequality invites some foundational questions:
  1. How does the KCBS inequality (related to the BKS theorem) compare with the CHSH inequalities (related to Bell's theorem)?
  2. Since the BKS theorem is a logical impossibility proof, isn't the KCBS inequality somewhat unnecessary?
  3. By contrast, doesn't the CHSH inequality look integral to Bell's theorem?

# Experimental violation of the KCBS inequality

- ▶ According to quantum theory, for any measurement basis you pick,  $\Pr(\text{Outer} = R) < \frac{1}{5}$ .
- ▶ So non-contextual hidden variable theories are ruled out on the basis of experiments!
- ▶ But the KCBS inequality invites some foundational questions:
  1. How does the KCBS inequality (related to the BKS theorem) compare with the CHSH inequalities (related to Bell's theorem)?
  2. Since the BKS theorem is a logical impossibility proof, isn't the KCBS inequality somewhat unnecessary?
  3. By contrast, doesn't the CHSH inequality look integral to Bell's theorem?
  4. In light of this, do the two inequalities have the same role to play in 'experimental metaphysics'?

# Experimental violation of the KCBS inequality

- ▶ According to quantum theory, for any measurement basis you pick,  $\Pr(\text{Outer} = R) < \frac{1}{5}$ .
- ▶ So non-contextual hidden variable theories are ruled out on the basis of experiments!
- ▶ But the KCBS inequality invites some foundational questions:
  1. How does the KCBS inequality (related to the BKS theorem) compare with the CHSH inequalities (related to Bell's theorem)?
  2. Since the BKS theorem is a logical impossibility proof, isn't the KCBS inequality somewhat unnecessary?
  3. By contrast, doesn't the CHSH inequality look integral to Bell's theorem?
  4. In light of this, do the two inequalities have the same role to play in 'experimental metaphysics'?
- ▶ The KCBS inequality at least helps us to *quantify* quantum non-contextuality (see e.g. Cabello 2013).

# Today

Contextuality

The Bell-Kochen-Specker theorem introduced

The Clifton-Stairs state-dependent proof

Proof of the Bell-Kochen-Specker theorem

The Klyachko-Can-Binicioglu-Shumovsky inequality

Coda: Gleason's theorem

# Gleason's theorem

## Theorem (Gleason, 1957)

*Let  $f$  be any function on projectors on a Hilbert space  $\mathcal{H}$  of dimension  $d > 2$  to the unit interval which is additive for any set of pairwise disjoint projectors on  $\mathcal{H}$ . Then there exists a unique density matrix  $\hat{\rho}$  such that for any  $\hat{P}$  on  $\mathcal{H}$ ,  $f(\hat{P}) = \text{Tr}(\hat{\rho}\hat{P})$ .*



# Gleason's theorem

## Theorem (Gleason, 1957)

*Let  $f$  be any function on projectors on a Hilbert space  $\mathcal{H}$  of dimension  $d > 2$  to the unit interval which is additive for any set of pairwise disjoint projectors on  $\mathcal{H}$ . Then there exists a unique density matrix  $\hat{\rho}$  such that for any  $\hat{P}$  on  $\mathcal{H}$ ,  $f(\hat{P}) = \text{Tr}(\hat{\rho}\hat{P})$ .*

- (The proof of Gleason's theorem is notoriously involved and I won't present it here.)

# Gleason's theorem

## Theorem (Gleason, 1957)

*Let  $f$  be any function on projectors on a Hilbert space  $\mathcal{H}$  of dimension  $d > 2$  to the unit interval which is additive for any set of pairwise disjoint projectors on  $\mathcal{H}$ . Then there exists a unique density matrix  $\hat{\rho}$  such that for any  $\hat{P}$  on  $\mathcal{H}$ ,  $f(\hat{P}) = \text{Tr}(\hat{\rho}\hat{P})$ .*

- ▶ (The proof of Gleason's theorem is notoriously involved and I won't present it here.)
- ▶ One might think that Gleason's theorem counts as a derivation of the Born rule in unitary quantum mechanics—thereby (e.g.) solving (?) Everettians' worries about probability without the need to invoke decision theory etc.

# Doubts about Gleason's theorem and quantum probabilities

However, as Saunders points out,

*Gleason's theorem is a derivation of part of the Born rule, but of course it says nothing about 'measurements' or 'experiments'; nor, on reflection, is the premise of the theorem so clearly motivated. (Saunders 2005, p. 213)*

# Doubts about Gleason's theorem and quantum probabilities

However, as Saunders points out,

*Gleason's theorem is a derivation of part of the Born rule, but of course it says nothing about 'measurements' or 'experiments'; nor, on reflection, is the premise of the theorem so clearly motivated.* (Saunders 2005, p. 213)

**Question:** To what extent really is Gleason's theorem significant in the philosophy of quantum probabilities?

# Gleason's theorem and the BKS theorem

- ▶ A direct consequence of Gleason's theorem is that when the Hilbert space  $\mathcal{H}$  has dimension  $d \geq 3$  then there does not exist a function  $f$  that assigns only the values 0 or 1 to projections  $\hat{P} : \mathcal{H} \rightarrow \mathcal{H}$ .

# Gleason's theorem and the BKS theorem

- ▶ A direct consequence of Gleason's theorem is that when the Hilbert space  $\mathcal{H}$  has dimension  $d \geq 3$  then there does not exist a function  $f$  that assigns only the values 0 or 1 to projections  $\hat{P} : \mathcal{H} \rightarrow \mathcal{H}$ .
- ▶ Loosely speaking, this means that it is not possible to assign true/false values to all elements of the Hilbert space that represent yes/no events (projections) in such a way that is compatible with their Boolean properties.

# Gleason's theorem and the BKS theorem

- ▶ A direct consequence of Gleason's theorem is that when the Hilbert space  $\mathcal{H}$  has dimension  $d \geq 3$  then there does not exist a function  $f$  that assigns only the values 0 or 1 to projections  $\hat{P} : \mathcal{H} \rightarrow \mathcal{H}$ .
- ▶ Loosely speaking, this means that it is not possible to assign true/false values to all elements of the Hilbert space that represent yes/no events (projections) in such a way that is compatible with their Boolean properties.
- ▶ The BKS theorem is similar in spirit; indeed, the BKS theorem is provably a special case of Gleason's theorem.

# Gleason's theorem and the BKS theorem

- ▶ A direct consequence of Gleason's theorem is that when the Hilbert space  $\mathcal{H}$  has dimension  $d \geq 3$  then there does not exist a function  $f$  that assigns only the values 0 or 1 to projections  $\hat{P} : \mathcal{H} \rightarrow \mathcal{H}$ .
- ▶ Loosely speaking, this means that it is not possible to assign true/false values to all elements of the Hilbert space that represent yes/no events (projections) in such a way that is compatible with their Boolean properties.
- ▶ The BKS theorem is similar in spirit; indeed, the BKS theorem is provably a special case of Gleason's theorem.
- ▶ Therefore, both theorems rule out non-contextual hidden variable theories (though, being strictly stronger, Gleason's theorem does more).



# Summary

Today, I've:

# Summary

Today, I've:

1. Introduced quantum contextuality.

# Summary

Today, I've:

1. Introduced quantum contextuality.
2. Presented the Clifton-Stairs state-dependent no-go theorem on quantum non-contextuality, and the BKS state-independent theorem.

# Summary

Today, I've:

1. Introduced quantum contextuality.
2. Presented the Clifton-Stairs state-dependent no-go theorem on quantum non-contextuality, and the BKS state-independent theorem.
3. Presented the KCBS inequality, related to experimental tests of and bounds on quantum contextuality.

# Summary

Today, I've:

1. Introduced quantum contextuality.
2. Presented the Clifton-Stairs state-dependent no-go theorem on quantum non-contextuality, and the BKS state-independent theorem.
3. Presented the KCBS inequality, related to experimental tests of and bounds on quantum contextuality.
4. Introduced Gleason's theorem and indicated briefly how it relates to the BKS theorem.







# Summary

Today, I've:





1. Introduced quantum contextuality.
2. Presented the Clifton-Stairs state-dependent no-go theorem on quantum non-contextuality, and the BKS state-independent theorem.
3. Presented the KCBS inequality, related to experimental tests of and bounds on quantum contextuality.
4. Introduced Gleason's theorem and indicated briefly how it relates to the BKS theorem.

Next time:  $\Psi$ -ontic and  $\Psi$ -epistemic theories and the PBR theorem.

# References I

-  John S. Bell, “On the Problem of Hidden Variables in Quantum Mechanics”, *Reviews of Modern Physics* 38, pp. 447–52, 1966.
-  Adán Cabello, “Simple Explanation of the Quantum Violation of a Fundamental Inequality”, *Physical Review Letters* 110, 060402, 2013.
-  Rob Clifton, “Getting Contextual and Nonlocal Elements-of-Reality the Easy Way”, *American Journal of Physics* 61, pp. 443–47, 1993.
-  A. M. Gleason, “Measures on the Closed Subspaces of a Hilbert Space”, *Journal of Mathematics and Mechanics* 6, pp. 885–93, 1957.
-  Alexander A. Klyachko, M. Ali Can, Sinem Binicioglu, and Alexander S. Shumovsky, “Simple Test for Hidden Variables in Spin-1 Systems”, *Physical Review Letters* 101, 020403, 2008.
-  S. Kochen and E. Specker, “The Problem of Hidden Variables in Quantum Mechanics”, *Journal of Mathematics and Mechanics* 17, pp. 59–87, 1967.

# References II

-  N. David Mermin, “Hidden Variables and the Two Theorems of John Bell”, *Reviews of Modern Physics* 65(3), 1993.
-  A. Peres, “Two Simple Proofs of the Kochen-Specker Theorem”, *Journal of Physics A: Mathematical and General* 24(4), pp. L175–L178, 1991.
-  Simon Saunders, “What is Probability?” in A. Elitzur, S. Dolev and N. Kolenda (eds.), *Quo Vadis Quantum Mechanics?*, Berlin: Springer, 2005.
-  A. Stairs, “Value Definiteness and Contextualism: Cut and Paste with Hilbert Space”, *PSA* 1992 pp. 91–103, 1992.