

IPP-QM-5: Dynamical collapse theories

James Read¹

¹Faculty of Philosophy, University of Oxford, UK, OX2 6GG

MT24

The course

1. Basic quantum formalism
2. Density operators and entanglement
3. Decoherence
4. The measurement problem
5. Dynamical collapse theories
6. Bohmian mechanics
7. Everettian structure
8. Everettian probability
9. EPR and Bell's theorem
10. The Bell-CHSH inequalities and possible responses
11. Contextuality
12. The PBR theorem
13. Quantum logic
14. Pragmatism and QBism
15. Relational quantum mechanics
16. Wavefunction realism

Today

Dynamical collapse theories

The GRW theory

Objections to GRW

Modifications/alternatives to GRW

Today

Dynamical collapse theories

The GRW theory

Objections to GRW

Modifications/alternatives to GRW

Collapse-based quantum mechanics

There are two kinds of dynamics in collapse-based quantum mechanics *à la* Dirac/von Neumann:

Collapse-based quantum mechanics

There are two kinds of dynamics in collapse-based quantum mechanics *à la* Dirac/von Neumann:

1. Unitary, deterministic, linear Schrödinger dynamics.

Collapse-based quantum mechanics

There are two kinds of dynamics in collapse-based quantum mechanics *à la* Dirac/von Neumann:

1. Unitary, deterministic, linear Schrödinger dynamics.
2. Non-unitary, non-deterministic, non-linear collapse dynamics.

Collapse-based quantum mechanics

There are two kinds of dynamics in collapse-based quantum mechanics *à la* Dirac/von Neumann:

1. Unitary, deterministic, linear Schrödinger dynamics.
2. Non-unitary, non-deterministic, non-linear collapse dynamics.

Textbooks would have us apply collapse dynamics when we perform measurements, but nothing in our physics tells us *what* counts as a measurement, or *why*.

Collapse-based quantum mechanics

There are two kinds of dynamics in collapse-based quantum mechanics *à la* Dirac/von Neumann:

1. Unitary, deterministic, linear Schrödinger dynamics.
2. Non-unitary, non-deterministic, non-linear collapse dynamics.

Textbooks would have us apply collapse dynamics when we perform measurements, but nothing in our physics tells us *what* counts as a measurement, or *why*.

Moreover, we don't have any idea how collapse is supposed to work! (See Bell's 'Against 'Measurement''.)

Dynamical collapse theories introduced

We might seek to tackle these issues directly by modifying the Schrödinger dynamics to include collapse. The resulting theories are (unsurprisingly!) known as *dynamical collapse theories*. Any such theory must respect the following three constraints:

Dynamical collapse theories introduced

We might seek to tackle these issues directly by modifying the Schrödinger dynamics to include collapse. The resulting theories are (unsurprisingly!) known as *dynamical collapse theories*. Any such theory must respect the following three constraints:

1. Modification must have no noticeable effects on microscopic systems.

Dynamical collapse theories introduced

We might seek to tackle these issues directly by modifying the Schrödinger dynamics to include collapse. The resulting theories are (unsurprisingly!) known as *dynamical collapse theories*. Any such theory must respect the following three constraints:

1. Modification must have no noticeable effects on microscopic systems.
2. Modification must prevent macroscopic superpositions.

Dynamical collapse theories introduced

We might seek to tackle these issues directly by modifying the Schrödinger dynamics to include collapse. The resulting theories are (unsurprisingly!) known as *dynamical collapse theories*. Any such theory must respect the following three constraints:

1. Modification must have no noticeable effects on microscopic systems.
2. Modification must prevent macroscopic superpositions.
3. Collapse must occur in accordance with the Born rule. (I.e., $\alpha |\uparrow\rangle + \beta |\downarrow\rangle$ must collapse to either $|\uparrow\rangle$ with probability $|\alpha|^2$, or $|\downarrow\rangle$ with probability $|\beta|^2$.)

Toy dynamical collapse theory

Here's a toy dynamical collapse theory:

Toy dynamical collapse theory

Here's a toy dynamical collapse theory:

- ▶ Particles have probability τ per second of collapsing to some position eigenstate. (For example, $\alpha |x\rangle + \beta |y\rangle$ will collapse to $|x\rangle$ with probability $|\alpha|^2$, or $|y\rangle$ with probability $|\beta|^2$, where $|x\rangle$ and $|y\rangle$ represent different position eigenstates.)

Toy dynamical collapse theory

Here's a toy dynamical collapse theory:

- ▶ Particles have probability τ per second of collapsing to some position eigenstate. (For example, $\alpha |x\rangle + \beta |y\rangle$ will collapse to $|x\rangle$ with probability $|\alpha|^2$, or $|y\rangle$ with probability $|\beta|^2$, where $|x\rangle$ and $|y\rangle$ represent different position eigenstates.)
- ▶ If particles are entangled, then if any one particle collapses, the quantum state for the whole system collapses.

Toy dynamical collapse theory

Here's a toy dynamical collapse theory:

- ▶ Particles have probability τ per second of collapsing to some position eigenstate. (For example, $\alpha|x\rangle + \beta|y\rangle$ will collapse to $|x\rangle$ with probability $|\alpha|^2$, or $|y\rangle$ with probability $|\beta|^2$, where $|x\rangle$ and $|y\rangle$ represent different position eigenstates.)
- ▶ If particles are entangled, then if any one particle collapses, the quantum state for the whole system collapses.
- ▶ If τ is set correctly (e.g. $\tau = 10^{-16}\text{s}^{-1}$), this will ensure that isolated particles collapse very slowly (roughly, after 100 million years), but large, *entangled* systems will collapse much faster.

Technical problem with the toy model

The collapses described above leave the particles which undergo them in perfect eigenstates of the position operator, and of course that entails that the momenta and the energies of those particles (whatever their values may have been just prior to those collapses) will be completely uncertain just following those collapses, and that will give rise to a host of problems: The momenta which electrons in atoms might sometimes acquire in the course of such collapses, for example, would be enough to knock them right out of their orbits; and the energies which certain of the molecules of a gas might sometimes acquire in the course of such collapses would be enough to spontaneously heat those gases up, and those sorts of things are experimentally known not to occur. (Albert 1992, p. 97)

A fix

The fix is to say that we don't collapse onto an *exact position*, but rather onto some *Gaussian function of position*:

A fix

The fix is to say that we don't collapse onto an *exact position*, but rather onto some *Gaussian function of position*:

And it turns out (and this is the punch line) that these curves can nonetheless be made wide enough (at the same time) so that the violations of the conservation of energy and of momentum which the multiplications by these curves will produce will be too small to be observed. (Albert 1992, p. 98)

A fix

The fix is to say that we don't collapse onto an *exact position*, but rather onto some *Gaussian function of position*:

And it turns out (and this is the punch line) that these curves can nonetheless be made wide enough (at the same time) so that the violations of the conservation of energy and of momentum which the multiplications by these curves will produce will be too small to be observed. (Albert 1992, p. 98)

This is the essence of what the GRW dynamical collapse theory seeks to do.

Today

Dynamical collapse theories

The GRW theory

Objections to GRW

Modifications/alternatives to GRW

The GRW theory

- ▶ Collapse: $\psi(x) \rightarrow \mathcal{N}\psi(x) f(x - a)$.

The GRW theory

- ▶ Collapse: $\psi(x) \rightarrow \mathcal{N}\psi(x) f(x - a)$.
- ▶ $f(x - a) = \exp\left(\frac{-(x-a)^2}{2L^2}\right)$ is a Gaussian function centered at $x = a$.

The GRW theory

- ▶ Collapse: $\psi(x) \rightarrow \mathcal{N}\psi(x) f(x - a)$.
- ▶ $f(x - a) = \exp\left(\frac{-(x-a)^2}{2L^2}\right)$ is a Gaussian function centered at $x = a$.
- ▶ a is stochastically selected in line with the Born rule: $\text{Prob}(x_0 < x < x_0 + \delta x) = |\psi(x_0)|^2 \delta x$.

The GRW theory

- ▶ Collapse: $\psi(x) \rightarrow \mathcal{N}\psi(x) f(x - a)$.
- ▶ $f(x - a) = \exp\left(\frac{-(x-a)^2}{2L^2}\right)$ is a Gaussian function centered at $x = a$.
- ▶ a is stochastically selected in line with the Born rule: $\text{Prob}(x_0 < x < x_0 + \delta x) = |\psi(x_0)|^2 \delta x$.
- ▶ \mathcal{N} is a normalisation factor.

The GRW theory

- ▶ Collapse: $\psi(x) \rightarrow \mathcal{N} \psi(x) f(x - a)$.
- ▶ $f(x - a) = \exp\left(\frac{-(x-a)^2}{2L^2}\right)$ is a Gaussian function centered at $x = a$.
- ▶ a is stochastically selected in line with the Born rule: $\text{Prob}(x_0 < x < x_0 + \delta x) = |\psi(x_0)|^2 \delta x$.
- ▶ \mathcal{N} is a normalisation factor.
- ▶ τ and L are new constants of nature.

Dynamical collapse timescales

If we choose $\tau = 10^{-16}\text{s}^{-1}$ and $L = 10^{-5}\text{cm}^{-1}$, then:

Dynamical collapse timescales

If we choose $\tau = 10^{-16}\text{s}^{-1}$ and $L = 10^{-5}\text{cm}^{-1}$, then:

- ▶ Isolated particles will collapse after approximately 100 million years.

Dynamical collapse timescales

If we choose $\tau = 10^{-16}\text{s}^{-1}$ and $L = 10^{-5}\text{cm}^{-1}$, then:

- ▶ Isolated particles will collapse after approximately 100 million years.
- ▶ A dust mote of 10^{-5}cm^{-1} in diameter will collapse after about 1 second.

Dynamical collapse timescales

If we choose $\tau = 10^{-16}\text{s}^{-1}$ and $L = 10^{-5}\text{cm}^{-1}$, then:

- ▶ Isolated particles will collapse after approximately 100 million years.
- ▶ A dust mote of 10^{-5}cm^{-1} in diameter will collapse after about 1 second.
- ▶ A macroscopic object (e.g. a grain of sand less than 1mm across) will collapse extremely quickly (less than 10^{-5}s for the grain of sand).

Dynamical collapse timescales

If we choose $\tau = 10^{-16}\text{s}^{-1}$ and $L = 10^{-5}\text{cm}^{-1}$, then:

- ▶ Isolated particles will collapse after approximately 100 million years.
- ▶ A dust mote of 10^{-5}cm^{-1} in diameter will collapse after about 1 second.
- ▶ A macroscopic object (e.g. a grain of sand less than 1mm across) will collapse extremely quickly (less than 10^{-5}s for the grain of sand).

Exercise: Think about how the GRW dynamical collapse timescale interacts with the decoherence timescales discussed in Lecture 3.

Today

Dynamical collapse theories

The GRW theory

Objections to GRW

Modifications/alternatives to GRW

Objection 1: The problem of tails

GRW dynamical collapse does not get rid of superpositions completely. It takes states like

$$\alpha |\text{here}\rangle + \beta |\text{there}\rangle$$

Objection 1: The problem of tails

GRW dynamical collapse does not get rid of superpositions completely. It takes states like

$$\alpha |\text{here}\rangle + \beta |\text{there}\rangle$$

to states like

$$\sqrt{1 - \epsilon^2} |\text{here}\rangle + \epsilon |\text{there}\rangle ,$$

where ϵ is small.

Objection 1: The problem of tails

GRW dynamical collapse does not get rid of superpositions completely. It takes states like

$$\alpha |\text{here}\rangle + \beta |\text{there}\rangle$$

to states like

$$\sqrt{1 - \epsilon^2} |\text{here}\rangle + \epsilon |\text{there}\rangle ,$$

where ϵ is small.

But there are obvious problems with this....

The problem of structured tails

- ▶ Suppose the system is initially in a state such as $\alpha |\text{here}\rangle + \beta |\text{there}\rangle$, but then collapses to a state such as $\sqrt{1 - \epsilon^2} |\text{here}\rangle + \epsilon |\text{there}\rangle$.

The problem of structured tails

- ▶ Suppose the system is initially in a state such as $\alpha |\text{here}\rangle + \beta |\text{there}\rangle$, but then collapses to a state such as $\sqrt{1 - \epsilon^2} |\text{here}\rangle + \epsilon |\text{there}\rangle$.
- ▶ Post-collapse, the ‘collapsed’ part of the superposition (above, $|\text{here}\rangle$) is ‘damped down’, but does not disappear completely.

The problem of structured tails

- ▶ Suppose the system is initially in a state such as $\alpha | \text{here} \rangle + \beta | \text{there} \rangle$, but then collapses to a state such as $\sqrt{1 - \epsilon^2} | \text{here} \rangle + \epsilon | \text{there} \rangle$.
- ▶ Post-collapse, the ‘collapsed’ part of the superposition (above, $| \text{here} \rangle$) is ‘damped down’, but does not disappear completely.

As Wallace writes:

Why should the continued presence of the ‘there’ term in the superposition—the continued indefiniteness of the system between ‘here’ and ‘there’—be ameliorated in any way at all just because the ‘there’ term has low amplitude? (Wallace 2008, p. 43)

The problem of structured tails

- ▶ Suppose the system is initially in a state such as $\alpha |\text{here}\rangle + \beta |\text{there}\rangle$, but then collapses to a state such as $\sqrt{1 - \epsilon^2} |\text{here}\rangle + \epsilon |\text{there}\rangle$.
- ▶ Post-collapse, the ‘collapsed’ part of the superposition (above, $|\text{here}\rangle$) is ‘damped down’, but does not disappear completely.

As Wallace writes:

Why should the continued presence of the ‘there’ term in the superposition—the continued indefiniteness of the system between ‘here’ and ‘there’—be ameliorated in any way at all just because the ‘there’ term has low amplitude? (Wallace 2008, p. 43)

This is the ‘problem of structured tails’, which is an issue specific to dynamical collapse theories such as GRW.

The problem of bare tails

- ▶ Even if we ignore the 'there' state, the wavefunction of $|\text{here}\rangle$ is itself spatially highly delocalised.

The problem of bare tails

- ▶ Even if we ignore the 'there' state, the wavefunction of $|\text{here}\rangle$ is itself spatially highly delocalised.
- ▶ Its centre-of-mass wavefunction is no doubt a Gaussian, and Gaussians are completely delocalised in space, for all that they may be concentrated in one region or another.

The problem of bare tails

- ▶ Even if we ignore the 'there' state, the wavefunction of $|\text{here}\rangle$ is itself spatially highly delocalised.
- ▶ Its centre-of-mass wavefunction is no doubt a Gaussian, and Gaussians are completely delocalised in space, for all that they may be concentrated in one region or another.
- ▶ So how can a delocalised wave packet possibly count as a localised particle?

The problem of bare tails

- ▶ Even if we ignore the ‘there’ state, the wavefunction of $|\text{here}\rangle$ is itself spatially highly delocalised.
- ▶ Its centre-of-mass wavefunction is no doubt a Gaussian, and Gaussians are completely delocalised in space, for all that they may be concentrated in one region or another.
- ▶ So how can a delocalised wave packet possibly count as a localised particle?

But, as Wallace writes on this problem:

This problem has little or nothing to do with the GRW theory. Rather, it is an unavoidable consequence of using wave packets to stand in for localised particles. For no wave packet evolving unitarily will remain in any finite spatial region for more than an instant. (Wallace 2008, p. 43)

The problem of bare tails

- ▶ Even if we ignore the ‘there’ state, the wavefunction of $|here\rangle$ is itself spatially highly delocalised.
- ▶ Its centre-of-mass wavefunction is no doubt a Gaussian, and Gaussians are completely delocalised in space, for all that they may be concentrated in one region or another.
- ▶ So how can a delocalised wave packet possibly count as a localised particle?

But, as Wallace writes on this problem:

This problem has little or nothing to do with the GRW theory. Rather, it is an unavoidable consequence of using wave packets to stand in for localised particles. For no wave packet evolving unitarily will remain in any finite spatial region for more than an instant. (Wallace 2008, p. 43)

This is the ‘problem of bare tails’, which is *not* an issue specific to dynamical collapse theories such as GRW.

The two problems

Strangely, most of the literature on the problem of tails addresses the problem of bare tails (which is not specific to dynamical collapse theories), rather than the problem of structured tails (which *is* specific to dynamical collapse theories).

The two problems

Strangely, most of the literature on the problem of tails addresses the problem of bare tails (which is not specific to dynamical collapse theories), rather than the problem of structured tails (which *is* specific to dynamical collapse theories).

Still, let's look at how people have tried to tackle the problem of bare tails.

Addressing bare tails

We want to say that, after collapse, systems and particles have definite positions, even though the wavefunction is non-zero everywhere in space.

Addressing bare tails

We want to say that, after collapse, systems and particles have definite positions, even though the wavefunction is non-zero everywhere in space.

Albert and Lower (1996) suggest we get achieve this by dropping the *eigenvector-eigenvalue link*, and replacing it with a so-called *fuzzy link*.

The fuzzy link

Eigenvector-eigenvalue link: A system in a state $|\psi\rangle$ has a definite value of the physical quantity associated to some observable \hat{X} , iff

$$\hat{X}|\psi\rangle = x|\psi\rangle.$$

The fuzzy link

Eigenvector-eigenvalue link: A system in a state $|\psi\rangle$ has a definite value of the physical quantity associated to some observable \hat{X} , iff

$$\hat{X}|\psi\rangle = x|\psi\rangle.$$

Fuzzy link: A system in a state $|\psi\rangle$ has a definite value of the physical quantity associated to some observable \hat{X} , iff

$$\left| \hat{X}|\psi\rangle - x|\psi\rangle \right| \leq \lambda,$$

for some small λ .

The counting anomaly

The fuzzy link solves the problem of bare tails, but it leads to the so-called *counting anomaly*:

The counting anomaly

The fuzzy link solves the problem of bare tails, but it leads to the so-called *counting anomaly*:

- ▶ Particle 1 is in some box (according to the fuzzy link).
- ▶ Particle 2 is in that box (according to the fuzzy link) ...
- ▶ ... Particle n is in that box (according to the fuzzy link).
- ▶ On the fuzzy link, this *does not imply*: N particles are in the box.

The counting anomaly in more detail

- ▶ Suppose that the wavefunction of each particle is very strongly peaked inside the box, so that if \hat{X}_i is the 'particle i is in the box' operator, then $|\hat{X}_i |\psi\rangle - x |\psi\rangle| \sim \epsilon$, for $\epsilon \ll \lambda$, so that each particle should be counted as being inside the box.

The counting anomaly in more detail

- ▶ Suppose that the wavefunction of each particle is very strongly peaked inside the box, so that if \hat{X}_i is the 'particle i is in the box' operator, then $\left| \hat{X}_i |\psi\rangle - x |\psi\rangle \right| \sim \epsilon$, for $\epsilon \ll \lambda$, so that each particle should be counted as being inside the box.
- ▶ Heuristically, $\left| \hat{X}_i |\psi\rangle \right| \sim 1 - \epsilon$.

The counting anomaly in more detail

- ▶ Suppose that the wavefunction of each particle is very strongly peaked inside the box, so that if \hat{X}_i is the 'particle i is in the box' operator, then $|\hat{X}_i |\psi\rangle - x |\psi\rangle| \sim \epsilon$, for $\epsilon \ll \lambda$, so that each particle should be counted as being inside the box.
- ▶ Heuristically, $|\hat{X}_i |\psi\rangle| \sim 1 - \epsilon$.
- ▶ But now consider the proposition 'all N particles are in the box'. By definition, this is represented by the operator $\hat{X} = \prod_{i=1}^N \hat{X}_i$.

The counting anomaly in more detail

- ▶ Suppose that the wavefunction of each particle is very strongly peaked inside the box, so that if \hat{X}_i is the 'particle i is in the box' operator, then $|\hat{X}_i |\psi\rangle - x |\psi\rangle| \sim \epsilon$, for $\epsilon \ll \lambda$, so that each particle should be counted as being inside the box.
- ▶ Heuristically, $|\hat{X}_i |\psi\rangle| \sim 1 - \epsilon$.
- ▶ But now consider the proposition 'all N particles are in the box'. By definition, this is represented by the operator $\hat{X} = \prod_{i=1}^N \hat{X}_i$.
- ▶ Suppose each particle has identical state $|\psi\rangle$; i.e. suppose that each $|\psi\rangle$ is highly localised in the box, as above. Then the overall state of the N particles is $|\Psi\rangle = \bigotimes_{i=1}^N |\psi\rangle$.

The counting anomaly in more detail

- ▶ Suppose that the wavefunction of each particle is very strongly peaked inside the box, so that if \hat{X}_i is the ‘particle i is in the box’ operator, then $|\hat{X}_i |\psi\rangle - x |\psi\rangle| \sim \epsilon$, for $\epsilon \ll \lambda$, so that each particle should be counted as being inside the box.
- ▶ Heuristically, $|\hat{X}_i |\psi\rangle| \sim 1 - \epsilon$.
- ▶ But now consider the proposition ‘all N particles are in the box’. By definition, this is represented by the operator $\hat{X} = \prod_{i=1}^N \hat{X}_i$.
- ▶ Suppose each particle has identical state $|\psi\rangle$; i.e. suppose that each $|\psi\rangle$ is highly localised in the box, as above. Then the overall state of the N particles is $|\Psi\rangle = \bigotimes_{i=1}^N |\psi\rangle$.
- ▶ Then $|\hat{X} |\Psi\rangle| = \prod_{i=1}^N |\hat{X}_i |\psi\rangle| = (1 - \epsilon)^N$.

Wallace on the counting anomaly

[T]his is unfortunate for the Fuzzy Link. For no matter how small ϵ may be, there will be some value of N for which $(1 - \epsilon)^N < \lambda$. And for that value of N , the Fuzzy Link tells us that it is false that all N particles are in the box, even as it tells us that, for each of the N particles, it is true that that particle is in the box. (Wallace 2008, p. 44)

The mass density link

An alternative to the fuzzy link is the 'mass density link'.

The mass density link

An alternative to the fuzzy link is the 'mass density link'.

According to the mass density link, a particle is in a box iff some sufficiently high fraction $1 - \epsilon$ of its mass is in the box.

The meaning of 'all N particles are in the box' is 'particle 1 is in the box and particle 2 is in the box and ...', and this is true iff all the constituent propositions are true.

The mass density link

An alternative to the fuzzy link is the ‘mass density link’.

According to the mass density link, a particle is in a box iff some sufficiently high fraction $1 - \epsilon$ of its mass is in the box.

The meaning of ‘all N particles are in the box’ is ‘particle 1 is in the box and particle 2 is in the box and ...’, and this is true iff all the constituent propositions are true.

The mass density link doesn’t face the counting anomaly, but some argue that it has problems of its own...

The location anomaly

It's sometimes claimed that the mass density link evades the counting anomaly only at the cost of a so-called 'location anomaly'. Here's how Wallace summarises the situation:

The location anomaly

It's sometimes claimed that the mass density link evades the counting anomaly only at the cost of a so-called 'location anomaly'. Here's how Wallace summarises the situation:

This anomaly arises when we consider the process of looking at the box and physically counting the number of particles in it. The ordinary quantum theory—which the GRW theory is supposed to reproduce—then predicts that the expected number of particles found in the box will be somewhat less than N . Lewis claims that this clash between the predictions of how many particles are found in the box and how many are actually in the box “violates the entailments of ordinary language” (Lewis 2005, p. 174). (Wallace 2008, pp. 46-47)

Responding to the location anomaly

Ghirardi and Bassi (1993, 2003), and separately Wallace (2008), are bemused by this criticism:

Responding to the location anomaly

Ghirardi and Bassi (1993, 2003), and separately Wallace (2008), are bemused by this criticism:

[W]e have a theory which (a) gives a perfectly well-defined description of how many particles are in the box; (b) allows a precise description, in terms acceptable to the realist, of the measurement process by which we determine how many particles are in the box; (c) predicts that if the number of particles is sufficiently (i.e., ridiculously) large there will be tiny deviations between the actual number of particles and the recorded number of particles. They, and I, fail to see what the problem is here; I leave readers to reach their own conclusions. (Wallace 2008, p. 47)

Status of the link principles

- ▶ Let's step back a bit. What's the actual status of the link principles?

Status of the link principles

- ▶ Let's step back a bit. What's the actual status of the link principles?
- ▶ We've been treating them like elements of physical law.

Status of the link principles

- ▶ Let's step back a bit. What's the actual status of the link principles?
- ▶ We've been treating them like elements of physical law.
- ▶ As Wallace (2008, p. 47) points out, this would seem to invite a dualistic ontology, of (a) the wavefunction, and (b) either mass densities or fuzzy-link-defined classical properties.

Status of the link principles

- ▶ Let's step back a bit. What's the actual status of the link principles?
- ▶ We've been treating them like elements of physical law.
- ▶ As Wallace (2008, p. 47) points out, this would seem to invite a dualistic ontology, of (a) the wavefunction, and (b) either mass densities or fuzzy-link-defined classical properties.
- ▶ This is unattractive—better to view the links as “perspicacious ways of picking out certain properties of [the] wavefunction”; that is, as linguistic tools.

Status of the link principles

- ▶ Let's step back a bit. What's the actual status of the link principles?
- ▶ We've been treating them like elements of physical law.
- ▶ As Wallace (2008, p. 47) points out, this would seem to invite a dualistic ontology, of (a) the wavefunction, and (b) either mass densities or fuzzy-link-defined classical properties.
- ▶ This is unattractive—better to view the links as “perspicacious ways if picking out certain properties of [the] wavefunction”; that is, as linguistic tools.
- ▶ But then they are not going to be sufficient to help with tails problems, and one worries that GRW is just ‘Everett in denial’.

More on 'Everett-in-denial'

Consider a post-GRW collapsed state, such as

$$\sqrt{1 - \epsilon^2} |\text{alive}\rangle + \epsilon |\text{dead}\rangle .$$

More on 'Everett-in-denial'

Consider a post-GRW collapsed state, such as

$$\sqrt{1 - \epsilon^2} |\text{alive}\rangle + \epsilon |\text{dead}\rangle .$$

- ▶ On the Everett interpretation, a structural/functional definition of macro-ontology is given (more on this in Lecture 7).

More on 'Everett-in-denial'

Consider a post-GRW collapsed state, such as

$$\sqrt{1 - \epsilon^2} |\text{alive}\rangle + \epsilon |\text{dead}\rangle .$$

- ▶ On the Everett interpretation, a structural/functional definition of macro-ontology is given (more on this in Lecture 7).
- ▶ Suppose we accept this. Then all there is to being a cat is being something structured like a cat.

More on ‘Everett-in-denial’

Consider a post-GRW collapsed state, such as

$$\sqrt{1 - \epsilon^2} |\text{alive}\rangle + \epsilon |\text{dead}\rangle .$$

- ▶ On the Everett interpretation, a structural/functional definition of macro-ontology is given (more on this in Lecture 7).
- ▶ Suppose we accept this. Then all there is to being a cat is being something structured like a cat.
- ▶ There’s still something structured like an (alive!) cat in the above state—it’s just very low amplitude.

More on ‘Everett-in-denial’

Consider a post-GRW collapsed state, such as

$$\sqrt{1 - \epsilon^2} |\text{alive}\rangle + \epsilon |\text{dead}\rangle .$$

- ▶ On the Everett interpretation, a structural/functional definition of macro-ontology is given (more on this in Lecture 7).
- ▶ Suppose we accept this. Then all there is to being a cat is being something structured like a cat.
- ▶ There’s still something structured like an (alive!) cat in the above state—it’s just very low amplitude.
- ▶ On Everettian ontology, a low amplitude cat is still a cat.

More on ‘Everett-in-denial’

Consider a post-GRW collapsed state, such as

$$\sqrt{1 - \epsilon^2} |\text{alive}\rangle + \epsilon |\text{dead}\rangle .$$

- ▶ On the Everett interpretation, a structural/functional definition of macro-ontology is given (more on this in Lecture 7).
- ▶ Suppose we accept this. Then all there is to being a cat is being something structured like a cat.
- ▶ There’s still something structured like an (alive!) cat in the above state—it’s just very low amplitude.
- ▶ On Everettian ontology, a low amplitude cat is still a cat.
- ▶ So, there are still two cats (one alive and one dead!) in the above state.

More on 'Everett-in-denial'

Consider a post-GRW collapsed state, such as

$$\sqrt{1 - \epsilon^2} |\text{alive}\rangle + \epsilon |\text{dead}\rangle .$$

- ▶ On the Everett interpretation, a structural/functional definition of macro-ontology is given (more on this in Lecture 7).
- ▶ Suppose we accept this. Then all there is to being a cat is being something structured like a cat.
- ▶ There's still something structured like an (alive!) cat in the above state—it's just very low amplitude.
- ▶ On Everettian ontology, a low amplitude cat is still a cat.
- ▶ So, there are still two cats (one alive and one dead!) in the above state.
- ▶ So is GRW just (some messy version of) Everett, in disguise?

Objection 2: Probability in GRW

In some ways, GRW collapse provides but the perfect example of objective chance. But what does τ *mean*? Is it only analysable in terms of propensities, or are other options available?

Objection 3: New constants of nature

- ▶ GRW theorists will aver that the constants of nature are not (\hbar, G_N, c) , but (\hbar, G_N, c, L, τ) .

Objection 3: New constants of nature

- ▶ GRW theorists will aver that the constants of nature are not (\hbar, G_N, c) , but (\hbar, G_N, c, L, τ) .
- ▶ Is the introduction of these two further constants of nature *ad hoc*?

Objection 3: New constants of nature

- ▶ GRW theorists will aver that the constants of nature are not (\hbar, G_N, c) , but (\hbar, G_N, c, L, τ) .
- ▶ Is the introduction of these two further constants of nature *ad hoc*?
- ▶ Some, e.g. Smolin (2013), aver that *everything* in a physics needs to have some explanation; the introduction of extra constants of nature would seem to run against this.

Objection 4: Compatibility with relativity

Is GRW compatible with relativity theory?

Objection 4: Compatibility with relativity

Is GRW compatible with relativity theory?

When I saw this theory first, I thought that I could blow it out of the water, by showing that it was grossly in violation of Lorentz invariance. That's connected with the problem of 'quantum entanglement', the EPR paradox. (Bell 1989, p. 1)

Tumulka's development

- ▶ Tumulka (2006) succeeded in proposing a relativistic version of the GRW theory for N non-interacting distinguishable particles.

Tumulka's development

- ▶ Tumulka (2006) succeeded in proposing a relativistic version of the GRW theory for N non-interacting distinguishable particles.
- ▶ This was the first proposal of a relativistic dynamical reduction mechanism, which satisfies all relativistic requirements. In particular it is free from divergences and foliation independent.

Tumulka's development

- ▶ Tumulka (2006) succeeded in proposing a relativistic version of the GRW theory for N non-interacting distinguishable particles.
- ▶ This was the first proposal of a relativistic dynamical reduction mechanism, which satisfies all relativistic requirements. In particular it is free from divergences and foliation independent.
- ▶ However, it is formulated only for systems containing a fixed number of noninteracting fermions.

Tumulka's development

- ▶ Tumulka (2006) succeeded in proposing a relativistic version of the GRW theory for N non-interacting distinguishable particles.
- ▶ This was the first proposal of a relativistic dynamical reduction mechanism, which satisfies all relativistic requirements. In particular it is free from divergences and foliation independent.
- ▶ However, it is formulated only for systems containing a fixed number of noninteracting fermions.
- ▶ Extending to QFT is another matter: Wallace (2022) argues that there is (as-yet) no viable dynamical collapse theory for QFT, *so there is not even a genuine case of underdetermination between dynamical collapse theories (or hidden variable theories, for that matter) and Everett!*

Today

Dynamical collapse theories

The GRW theory

Objections to GRW

Modifications/alternatives to GRW

Modifications/alternatives to GRW

Let's close by thinking about some dynamical collapse theories which are modifications/alternatives to GRW.

GRW flash ontology

The GRW 'flash ontology' approach works as follows:

GRW flash ontology

The GRW 'flash ontology' approach works as follows:

- ▶ Treat the GRW collapses ('flashes') as the basic elements of one's ontology.

GRW flash ontology

The GRW 'flash ontology' approach works as follows:

- ▶ Treat the GRW collapses ('flashes') as the basic elements of one's ontology.
- ▶ I.e., treat the collapses as a localisation 'event' at (x^i, t) .

GRW flash ontology

The GRW 'flash ontology' approach works as follows:

- ▶ Treat the GRW collapses ('flashes') as the basic elements of one's ontology.
- ▶ I.e., treat the collapses as a localisation 'event' at (x^i, t) .
- ▶ Flashes are located in spacetime.

GRW flash ontology

The GRW 'flash ontology' approach works as follows:

- ▶ Treat the GRW collapses ('flashes') as the basic elements of one's ontology.
- ▶ I.e., treat the collapses as a localisation 'event' at (x^i, t) .
- ▶ Flashes are located in spacetime.
- ▶ Given a sufficiently long sequence of flashes, the final state is not sensitive to the original state.

GRW flash ontology

The GRW 'flash ontology' approach works as follows:

- ▶ Treat the GRW collapses ('flashes') as the basic elements of one's ontology.
- ▶ I.e., treat the collapses as a localisation 'event' at (x^i, t) .
- ▶ Flashes are located in spacetime.
- ▶ Given a sufficiently long sequence of flashes, the final state is not sensitive to the original state.
- ▶ Forget the wavefunction altogether: future flashes depend only on past history of flashes.

GRW flash ontology

The GRW 'flash ontology' approach works as follows:

- ▶ Treat the GRW collapses ('flashes') as the basic elements of one's ontology.
- ▶ I.e., treat the collapses as a localisation 'event' at (x^i, t) .
- ▶ Flashes are located in spacetime.
- ▶ Given a sufficiently long sequence of flashes, the final state is not sensitive to the original state.
- ▶ Forget the wavefunction altogether: future flashes depend only on past history of flashes.

Arguably, this solves the problem of tails (but does it really?), although (a) it's a revisionary ontology, and (b) the other problems persist.

GRW flash ontology

The GRW 'flash ontology' approach works as follows:

- ▶ Treat the GRW collapses ('flashes') as the basic elements of one's ontology.
- ▶ I.e., treat the collapses as a localisation 'event' at (x^i, t) .
- ▶ Flashes are located in spacetime.
- ▶ Given a sufficiently long sequence of flashes, the final state is not sensitive to the original state.
- ▶ Forget the wavefunction altogether: future flashes depend only on past history of flashes.

Arguably, this solves the problem of tails (but does it really?), although (a) it's a revisionary ontology, and (b) the other problems persist.

(See (Esfeld & Gisin 2013) for more.)

Continuous spontaneous localisation models

- ▶ In 'continuous spontaneous localisation' (CSL) models (Pearle 1989), the discontinuous jumps which characterize dynamical collapse theories such as GRW are replaced by a *continuous* stochastic evolution in the Hilbert space.

Continuous spontaneous localisation models

- ▶ In ‘continuous spontaneous localisation’ (CSL) models (Pearle 1989), the discontinuous jumps which characterize dynamical collapse theories such as GRW are replaced by a *continuous* stochastic evolution in the Hilbert space.
- ▶ For any CSL dynamics there is a hitting dynamics which, from a physical point of view, is ‘as close to it as one wants’.

Continuous spontaneous localisation models

- ▶ In ‘continuous spontaneous localisation’ (CSL) models (Pearle 1989), the discontinuous jumps which characterize dynamical collapse theories such as GRW are replaced by a *continuous* stochastic evolution in the Hilbert space.
- ▶ For any CSL dynamics there is a hitting dynamics which, from a physical point of view, is ‘as close to it as one wants’.
- ▶ CSL offers various empirically detectable differences from standard QM—see (Ghirardi 2011, §7).

Continuous spontaneous localisation models

- ▶ In ‘continuous spontaneous localisation’ (CSL) models (Pearle 1989), the discontinuous jumps which characterize dynamical collapse theories such as GRW are replaced by a *continuous* stochastic evolution in the Hilbert space.
- ▶ For any CSL dynamics there is a hitting dynamics which, from a physical point of view, is ‘as close to it as one wants’.
- ▶ CSL offers various empirically detectable differences from standard QM—see (Ghirari 2011, §7).
- ▶ The hope is that continuous localisation will be more easily reconcilable with relativity theory and QFT than the GRW-style ‘hits’, but there remains much to be done on this front.

Continuous spontaneous localisation models

- ▶ In ‘continuous spontaneous localisation’ (CSL) models (Pearle 1989), the discontinuous jumps which characterize dynamical collapse theories such as GRW are replaced by a *continuous* stochastic evolution in the Hilbert space.
- ▶ For any CSL dynamics there is a hitting dynamics which, from a physical point of view, is ‘as close to it as one wants’.
- ▶ CSL offers various empirically detectable differences from standard QM—see (Ghirardi 2011, §7).
- ▶ The hope is that continuous localisation will be more easily reconcilable with relativity theory and QFT than the GRW-style ‘hits’, but there remains much to be done on this front.
- ▶ In any case, it’s not completely obvious that CSL will offer a satisfactory resolution to the problem of tails.

Summary

Today, I've:

Summary

Today, I've:

- ▶ Introduced dynamical collapse theories, which take one horn of Bell's dilemma and *modify* the formalism of quantum mechanics.

Summary

Today, I've:

- ▶ Introduced dynamical collapse theories, which take one horn of Bell's dilemma and *modify* the formalism of quantum mechanics.
- ▶ Presented how dynamical collapse theories work in practice.

Summary

Today, I've:

- ▶ Introduced dynamical collapse theories, which take one horn of Bell's dilemma and *modify* the formalism of quantum mechanics.
- ▶ Presented how dynamical collapse theories work in practice.
- ▶ Presented the problem of tails in its various guises, and how dynamical collapse theories have tried to respond.

Summary

Today, I've:

- ▶ Introduced dynamical collapse theories, which take one horn of Bell's dilemma and *modify* the formalism of quantum mechanics.
- ▶ Presented how dynamical collapse theories work in practice.
- ▶ Presented the problem of tails in its various guises, and how dynamical collapse theories have tried to respond.
- ▶ Presented other problems for dynamical collapse theories, namely those to do with probabilities, constants of nature, and compatibility with relativity.

Summary

Today, I've:

- ▶ Introduced dynamical collapse theories, which take one horn of Bell's dilemma and *modify* the formalism of quantum mechanics.
- ▶ Presented how dynamical collapse theories work in practice.
- ▶ Presented the problem of tails in its various guises, and how dynamical collapse theories have tried to respond.
- ▶ Presented other problems for dynamical collapse theories, namely those to do with probabilities, constants of nature, and compatibility with relativity.
- ▶ Presented some alternative dynamical collapse models (flash and CSL).

Summary

Today, I've:

- ▶ Introduced dynamical collapse theories, which take one horn of Bell's dilemma and *modify* the formalism of quantum mechanics.
- ▶ Presented how dynamical collapse theories work in practice.
- ▶ Presented the problem of tails in its various guises, and how dynamical collapse theories have tried to respond.
- ▶ Presented other problems for dynamical collapse theories, namely those to do with probabilities, constants of nature, and compatibility with relativity.
- ▶ Presented some alternative dynamical collapse models (flash and CSL).

Next time: *hidden variable theories*.

References I

-  D. Z. Albert, *Quantum Mechanics and Experience*, Cambridge, MA: Harvard University Press, 1992.
-  D. Z. Albert and B. Loewer, *Tails of Schrödinger's Cat*, in R. Clifton (ed.), *Perspectives on Quantum Reality*, Dordrecht: Kluwer Academic Publishers, 1996.
-  John S. Bell, “Towards an Exact Quantum mechanics”, in S. Deser and R. J. Finkelstein (eds.), *Themes in Contemporary Physics II*, Singapore: World Scientific, 1989.
-  Michael Esfeld and Nicolas Gisin, “The GRW Flash Theory: A Relativistic Quantum Ontology of Matter in Space-Time?”, *Philosophy of Science* 81(2), pp. 248–64, 2014.
-  Giancarlo Ghirardi, “Collapse Theories”, in E. N. Zalta (ed.), *The Stanford Encyclopedia of Philosophy*, 2011.

References II

-  P. Pearle, “Combining Stochastic Dynamical State-Vector Reduction with Spontaneous Localization”, *Physical Review A* 39, 2277, 1989.
-  Lee Smolin, “A Perspective on the Landscape Problem”, *Foundations of Physics* 43(1), pp. 21–45, 2013.
-  R. Tumulka, “A Relativistic Version of the Ghirardi-Rimini-Weber Model”, *Journal of Statistical Physics* 125, 821, 2006.
-  David Wallace, “Philosophy of Quantum Mechanics”, in D. Rickles (ed.), *The Ashgate Companion to Contemporary Philosophy of Physics*, London: Routledge, 2008.
-  David Wallace, “The Sky is Blue, and Other Reasons Quantum Mechanics is Not Underdetermined by Evidence”, *European Journal for Philosophy of Science* 13(54), 2023.