

# IPP-QM-9: EPR and Bell's theorem

James Read<sup>1</sup>

<sup>1</sup>Faculty of Philosophy, University of Oxford, UK, OX2 6GG

MT24

# The course

1. Basic quantum formalism
2. Density operators and entanglement
3. Decoherence
4. The measurement problem
5. Dynamical collapse theories
6. Bohmian mechanics
7. Everettian structure
8. Everettian probability
9. EPR and Bell's theorem
10. The Bell-CHSH inequalities and possible responses
11. Contextuality
12. The PBR theorem
13. Quantum logic
14. Pragmatism and QBism
15. Relational quantum mechanics
16. Wavefunction realism

# Today

Locality and separability

The Einstein-Podolsky-Rosen argument

Bell's 1964 theorem

# Today

Locality and separability

The Einstein-Podolsky-Rosen argument

Bell's 1964 theorem

# Locality and separability

The first thing to get clear on today are two key terms which are ubiquitous in the foundations of quantum mechanics: *locality* and *separability*.

# Locality

What does it mean for a theory to be *local*?

# Locality

What does it mean for a theory to be *local*?

- ▶ The objects of the world can only be affected by objects in the neighbourhood of their location?

# Locality

What does it mean for a theory to be *local*?

- ▶ The objects of the world can only be affected by objects in the neighbourhood of their location?
- ▶ The objects and properties of the world have well defined spatial locations?



# Locality

What does it mean for a theory to be *local*?

- ▶ The objects of the world can only be affected by objects in the neighbourhood of their location?
- ▶ The objects and properties of the world have well defined spatial locations?
- ▶ We cannot produce observable effects on objects not in our neighbourhood?

# Locality

What does it mean for a theory to be *local*?

- ▶ The objects of the world can only be affected by objects in the neighbourhood of their location?
- ▶ The objects and properties of the world have well defined spatial locations?
- ▶ We cannot produce observable effects on objects not in our neighbourhood?

Is locality the conjunction of these? Or some subset or other?

# Action-at-a-distance

- ▶ One thing which locality could mean is this: *no action at a distance*.

# Action-at-a-distance

- ▶ One thing which locality could mean is this: *no action at a distance*.
- ▶ If a physical theory allows one system to directly change, influence, alter, or otherwise interact with another system at a remote location, unmediated by some material connection, this is action-at-a-distance.

# Action-at-a-distance

- ▶ One thing which locality could mean is this: *no action at a distance*.
- ▶ If a physical theory allows one system to directly change, influence, alter, or otherwise interact with another system at a remote location, unmediated by some material connection, this is action-at-a-distance.
  - ▶ Newtonian mechanics allows action-at-a-distance.
  - ▶ Maxwell's equations and Einstein's relativity: mediated by fields and potentials with finite propagation speeds, so no action-at-a-distance.

# Action-at-a-distance

- ▶ One thing which locality could mean is this: *no action at a distance*.
- ▶ If a physical theory allows one system to directly change, influence, alter, or otherwise interact with another system at a remote location, unmediated by some material connection, this is action-at-a-distance.
  - ▶ Newtonian mechanics allows action-at-a-distance.
  - ▶ Maxwell's equations and Einstein's relativity: mediated by fields and potentials with finite propagation speeds, so no action-at-a-distance.
- ▶ What about quantum mechanics?

# Action-at-a-distance

- ▶ One thing which locality could mean is this: *no action at a distance*.
- ▶ If a physical theory allows one system to directly change, influence, alter, or otherwise interact with another system at a remote location, unmediated by some material connection, this is action-at-a-distance.
  - ▶ Newtonian mechanics allows action-at-a-distance.
  - ▶ Maxwell's equations and Einstein's relativity: mediated by fields and potentials with finite propagation speeds, so no action-at-a-distance.
- ▶ What about quantum mechanics?
  - ▶ Dirac/von Neumann collapse dynamics is non-local in this sense.
  - ▶ Bohmian dynamics involves action-at-a-distance, as we saw in Lecture 6.

# Signal locality

- ▶ If there is action-at-a-distance why can't we observe it?



# Signal locality

- ▶ If there is action-at-a-distance why can't we observe it?
- ▶ Why can't we use it to send superluminal signals?

# Signal locality

- ▶ If there is action-at-a-distance why can't we observe it?
- ▶ Why can't we use it to send superluminal signals?
- ▶ In particular contexts (of e.g. particular collapse mechanisms), people show that this is not possible by proving 'no signalling theorems'.

# Signal locality

- ▶ If there is action-at-a-distance why can't we observe it?
- ▶ Why can't we use it to send superluminal signals?
- ▶ In particular contexts (of e.g. particular collapse mechanisms), people show that this is not possible by proving 'no signalling theorems'.
- ▶ So you might have action-at-a-distance at the fundamental ontological level, but still signal locality at the empirical/operational level.

# Three senses of 'at-a-distance'

Einstein locality: 'At-a-distance' means at a spacelike separation between spacetime events. This assumes at minimum special relativity theory.

# Three senses of 'at-a-distance'

Einstein locality: 'At-a-distance' means at a spacelike separation between spacetime events. This assumes at minimum special relativity theory.

Bell locality: 'At-a-distance' means beyond causal influences recognised by current physical theories.

# Three senses of 'at-a-distance'

- Einstein locality: 'At-a-distance' means at a spacelike separation between spacetime events. This assumes at minimum special relativity theory.
- Bell locality: 'At-a-distance' means beyond causal influences recognised by current physical theories.
- EPR locality: "A physical theory is EPR-local iff according to the theory procedures carried out in one region do not immediately disturb the physical state of systems in sufficiently distant regions in any significant way."

# Three senses of 'at-a-distance'

Einstein locality: 'At-a-distance' means at a spacelike separation between spacetime events. This assumes at minimum special relativity theory.

Bell locality: 'At-a-distance' means beyond causal influences recognised by current physical theories.

EPR locality: "A physical theory is EPR-local iff according to the theory procedures carried out in one region do not immediately disturb the physical state of systems in sufficiently distant regions in any significant way."

(The first two of these are from Redhead (1987), while the third is from Maudlin (2014).)

# Comparing the three senses of action-at-a-distance

- ▶ Einstein locality imposes the strongest restrictions on locality; EPR locality is the weakest due e.g. to the vagueness of 'do not immediately disturb' and 'sufficiently distant'.



# Comparing the three senses of action-at-a-distance

- ▶ Einstein locality imposes the strongest restrictions on locality; EPR locality is the weakest due e.g. to the vagueness of 'do not immediately disturb' and 'sufficiently distant'.
- ▶ A failure of EPR locality implies a failure of Bell locality and Einstein locality.

# Comparing the three senses of action-at-a-distance

- ▶ Einstein locality imposes the strongest restrictions on locality; EPR locality is the weakest due e.g. to the vagueness of 'do not immediately disturb' and 'sufficiently distant'.
- ▶ A failure of EPR locality implies a failure of Bell locality and Einstein locality.
- ▶ If it is assumed there can be causal influences between events only if those events are not separated by a spacelike spacetime interval, then Bell locality reduces to Einstein locality.

# Comparing the three senses of action-at-a-distance

- ▶ Einstein locality imposes the strongest restrictions on locality; EPR locality is the weakest due e.g. to the vagueness of 'do not immediately disturb' and 'sufficiently distant'.
- ▶ A failure of EPR locality implies a failure of Bell locality and Einstein locality.
- ▶ If it is assumed there can be causal influences between events only if those events are not separated by a spacelike spacetime interval, then Bell locality reduces to Einstein locality.
- ▶ Thus, Bell locality can be distinguished from Einstein locality in that it does not explicitly assume relativity theory.

# Separability

- ▶ Unlike locality (which, roughly, is a thesis about the action that one system can exert upon another distant system), *separability* is a metaphysical thesis.

# Separability

- ▶ Unlike locality (which, roughly, is a thesis about the action that one system can exert upon another distant system), *separability* is a metaphysical thesis.
- ▶ Roughly, separability says something like this: *the properties of a system supervene upon the properties of its subsystems.*

# Separability

- ▶ Unlike locality (which, roughly, is a thesis about the action that one system can exert upon another distant system), *separability* is a metaphysical thesis.
- ▶ Roughly, separability says something like this: *the properties of a system supervene upon the properties of its subsystems*.
- ▶ There is clearly, then, a strong connection between quantum entanglement and non-separability.

# Separability

- ▶ Unlike locality (which, roughly, is a thesis about the action that one system can exert upon another distant system), *separability* is a metaphysical thesis.
- ▶ Roughly, separability says something like this: *the properties of a system supervene upon the properties of its subsystems*.
- ▶ There is clearly, then, a strong connection between quantum entanglement and non-separability.
- ▶ That said, some authors (e.g. Ney (2021), see Lecture 16) do try to maintain that one can have entanglement without non-separability.

# Today

Locality and separability

The Einstein-Podolsky-Rosen argument

Bell's 1964 theorem



# The Einstein-Podolsky-Rosen argument

- ▶ In 1935, Einstein, Podolsky and Rosen (EPR) presented a dilemma: *either quantum mechanics is incomplete, or it is non-local.*

# The Einstein-Podolsky-Rosen argument

- ▶ In 1935, Einstein, Podolsky and Rosen (EPR) presented a dilemma: *either quantum mechanics is incomplete, or it is non-local*.
- ▶ EPR took the 'incomplete' horn of the dilemma.

# The Einstein-Podolsky-Rosen argument

- ▶ In 1935, Einstein, Podolsky and Rosen (EPR) presented a dilemma: *either quantum mechanics is incomplete, or it is non-local*.
- ▶ EPR took the 'incomplete' horn of the dilemma.
- ▶ Let's see how their argument goes.

# The EPR dilemma

Consider the singlet two-electron spin state (NB: this is Bohm's (1951) version of the EPR setup),

$$|\psi^-\rangle_{12} = \frac{1}{\sqrt{2}} (|\uparrow_z\rangle_1 |\downarrow_z\rangle_2 - |\downarrow_z\rangle_1 |\uparrow_z\rangle_2).$$

# The EPR dilemma

Consider the singlet two-electron spin state (NB: this is Bohm's (1951) version of the EPR setup),

$$|\psi^-\rangle_{12} = \frac{1}{\sqrt{2}} (|\uparrow_z\rangle_1 |\downarrow_z\rangle_2 - |\downarrow_z\rangle_1 |\uparrow_z\rangle_2).$$

Recalling that

$$|\uparrow_z\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle + |\downarrow_x\rangle),$$

$$|\downarrow_z\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle - |\downarrow_x\rangle),$$

we see that  $|\psi^-\rangle_{12}$  could also be written

# The EPR dilemma

Consider the singlet two-electron spin state (NB: this is Bohm's (1951) version of the EPR setup),

$$|\psi^-\rangle_{12} = \frac{1}{\sqrt{2}} (|\uparrow_z\rangle_1 |\downarrow_z\rangle_2 - |\downarrow_z\rangle_1 |\uparrow_z\rangle_2).$$

Recalling that

$$|\uparrow_z\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle + |\downarrow_x\rangle),$$

$$|\downarrow_z\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle - |\downarrow_x\rangle),$$

we see that  $|\psi^-\rangle_{12}$  could also be written

$$|\psi^-\rangle_{12} = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle_1 |\downarrow_x\rangle_2 - |\downarrow_x\rangle_1 |\uparrow_x\rangle_2)$$

# The EPR dilemma

Consider the singlet two-electron spin state (NB: this is Bohm's (1951) version of the EPR setup),

$$|\psi^-\rangle_{12} = \frac{1}{\sqrt{2}} (|\uparrow_z\rangle_1 |\downarrow_z\rangle_2 - |\downarrow_z\rangle_1 |\uparrow_z\rangle_2).$$

Recalling that

$$|\uparrow_z\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle + |\downarrow_x\rangle),$$

$$|\downarrow_z\rangle = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle - |\downarrow_x\rangle),$$

we see that  $|\psi^-\rangle_{12}$  could also be written

$$|\psi^-\rangle_{12} = \frac{1}{\sqrt{2}} (|\uparrow_x\rangle_1 |\downarrow_x\rangle_2 - |\downarrow_x\rangle_1 |\uparrow_x\rangle_2)$$

(In fact, the singlet is spherically symmetric, i.e. it takes the same form for all spin directions.)

# The EPR dilemma

- Now consider an experiment in which a pair of electrons is prepared in this joint state at one location at time  $t_0$ , before being widely separated.



# The EPR dilemma

- ▶ Now consider an experiment in which a pair of electrons is prepared in this joint state at one location at time  $t_0$ , before being widely separated.
- ▶ At time  $t_1$  the systems 1 and 2 are far apart and no longer interacting and we consider performing measurements on one of them, say system 1.

# The EPR dilemma

- ▶ Now consider an experiment in which a pair of electrons is prepared in this joint state at one location at time  $t_0$ , before being widely separated.
- ▶ At time  $t_1$  the systems 1 and 2 are far apart and no longer interacting and we consider performing measurements on one of them, say system 1.
- ▶ z-spin measurement: before measurement, neither system has a definite value of spin in the z-direction. At  $t > t_1$  we obtain a definite value of spin for system 1. But not only has the property of system 1 changed, but so also have the properties of the far away system 2 (“instantaneous action at a distance”—Einstein: “spooky action”).

# The EPR dilemma

- ▶ Now consider an experiment in which a pair of electrons is prepared in this joint state at one location at time  $t_0$ , before being widely separated.
- ▶ At time  $t_1$  the systems 1 and 2 are far apart and no longer interacting and we consider performing measurements on one of them, say system 1.
- ▶ z-spin measurement: before measurement, neither system has a definite value of spin in the z-direction. At  $t > t_1$  we obtain a definite value of spin for system 1. But not only has the property of system 1 changed, but so also have the properties of the far away system 2 (“instantaneous action at a distance”—Einstein: “spooky action”).
- ▶ Similarly if we had performed an x-spin measurement on system 1 at time  $t > t_1$ . *Our choice of which measurement to perform in the region of system 1 affects at a distance the properties of system 2.*

# Two possibilities

So two possibilities:

# Two possibilities

So two possibilities:

Either: Quantum mechanics is non-local: the effect of measurement is to cause a collapse which has an instantaneous causal effect at a distance.

Or: Quantum mechanics is incomplete: the genuine properties of systems should not behave like this, which means that the quantum state cannot be the whole story. (This was EPR's preferred horn of the dilemma.)

# Two possibilities

So two possibilities:

Either: Quantum mechanics is non-local: the effect of measurement is to cause a collapse which has an instantaneous causal effect at a distance.

Or: Quantum mechanics is incomplete: the genuine properties of systems should not behave like this, which means that the quantum state cannot be the whole story. (This was EPR's preferred horn of the dilemma.)

Naturally, the second of these options invites thinking about quantum mechanics in terms of hidden variable theories.

# EINSTEIN ATTACKS QUANTUM THEORY

---

Scientist and Two Colleagues  
Find It Is Not 'Complete'  
Even Though 'Correct.'

---

SEE FULLER ONE POSSIBLE

---

Believe a Whole Description of  
'the Physical Reality' Can Be  
Provided Eventually.

# Hidden variables

- ▶ Recall that hidden variables are additional ‘elements of reality’ about whose values we are ignorant (hence ‘hidden’) and which explain why we get the particular outcomes we do when we perform (quantum) experiments.



# Hidden variables

- ▶ Recall that hidden variables are additional ‘elements of reality’ about whose values we are ignorant (hence ‘hidden’) and which explain why we get the particular outcomes we do when we perform (quantum) experiments.
- ▶ On hidden variable theories, the statistical features of quantum mechanics just arise as ‘ignorance’ probabilities regarding the hidden variables.

# Hidden variables

- ▶ Recall that hidden variables are additional ‘elements of reality’ about whose values we are ignorant (hence ‘hidden’) and which explain why we get the particular outcomes we do when we perform (quantum) experiments.
- ▶ On hidden variable theories, the statistical features of quantum mechanics just arise as ‘ignorance’ probabilities regarding the hidden variables.
- ▶ In general, incompatible properties such as position and momentum *are* definite at the same time, it’s just that we are *ignorant* of what their values are.

# Hidden variables

- ▶ Recall that hidden variables are additional ‘elements of reality’ about whose values we are ignorant (hence ‘hidden’) and which explain why we get the particular outcomes we do when we perform (quantum) experiments.
- ▶ On hidden variable theories, the statistical features of quantum mechanics just arise as ‘ignorance’ probabilities regarding the hidden variables.
- ▶ In general, incompatible properties such as position and momentum *are* definite at the same time, it’s just that we are *ignorant* of what their values are.
- ▶ Introducing hidden variables might help us with the measurement problem: the value of the hidden variable will determine which outcome of the experiment is definite, even if the *quantum state* happens to be left in a superposition of distinct measurement outcomes.

# Hidden variables

- ▶ Recall that hidden variables are additional ‘elements of reality’ about whose values we are ignorant (hence ‘hidden’) and which explain why we get the particular outcomes we do when we perform (quantum) experiments.
- ▶ On hidden variable theories, the statistical features of quantum mechanics just arise as ‘ignorance’ probabilities regarding the hidden variables.
- ▶ In general, incompatible properties such as position and momentum *are* definite at the same time, it’s just that we are *ignorant* of what their values are.
- ▶ Introducing hidden variables might help us with the measurement problem: the value of the hidden variable will determine which outcome of the experiment is definite, even if the *quantum state* happens to be left in a superposition of distinct measurement outcomes.
- ▶ Of course, we saw a lot of this already in Lecture 6 in the context of Bohmian mechanics.

# Today

Locality and separability

The Einstein-Podolsky-Rosen argument

Bell's 1964 theorem

# Enter Bell



# Bell's 1964 theorem

- ▶ In 1964 the landscape changed: *embracing incompleteness cannot avoid non-locality.*

# Bell's 1964 theorem

- ▶ In 1964 the landscape changed: *embracing incompleteness cannot avoid non-locality*.
- ▶ Bell (1964) made the discovery that if a hidden variable theory is to be consistent with the empirical predictions of quantum mechanics, it *must* be non-local ('Bell's theorem').



# Bell's 1964 theorem

- ▶ In 1964 the landscape changed: *embracing incompleteness cannot avoid non-locality*.
- ▶ Bell (1964) made the discovery that if a hidden variable theory is to be consistent with the empirical predictions of quantum mechanics, it *must* be non-local ('Bell's theorem').
- ▶ Therefore, taking the incompleteness horn of the EPR dilemma does not lead us away from non-locality after all.

# Bell's 1964 theorem

- ▶ In 1964 the landscape changed: *embracing incompleteness cannot avoid non-locality*.
- ▶ Bell (1964) made the discovery that if a hidden variable theory is to be consistent with the empirical predictions of quantum mechanics, it *must* be non-local ('Bell's theorem').
- ▶ Therefore, taking the incompleteness horn of the EPR dilemma does not lead us away from non-locality after all.
- ▶ Today, I'll show you how Bell's 1964 theorem works; in the next lecture, I'll present some later, arguably more powerful (and intricate) versions of Bell's theorem.

# Setup for the 1964 result

- ▶ Begin by considering a deterministic theory which ascribes definite values to the results of all experiments we might perform on a system ('deterministic hidden variable theory').

# Setup for the 1964 result

- ▶ Begin by considering a deterministic theory which ascribes definite values to the results of all experiments we might perform on a system ('deterministic hidden variable theory').
- ▶ Specifying the initial state leads to deterministic predictions about what the results of any measurement will be.

# Setup for the 1964 result

- ▶ Begin by considering a deterministic theory which ascribes definite values to the results of all experiments we might perform on a system ('deterministic hidden variable theory').
- ▶ Specifying the initial state leads to deterministic predictions about what the results of any measurement will be.
- ▶ Assume, furthermore, that the theory satisfies a locality condition: the values for the outcomes of experiments on a system depend only on the initial state and on the particular experiment performed on that system.

# The aim of the game

- ▶ The aim is to show that the statistical predictions of any such theory will satisfy a certain inequality (a 'Bell inequality') regarding the correlations in EPR-type experiments.

# The aim of the game

- ▶ The aim is to show that the statistical predictions of any such theory will satisfy a certain inequality (a 'Bell inequality') regarding the correlations in EPR-type experiments.
- ▶ Such an inequality will be violated by the predictions of quantum mechanics.

# The aim of the game

- ▶ The aim is to show that the statistical predictions of any such theory will satisfy a certain inequality (a 'Bell inequality') regarding the correlations in EPR-type experiments.
- ▶ Such an inequality will be violated by the predictions of quantum mechanics.
- ▶ Conclusion: The results predicted by quantum mechanics cannot be modelled by any *local* deterministic hidden variable theory.



# The setup for Bell's theorem

- We will consider an experiment testing the correlations between various spin measurements on a pair of spin-half systems prepared in an initial entangled singlet state (Bohm's version of the EPR setup).

# The setup for Bell's theorem

- ▶ We will consider an experiment testing the correlations between various spin measurements on a pair of spin-half systems prepared in an initial entangled singlet state (Bohm's version of the EPR setup).
- ▶ Measurements of spin will be made in the direction  $\mathbf{a}$  or  $\mathbf{a}'$  on system 1; and in the direction  $\mathbf{b}$  or  $\mathbf{b}'$  on system 2.

# The setup for Bell's theorem

- ▶ We will consider an experiment testing the correlations between various spin measurements on a pair of spin-half systems prepared in an initial entangled singlet state (Bohm's version of the EPR setup).
- ▶ Measurements of spin will be made in the direction  $\mathbf{a}$  or  $\mathbf{a}'$  on system 1; and in the direction  $\mathbf{b}$  or  $\mathbf{b}'$  on system 2.
- ▶ The hidden variable theory will assign definite values to all four spin quantities at the same time, i.e.  $\mathbf{a}$  and  $\mathbf{a}'$  for system 1 possess a definite value just before measurement, as do  $\mathbf{b}$  and  $\mathbf{b}'$ .

# The setup for Bell's theorem

- Denote these definite values of quantities by  $a$ ,  $a'$ ,  $b$ ,  $b'$ , respectively. They represent the values that *would* be observed if one *were* to perform the measurement of that quantity on the given system.

# The setup for Bell's theorem

- ▶ Denote these definite values of quantities by  $a, a', b, b'$ , respectively. They represent the values that *would* be observed if one *were* to perform the measurement of that quantity on the given system.
- ▶ On each run of the experiment, all four spin quantities possess a definite value. For run  $n$ , denote these quantities by  $a_n, a'_n, b_n, b'_n$ .

# The setup for Bell's theorem

- ▶ Denote these definite values of quantities by  $a$ ,  $a'$ ,  $b$ ,  $b'$ , respectively. They represent the values that *would* be observed if one *were* to perform the measurement of that quantity on the given system.
- ▶ On each run of the experiment, all four spin quantities possess a definite value. For run  $n$ , denote these quantities by  $a_n$ ,  $a'_n$ ,  $b_n$ ,  $b'_n$ .
- ▶ Although the initial *quantum* state of the joint system is the same for each run, the hidden variable takes different values and therefore a spread of outcomes is observed.

# Average values, correlation coefficients, and $\gamma_n$

Average values and correlation coefficients are given by, respectively,

$$\bar{a} = \frac{1}{N} \sum_{n=1}^N a_n,$$

$$c(\mathbf{a}, \mathbf{b}) = \frac{1}{N} \sum_{n=1}^N a_n b_n.$$

# Average values, correlation coefficients, and $\gamma_n$

Average values and correlation coefficients are given by, respectively,

$$\bar{a} = \frac{1}{N} \sum_{n=1}^N a_n,$$
$$c(\mathbf{a}, \mathbf{b}) = \frac{1}{N} \sum_{n=1}^N a_n b_n.$$

Now consider the following function:

$$\begin{aligned}\gamma_n &:= a_n b_n + a_n b'_n + a'_n b_n - a'_n b'_n \\ &= a_n (b_n + b'_n) + a'_n (b_n - b'_n).\end{aligned}$$

Since each of  $a$ ,  $a'$ ,  $b$ ,  $b'$  can be equal to  $\pm 1$ ,  $\gamma_n$  must be equal to  $\pm 2$ .



# Deriving the Bell inequality

Now calculate the average value of  $\gamma_n$  for a large number of runs of the experiment,

$$\frac{1}{N} \sum_{n=1}^N \gamma_n = \frac{1}{N} \sum_{n=1}^N (a_n b_n + a_n b'_n + a'_n b_n - a'_n b'_n) .$$

# Deriving the Bell inequality

Now calculate the average value of  $\gamma_n$  for a large number of runs of the experiment,

$$\frac{1}{N} \sum_{n=1}^N \gamma_n = \frac{1}{N} \sum_{n=1}^N (a_n b_n + a_n b'_n + a'_n b_n - a'_n b'_n).$$

Given that on each run  $\gamma_n = \pm 2$ , this average must lie between  $\pm 2$ , so its modulus will be less than or equal to 2:

$$\left| \frac{1}{N} \sum_{n=1}^N (a_n b_n + a_n b'_n + a'_n b_n - a'_n b'_n) \right| \leq 2.$$

# Deriving the Bell inequality

Now calculate the average value of  $\gamma_n$  for a large number of runs of the experiment,

$$\frac{1}{N} \sum_{n=1}^N \gamma_n = \frac{1}{N} \sum_{n=1}^N (a_n b_n + a_n b'_n + a'_n b_n - a'_n b'_n).$$

Given that on each run  $\gamma_n = \pm 2$ , this average must lie between  $\pm 2$ , so its modulus will be less than or equal to 2:

$$\left| \frac{1}{N} \sum_{n=1}^N (a_n b_n + a_n b'_n + a'_n b_n - a'_n b'_n) \right| \leq 2.$$

In terms of correlation coefficients, this gives us the *Bell inequality*:

$$|c(\mathbf{a}, \mathbf{b}) + c(\mathbf{a}, \mathbf{b}') + c(\mathbf{a}', \mathbf{b}) - c(\mathbf{a}', \mathbf{b}')| \leq 2.$$

# Comparison with standard quantum mechanics

How does this compare with the predictions that standard quantum mechanics makes for the correlations between spin measurement outcomes that would be observed in the EPR-type experiment?

# Comparison with standard quantum mechanics

How does this compare with the predictions that standard quantum mechanics makes for the correlations between spin measurement outcomes that would be observed in the EPR-type experiment?

When the quantum mechanical values for the correlation coefficients are calculated, we instead obtain

$$|c(\mathbf{a}, \mathbf{b}) + c(\mathbf{a}, \mathbf{b}') + c(\mathbf{a}', \mathbf{b}) - c(\mathbf{a}', \mathbf{b}')| \leq 2\sqrt{2}.$$

# Comparison with standard quantum mechanics

How does this compare with the predictions that standard quantum mechanics makes for the correlations between spin measurement outcomes that would be observed in the EPR-type experiment?

When the quantum mechanical values for the correlation coefficients are calculated, we instead obtain

$$|c(\mathbf{a}, \mathbf{b}) + c(\mathbf{a}, \mathbf{b}') + c(\mathbf{a}', \mathbf{b}) - c(\mathbf{a}', \mathbf{b}')| \leq 2\sqrt{2}.$$

That is, the quantum mechanical correlations are *stronger* than any correlations that can be obtained in a local hidden variable model.

# Illustration of the quantum mechanical inequality

This can be illustrated in a special case—suppose for example that:

# Illustration of the quantum mechanical inequality

This can be illustrated in a special case—suppose for example that:

1. Measurements of **a** and **b** are along the same axis.
2. Measurements of **a** and **b'** or **b** and **a'** are at an angle  $\theta$ .
3. Measurements of **a'** and **b'** are at an angle  $2\theta$ .



# Illustration of the quantum mechanical inequality

This can be illustrated in a special case—suppose for example that:

1. Measurements of **a** and **b** are along the same axis.
2. Measurements of **a** and **b'** or **b** and **a'** are at an angle  $\theta$ .
3. Measurements of **a'** and **b'** are at an angle  $2\theta$ .

Quantum mechanics has its own predictions of correlation coefficients. If  $\phi$  is the angle between directions **a** and **b**, then  $c(\mathbf{a}, \mathbf{b}) = \cos \phi$ .

# Illustration of the quantum mechanical inequality

Using the above assumptions, we therefore have

$$\begin{aligned} |c(\mathbf{a}, \mathbf{b}) + c(\mathbf{a}, \mathbf{b}') + c(\mathbf{a}', \mathbf{b}) - c(\mathbf{a}', \mathbf{b}')| &= |\cos(0) + 2\cos(\theta) - \cos(2\theta)| \\ &= |2(1 + \cos\theta - \cos^2\theta)| \\ &> 2 \end{aligned}$$

# Illustration of the quantum mechanical inequality

Using the above assumptions, we therefore have

$$\begin{aligned} |c(\mathbf{a}, \mathbf{b}) + c(\mathbf{a}, \mathbf{b}') + c(\mathbf{a}', \mathbf{b}) - c(\mathbf{a}', \mathbf{b}')| &= |\cos(0) + 2\cos(\theta) - \cos(2\theta)| \\ &= |2(1 + \cos\theta - \cos^2\theta)| \\ &> 2 \end{aligned}$$

(Note: It's not that standard quantum mechanics predicts different *inequalities* to local hidden variable theories. It's that some of its specific predictions involve correlations that violate the Bell inequalities.)

# Illustration of the quantum mechanical inequality

Using the above assumptions, we therefore have

$$\begin{aligned} |c(\mathbf{a}, \mathbf{b}) + c(\mathbf{a}, \mathbf{b}') + c(\mathbf{a}', \mathbf{b}) - c(\mathbf{a}', \mathbf{b}')| &= |\cos(0) + 2\cos(\theta) - \cos(2\theta)| \\ &= |2(1 + \cos\theta - \cos^2\theta)| \\ &> 2 \end{aligned}$$

(Note: It's not that standard quantum mechanics predicts different *inequalities* to local hidden variable theories. It's that some of its specific predictions involve correlations that violate the Bell inequalities.)

*So no local hidden variable model can be empirically equivalent to quantum mechanics!*

# Empirical evidence for standard quantum mechanics

- ▶ Which is right—quantum mechanics, or local hidden variable theories?

# Empirical evidence for standard quantum mechanics

- ▶ Which is right—quantum mechanics, or local hidden variable theories?
- ▶ We need to do the experiment and see whether the quantum predictions or the local hidden variable predictions for the correlations hold.

# Empirical evidence for standard quantum mechanics

- ▶ Which is right—quantum mechanics, or local hidden variable theories?
- ▶ We need to do the experiment and see whether the quantum predictions or the local hidden variable predictions for the correlations hold.
- ▶ The overwhelming evidence is that *standard quantum mechanics is correct*.

# Empirical evidence for standard quantum mechanics

- ▶ Which is right—quantum mechanics, or local hidden variable theories?
- ▶ We need to do the experiment and see whether the quantum predictions or the local hidden variable predictions for the correlations hold.
- ▶ The overwhelming evidence is that *standard quantum mechanics is correct*.
- ▶ The most famous of these experiments is Aspect's experiment, which won the Nobel Prize in Physics in 2022.



# Empirical evidence for standard quantum mechanics

- ▶ Which is right—quantum mechanics, or local hidden variable theories?
- ▶ We need to do the experiment and see whether the quantum predictions or the local hidden variable predictions for the correlations hold.
- ▶ The overwhelming evidence is that *standard quantum mechanics is correct*.
- ▶ The most famous of these experiments is Aspect's experiment, which won the Nobel Prize in Physics in 2022.
- ▶ For a more comprehensive list of tests, see e.g. (Redhead 1987, p. 108) or (Myrvold *et al.* 2024).

# Empirical evidence for standard quantum mechanics

- ▶ Which is right—quantum mechanics, or local hidden variable theories?
- ▶ We need to do the experiment and see whether the quantum predictions or the local hidden variable predictions for the correlations hold.
- ▶ The overwhelming evidence is that *standard quantum mechanics is correct*.
- ▶ The most famous of these experiments is Aspect's experiment, which won the Nobel Prize in Physics in 2022.
- ▶ For a more comprehensive list of tests, see e.g. (Redhead 1987, p. 108) or (Myrvold *et al.* 2024).

*So no local hidden variable model is empirically adequate!*

# Conclusions so far

- ▶ Thus we conclude that any acceptable deterministic hidden variable theory must be non-local.

# Conclusions so far

- ▶ Thus we conclude that any acceptable deterministic hidden variable theory must be non-local.
- ▶ But note that this doesn't imply anything about *quantum mechanics* being non-local, just that any hidden variable theory put in its place must be.

# Conclusions so far

- ▶ Thus we conclude that any acceptable deterministic hidden variable theory must be non-local.
- ▶ But note that this doesn't imply anything about *quantum mechanics* being non-local, just that any hidden variable theory put in its place must be.
- ▶ We know that collapse in quantum mechanics gives rise to action-at-a-distance, but what if we deny that collapse is a real physical process?

# Conclusions so far

- ▶ Thus we conclude that any acceptable deterministic hidden variable theory must be non-local.
- ▶ But note that this doesn't imply anything about *quantum mechanics* being non-local, just that any hidden variable theory put in its place must be.
- ▶ We know that collapse in quantum mechanics gives rise to action-at-a-distance, but what if we deny that collapse is a real physical process?

NB: Bell's theorem does not force us to accept “the bare formalism of quantum mechanics” *sans* interpretation, as some (physics texts) claim. We still need to solve the measurement problem!

# Prospectus

Next time, I'll:

# Prospectus

Next time, I'll:

1. Present later versions of Bell's theorem.











# Prospectus

Next time, I'll:

1. Present later versions of Bell's theorem.
2. Think about what different interpretations of quantum mechanics have to say about the EPR scenario.

# References

-  John S. Bell, “On the Einstein–Podolsky–Rosen Paradox”, *Physics* 1, pp. 195–200, 1964.
-  John S. Bell, “Bertlmann’s Socks and the Nature of Reality”, in *Speakable and Unspeakable in Quantum Mechanics*, Cambridge: Cambridge University Press, 1987.
-  David Bohm, *Quantum Theory*, Englewood Cliffs, NJ: Prentice-Hall, 1951.
-  A. Einstein, B. Podolsky and N. Rosen, “Can Quantum-Mechanical Description of Physical Reality Be Considered Complete?”, *Physical Review* 47(10), pp. 777–80, 1935.
-  Tim Maudlin, “What Bell Did”, *Journal of Physics A: Mathematical and Theoretical* 47, 424010, 2014.
-  Wayne Myrvold, Marco Genovese and Abner Shimony, “Bell’s Theorem”, in E. N. Zalta (ed.), *The Stanford Encyclopedia of Philosophy*, 2024.
-  Alyssa Ney, *The World in the Wave Function*, Oxford: Oxford University Press, 2021.
-  Michael Redhead, *Incompleteness, Nonlocality and Realism: A Prolegomenon to the Philosophy of Quantum Mechanics*, Oxford: Clarendon Press, 1987.