


Syntax, Truth,
Paradox:

A map through
the land of dragons

Volker Halbach & Lorenzo Rossi

Lugano · 17 to 24 June 2024



Prelude: The paradoxes in philosophy


Day 1: A theory of expressions

Day 2: The paradoxes

Day 3: Possible-worlds analysis of the paradoxes

Day 4: Truth!

THE PARADOXES



The liar and other simple paradoxes

Paradoxes from Interaction

Yablo and Visser

methodology

McGee's ω -inconsistency

bibliography

For later reference here are the axioms again:

DEFINITION

All instances of the following schemas are axioms of E:

A1 $\bar{a} \wedge \bar{b} = \overline{ab}$, where a and b are arbitrary strings of symbols

A2 $q(\bar{a}) = \overline{\bar{a}}$

A3 $\text{sub}(\bar{a}, \bar{b}, \bar{c}) = \bar{d}$, where a and c are arbitrary strings of symbols, b is a symbol (or, equivalently, a length-1 string of symbols), and d is the string of symbols obtained from a by replacing all occurrences of the symbol b with c

A4 $\forall x \forall y \forall z ((x \wedge y) \wedge z) = (x \wedge (y \wedge z))$

A5 $\forall x \forall y (x \wedge y = \underline{0} \rightarrow x = \underline{0} \wedge y = \underline{0})$

A6 $\forall x \forall y (x \wedge y = x \leftrightarrow y = \underline{0}) \wedge \forall x \forall y (y \wedge x = x \leftrightarrow y = \underline{0})$

A7 $\forall x \forall y \text{sub}(x \wedge \bar{a}, \bar{a}, y) = \text{sub}(x, \bar{a}, y) \wedge y$, where a is a symbol

A8 $\forall x \forall y \forall z \forall w$

$(x \wedge y = z \wedge w \leftrightarrow \exists v_4 ((x = z \wedge v_4 \wedge v_4 \wedge y = w) \vee (x \wedge v_4 = z \wedge y = v_4 \wedge w)))$

The liar and other simple paradoxes

The theory E can contain more axioms beyond those explicitly stated.

Thus the following two claims are equivalent:

- (i) φ is inconsistent with E.
- (ii) E is inconsistent if it contains φ .

Instead of saying ‘ φ is inconsistent with E’ I will often only say ‘ φ is inconsistent’.

I do not assume that E is consistent. It is consistent with the stated axioms.

THEOREM (LIAR PARADOX)

The T-schema $\Box\bar{\psi} \leftrightarrow \psi$ for all sentences ψ of \mathcal{L} is inconsistent.

Proof: Apply the diagonal lemma to $\neg\Box v$.

THEOREM (TARSKI'S THEOREM ON THE UNDEFINABILITY OF TRUTH)

If E is consistent, there is no formula $\tau(x)$ such that $\tau(\bar{\psi}) \leftrightarrow \psi$ can be derived in E for all sentences ψ of \mathcal{L} .

However, we can axiomatically add a new truth predicate.

McGee (1992) proved:

THEOREM (MCGEE'S THEOREM ON T-SENTENCES)

Assume $E \not\vdash \neg\varphi$. Then there is a sentence γ such that $E + (\Box\bar{\gamma} \leftrightarrow \gamma)$ is consistent and $E + (\Box\bar{\gamma} \leftrightarrow \gamma) \vdash \varphi$.

Proof: Apply the diagonal lemma to $\Box v \leftrightarrow \varphi$.

This is a variant of Curry's paradox (Curry 1942). McGee used it against a 'solution' of the truth-theoretic paradoxes in (Horwich 1998).

THEOREM (MONTAGUE'S PARADOX, MONTAGUE 1963)

The schema $\Box\bar{\varphi} \rightarrow \varphi$ is inconsistent with the rule

(NEC)
$$\frac{\varphi}{\Box\bar{\varphi}}$$

for all sentences φ .

Which modal notions are affected?

THEOREM (MONTAGUE'S PARADOX, MONTAGUE 1963)

The schema $\Box\bar{\varphi} \rightarrow \varphi$ is inconsistent with the rule

(NEC)
$$\frac{\varphi}{\Box\bar{\varphi}}$$

for all sentences φ .

Which modal notions are affected?

THEOREM (GÖDEL'S FIRST INCOMPLETENESS THEOREM)

Suppose for all sentences φ of \mathcal{L} that $E \vdash \varphi$ iff $E \vdash \Box\bar{\varphi}$. Then there is a sentence γ such that neither γ itself nor its negation is derivable in E , or E is inconsistent.

This is just extracted from (Gödel 1931).

THEOREM

E is inconsistent if it contains the schema $\Box\Box\overline{\varphi} \leftrightarrow \varphi$.

PROOF.

For any for all sentences φ of \mathcal{L} and indeed any expression φ we have the following:

$$E \vdash \Box \wedge q(\overline{\varphi}) = \Box \wedge \overline{\overline{\varphi}} \quad \text{axiom A2}$$

$$E \vdash \Box \wedge q(\overline{\varphi}) = \overline{\Box\overline{\varphi}} \quad \text{axiom A1}$$

The diagonal lemma is applied to the formula $\neg\Box(\Box \wedge q(v))$:

$$E \vdash \gamma \leftrightarrow \neg\Box(\Box \wedge q(\overline{\gamma}))$$

$$E \vdash \gamma \leftrightarrow \neg\Box\overline{\overline{\gamma}} \quad \text{remark above}$$

$$E \vdash \gamma \leftrightarrow \neg\gamma \quad \text{assumption}$$

$$(K) \quad \Box \overline{\varphi \rightarrow \psi} \rightarrow (\Box \overline{\varphi} \rightarrow \Box \overline{\psi}).$$

For our abbreviation of \wedge we have:

LEMMA

$E \vdash \Box \overline{\varphi \wedge \psi} \leftrightarrow \Box \overline{\varphi} \wedge \Box \overline{\psi}$, if E contains K and is closed under NEC.

A theory \mathcal{T} is internally inconsistent (with respect to \Box) if and only if $\mathcal{T} \vdash \Box \bar{\varphi}$ and $\mathcal{T} \vdash \Box \overline{\neg \varphi}$ for some sentence φ .

LEMMA

Assume E is closed under NEC. Then every internally inconsistent theory containing E and K proves $\Box \bar{\psi}$ for all sentences ψ .

is the schema $\Box\bar{\varphi} \rightarrow \overline{\Box\bar{\varphi}}$.

This is similar to (Thomason 1980):

THEOREM (THOMASON 1980)

Assume E is closed under NEC. Then any theory \mathcal{T} containing E and the schemas K , 4, and $\overline{\Box\bar{\varphi} \rightarrow \varphi}$ is internally inconsistent and proves $\Box\bar{\psi}$ for all \mathcal{L} -sentences ψ .

Read the modal predicate as 'S knows x '.

Proof:

Roughly speaking, we run the proof of Montague's paradox in the scope of \Box . Assume that \mathcal{E} and \mathcal{T} have the properties mentioned in the theorem.

$\mathcal{E} \vdash \gamma \leftrightarrow \neg\Box\bar{\gamma}$	liar sentence
$\mathcal{E} \vdash \overline{\Box\Box\bar{\gamma} \rightarrow \neg\gamma}$	logic and NEC
$\mathcal{E} \vdash (\Box\bar{\gamma} \rightarrow \gamma) \rightarrow ((\Box\bar{\gamma} \rightarrow \neg\gamma) \rightarrow \neg\Box\bar{\gamma})$	logic
$\mathcal{E} \vdash (\Box\bar{\gamma} \rightarrow \gamma) \rightarrow ((\Box\bar{\gamma} \rightarrow \neg\gamma) \rightarrow \gamma)$	first line
$\mathcal{E} \vdash \overline{\Box\Box\bar{\gamma} \rightarrow \gamma} \rightarrow (\overline{\Box\Box\bar{\gamma} \rightarrow \neg\gamma} \rightarrow \Box\bar{\gamma})$	NEC and K
$\mathcal{T} \vdash (\overline{\Box\Box\bar{\gamma} \rightarrow \neg\gamma} \rightarrow \Box\bar{\gamma})$	$\overline{\Box\Box\bar{\varphi} \rightarrow \varphi}$
$\mathcal{T} \vdash \Box\bar{\gamma}$	second line

Now we invoke 4 to conclude $\mathcal{T} \vdash \overline{\square \bar{\gamma}}$ from $\mathcal{T} \vdash \square \bar{\gamma}$. From the first line above we also get $E \vdash \square \bar{\gamma} \rightarrow \overline{\square \bar{\gamma}}$ by NEC and K. Combining this with the last line, we obtain the following internal inconsistency:

$$\mathcal{T} \vdash \overline{\square \bar{\gamma}} \wedge \square \bar{\gamma}.$$

Since E and \mathcal{T} satisfy the conditions of 9, we have $\mathcal{T} \vdash \square \bar{\psi}$ for all sentences ψ .

THEOREM (LÖB'S THEOREM)

If E is closed under NEC and contains K and 4 , then we have
 $E \vdash \overline{\Box\Box\bar{\varphi} \rightarrow \varphi} \rightarrow \Box\bar{\varphi}$ *for all sentences φ of \mathcal{L} .*

Then there is the mystery of the de Jongh–Sambin fixed-point theorem.

THEOREM (LÖB'S THEOREM)

If E is closed under NEC and contains K and 4 , then we have
 $E \vdash \overline{\Box\Box\bar{\varphi} \rightarrow \varphi} \rightarrow \Box\bar{\varphi}$ *for all sentences φ of \mathcal{L} .*

Then there is the mystery of the de Jongh–Sambin fixed-point theorem.

COROLLARY

Assume that E is closed under NEC and contains K and 4 . Then for any φ of \mathcal{L} the following rule of inference holds: If $E \vdash \Box\bar{\varphi} \rightarrow \varphi$, then $E \vdash \varphi$.

PROOF.

Assume $E \vdash \Box\bar{\varphi} \rightarrow \varphi$ and reason as follows:

$E \vdash \Box\bar{\varphi} \rightarrow \varphi$	assumption
$E \vdash \Box\overline{\Box\bar{\varphi} \rightarrow \varphi}$	NEC
$E \vdash \Box\bar{\varphi}$	theorem above
$E \vdash \varphi$	assumption in first line



THEOREM (GÖDEL'S SECOND INCOMPLETENESS THEOREM)

Assume that E is closed under NEC and contains K and 4 . Then $E \vdash \neg \Box \perp$ implies that E is inconsistent.

Paradoxes from Interaction

Assume Tr is also a primitive unary predicate in the language.

THEOREM

Assume E satisfies the following three conditions:

- (i) If ψ is a sentence of \mathcal{L} not containing Tr , then E contains $\text{Tr}\bar{\psi} \leftrightarrow \psi$.
- (ii) If ψ is a sentence of \mathcal{L} not containing \Box , then E contains $\Box\bar{\psi} \rightarrow \psi$.
- (iii) If ψ is a sentence of \mathcal{L} not containing \Box with $E \vdash \psi$, then also $E \vdash \Box\bar{\psi}$ holds.

Then E is inconsistent.

PROOF.

We apply the diagonal lemma to the formula $\neg\text{Tr}(\bar{\square} \wedge q(x))$ and reason as follows:

$E \vdash \gamma \leftrightarrow \neg\text{Tr}(\bar{\square} \wedge q(\bar{\gamma}))$	diagonal lemma
$E \vdash \gamma \leftrightarrow \neg\text{Tr}\bar{\square}\bar{\gamma}$	axioms A1 and A2; cf. proof of 7
$E \vdash \text{Tr}\bar{\square}\bar{\gamma} \rightarrow \neg\gamma$	logic
$E \vdash \square\bar{\gamma} \rightarrow \neg\gamma$	(i)
$E \vdash \square\bar{\gamma} \rightarrow \gamma$	(ii)
$E \vdash \neg\square\bar{\gamma}$	two preceding lines
$E \vdash \neg\text{Tr}\bar{\square}\bar{\gamma}$	(i)
$E \vdash \gamma$	second line
$E \vdash \square\bar{\gamma}$	(iii)

The last line and the fourth line from the bottom establish the claim. □

Here is another application, which is not an inconsistency. See (Halbach and Horsten 2025). We assume that we have a predicate K for knowledge and a predicate JB for justified belief.

The knower sentence is a sentence γ with $E \vdash \gamma \leftrightarrow \neg K\bar{\gamma}$.

JTB CONCEPTION OF KNOWLEDGE

The following assumptions are jointly inconsistent:

schematic definition of knowledge: $K\bar{\varphi} \leftrightarrow (JB\bar{\varphi} \wedge \varphi)$

factivity of knowledge: $K\bar{\varphi} \rightarrow \varphi$

Crucial assumption: From a proof of φ conclude $JB\bar{\varphi}$

	$K\bar{\gamma} \rightarrow \gamma$	factivity
	$K\bar{\gamma} \rightarrow \neg\gamma$	knower
(1)	$\neg K\bar{\gamma}$	two preceding lines
(2)	γ	knower
(3)	$JB\bar{\gamma}$	crucial assumption
	$JB\bar{\gamma} \wedge \gamma$	from (2) and (3)
(4)	$K\bar{\gamma}$	def. of knowledge

JTB CONCEPTION OF KNOWLEDGE

The following assumptions are jointly inconsistent:

schematic definition of knowledge: $K\bar{\varphi} \leftrightarrow (JB\bar{\varphi} \wedge \varphi)$

factivity of knowledge: $K\bar{\varphi} \rightarrow \varphi$

Crucial assumption: From a proof of φ conclude $JB\bar{\varphi}$

	$K\bar{\gamma} \rightarrow \gamma$	factivity
	$K\bar{\gamma} \rightarrow \neg\gamma$	knower
(1)	$\neg K\bar{\gamma}$	two preceding lines
(2)	γ	knower
(3)	$JB\bar{\gamma}$	crucial assumption
	$JB\bar{\gamma} \wedge \gamma$	from (2) and (3)
(4)	$K\bar{\gamma}$	def. of knowledge

JTB CONCEPTION OF KNOWLEDGE

The following assumptions are jointly inconsistent:

schematic definition of knowledge: $K\bar{\varphi} \leftrightarrow (JB\bar{\varphi} \wedge \varphi)$

factivity of knowledge: $K\bar{\varphi} \rightarrow \varphi$

Crucial assumption: From a proof of φ conclude $JB\bar{\varphi}$

	$K\bar{\gamma} \rightarrow \gamma$	factivity
	$K\bar{\gamma} \rightarrow \neg\gamma$	knower
(1)	$\neg K\bar{\gamma}$	two preceding lines
(2)	γ	knower
(3)	$JB\bar{\gamma}$	crucial assumption
	$JB\bar{\gamma} \wedge \gamma$	from (2) and (3)
(4)	$K\bar{\gamma}$	def. of knowledge

JTB CONCEPTION OF KNOWLEDGE

The following assumptions are jointly inconsistent:

schematic definition of knowledge: $K\bar{\varphi} \leftrightarrow (JB\bar{\varphi} \wedge \varphi)$

factivity of knowledge: $K\bar{\varphi} \rightarrow \varphi$

Crucial assumption: From a proof of φ conclude $JB\bar{\varphi}$

	$K\bar{\gamma} \rightarrow \gamma$	factivity
	$K\bar{\gamma} \rightarrow \neg\gamma$	knower
(1)	$\neg K\bar{\gamma}$	two preceding lines
(2)	γ	knower
(3)	$JB\bar{\gamma}$	crucial assumption
	$JB\bar{\gamma} \wedge \gamma$	from (2) and (3)
(4)	$K\bar{\gamma}$	def. of knowledge

JTB CONCEPTION OF KNOWLEDGE

The following assumptions are jointly inconsistent:

schematic definition of knowledge: $K\bar{\varphi} \leftrightarrow (JB\bar{\varphi} \wedge \varphi)$

factivity of knowledge: $K\bar{\varphi} \rightarrow \varphi$

Crucial assumption: From a proof of φ conclude $JB\bar{\varphi}$

	$K\bar{\gamma} \rightarrow \gamma$	factivity
	$K\bar{\gamma} \rightarrow \neg\gamma$	knower
(1)	$\neg K\bar{\gamma}$	two preceding lines
(2)	γ	knower
(3)	$JB\bar{\gamma}$	crucial assumption
	$JB\bar{\gamma} \wedge \gamma$	from (2) and (3)
(4)	$K\bar{\gamma}$	def. of knowledge

JTB CONCEPTION OF KNOWLEDGE

The following assumptions are jointly inconsistent:

schematic definition of knowledge: $K\bar{\varphi} \leftrightarrow (JB\bar{\varphi} \wedge \varphi)$

factivity of knowledge: $K\bar{\varphi} \rightarrow \varphi$

Crucial assumption: From a proof of φ conclude $JB\bar{\varphi}$

	$K\bar{\gamma} \rightarrow \gamma$	factivity
	$K\bar{\gamma} \rightarrow \neg\gamma$	knower
(1)	$\neg K\bar{\gamma}$	two preceding lines
(2)	γ	knower
(3)	$JB\bar{\gamma}$	crucial assumption
	$JB\bar{\gamma} \wedge \gamma$	from (2) and (3)
(4)	$K\bar{\gamma}$	def. of knowledge

JTB CONCEPTION OF KNOWLEDGE

The following assumptions are jointly inconsistent:

schematic definition of knowledge: $K\bar{\varphi} \leftrightarrow (JB\bar{\varphi} \wedge \varphi)$

factivity of knowledge: $K\bar{\varphi} \rightarrow \varphi$

Crucial assumption: From a proof of φ conclude $JB\bar{\varphi}$

	$K\bar{\gamma} \rightarrow \gamma$	factivity
	$K\bar{\gamma} \rightarrow \neg\gamma$	knower
(1)	$\neg K\bar{\gamma}$	two preceding lines
(2)	γ	knower
(3)	$JB\bar{\gamma}$	crucial assumption
	$JB\bar{\gamma} \wedge \gamma$	from (2) and (3)
(4)	$K\bar{\gamma}$	def. of knowledge

Gx is 'x will always be the case', while Hx is 'x has always been the case'.
 Defined: Fx is 'x will be the case at some point (in the future)', and Px ,
 stands for 'x has been the case at some point (in the past)'.

$$(G1) \quad G\overline{\varphi \rightarrow \psi} \rightarrow (G\overline{\varphi} \rightarrow G\overline{\psi}),$$

$$(H1) \quad H\overline{\varphi \rightarrow \psi} \rightarrow (H\overline{\varphi} \rightarrow H\overline{\psi}),^1$$

$$(G2) \quad \varphi \rightarrow H\overline{F\overline{\varphi}},$$

$$(H2) \quad \varphi \rightarrow G\overline{P\overline{\varphi}},$$

$$(G3) \quad G\overline{\varphi} \leftrightarrow \neg F\overline{\neg\varphi},$$

$$(H3) \quad H\overline{\varphi} \leftrightarrow \neg P\overline{\neg\varphi},$$

$$(N) \quad \frac{\varphi}{G\overline{\varphi}} \text{ and } \frac{\varphi}{H\overline{\varphi}} \text{ for all sentences } \varphi.$$

¹In the original paper (Horsten and Leitgeb 2001, p. 260), there is a typo in the formulation of this axiom: the occurrence of G there should be an H , too.

THEOREM (NO FUTURE PARADOX, HORSTEN AND LEITGEB 2001)

If E contains $G_1, H_1, G_2, H_2, G_3,$ and H_3 and is closed under N , we have

$E \vdash H\bar{1} \wedge G\bar{1}.$

PROOF.

I shall only prove that there is no future, that is, $E \vdash G\bar{1}$. The first line is obtained as in the proof of 14:

- (5) $E \vdash \gamma \leftrightarrow \overline{G\overline{P\overline{\neg\gamma}}}$
 $E \vdash \neg\gamma \leftrightarrow \neg\overline{G\overline{P\overline{\neg\gamma}}}$
 $E \vdash \neg\gamma \rightarrow \overline{G\overline{P\overline{\neg\gamma}}}$ H2
- (6) $E \vdash \gamma$ preceding two lines
- (7) $E \vdash \overline{G\overline{P\overline{\neg\gamma}}}$ from (5) and previous line
 $E \vdash H\bar{\gamma}$ N and (6)
 $E \vdash \neg P\overline{\neg\gamma}$ H3
- (8) $E \vdash \overline{G\overline{P\overline{\neg\gamma}}}$ N
- (9) $E \vdash G\bar{1}$ (7), (8), and G1

The last line follows because, by N, we have $\overline{G\varphi \rightarrow (\neg\varphi \rightarrow \perp)}$ for all φ and in particular for $P\overline{\neg\gamma}$. □

Yablo and Visser

Kripke (1975) mentioned something about illfounded hierarchies, but said later in (2019) that this was not about the Yablo–Visser paradox. Visser (1989) and Yablo (1985, 1993) presented similar paradoxes. See also (Cook 2014).

Some authors argued that these are paradoxes without self-reference.

Kripke (1975) mentioned something about illfounded hierarchies, but said later in (2019) that this was not about the Yablo–Visser paradox. Visser (1989) and Yablo (1985, 1993) presented similar paradoxes. See also (Cook 2014).

Some authors argued that these are paradoxes without self-reference.

Yablo (1993) presented his paradox as an infinitely descending list of sentences.

(S₁) for all $k > 1$, S_k is untrue .

(S₂) for all $k > 2$, S_k is untrue .

(S₃) for all $k > 3$, S_k is untrue .

(S₄) for all $k > 4$, S_k is untrue .

⋮

Yablo (1993) presented his paradox as an infinitely descending list of sentences. Visser (1989) presented his paradox in a more formal way, but it is Yablo's paradox with typed truth predicates:

- (S₁) for all $k > 1$, S_k is untrue₁.
- (S₂) for all $k > 2$, S_k is untrue₂.
- (S₃) for all $k > 3$, S_k is untrue₃.
- (S₄) for all $k > 4$, S_k is untrue₄.
- ⋮

The use of infinite lists is dodgy.

The paradoxes can be formulated in syntax theory.

Before we can formulate the paradoxes, I need to explain how we can quantify into quotations.

I will present the paradox with very weak assumptions.

The use of infinite lists is dodgy.

The paradoxes can be formulated in syntax theory.

Before we can formulate the paradoxes, I need to explain how we can quantify into quotations.

I will present the paradox with very weak assumptions.

The use of infinite lists is dodgy.

The paradoxes can be formulated in syntax theory.

Before we can formulate the paradoxes, I need to explain how we can quantify into quotations.

I will present the paradox with very weak assumptions.

Assume \Box is read as 'necessary' and we want to say that every expression is necessarily identical with itself, that is, we want to say that for all expressions e the sentence $\bar{e} = \bar{e}$ is necessary. We cannot do this by writing $\forall x \Box \bar{x} = \bar{x}$, but we can formulate our claim in the following way:

For all expressions e : if we replace in the formula $x = x$ every occurrence of x by the quotation constant for e , then the resulting sentence is necessary.

Assume \Box is read as ‘necessary’ and we want to say that every expression is necessarily identical with itself, that is, we want to say that for all expressions e the sentence $\bar{e} = \bar{e}$ is necessary. We cannot do this by writing $\forall x \Box \bar{x} = \bar{x}$, but we can formulate our claim in the following way:

For all expressions e : if we replace in the formula $x = x$ every occurrence of x by the quotation constant for e , then the resulting sentence is necessary.

This can be expressed in \mathcal{L} using the following formula:

$$\forall x \Box \text{sub}(\overline{x=x}, \bar{x}, q(x)).$$

From this we can derive, for instance, $\Box \overline{\neg = \neg}$ in E in the following way:

$\forall x \Box \text{sub}(\overline{x=x}, \bar{x}, q(x))$	assumption
$\Box \text{sub}(\overline{x=x}, \bar{x}, q(\neg))$	logic
$\Box \text{sub}(\overline{x=x}, \bar{x}, \overline{\neg})$	A2
$\Box \overline{\neg = \neg}$	A3

Assume $\varphi(y)$ is a formula with no bound occurrences of the variable y , then

$$\overline{\varphi(x)} \text{ abbreviates } \text{sub}(\overline{\varphi(y)}, \bar{y}, q(x)).$$

In arithmetic the dot is placed above the variable; but we have already a bar there.

Of course, we can still generalize this and stipulate that only free occurrences of y are replaced; but we don't need this.

The proof of the diagonal lemma yields the following:

DIAGONAL LEMMA WITH FREE VARIABLES

If $\varphi(x, y)$ is a formula of \mathcal{L} with no bound occurrences of x , then one can find a formula $\theta(y)$ such that the following holds:

$$\mathbb{E} \vdash \forall y \left(\theta(y) \leftrightarrow \varphi(\overline{\theta(y)}, y) \right).$$

We want a dot under the overlined y .

Let $\varphi(x, y)$ be a formula with the two free variables x and y that does not contain a bound occurrence of y . Then there is a formula $\theta(y)$ such that

$$E \vdash \forall y \left(\theta(y) \leftrightarrow \varphi(\overline{\theta(y)}, y) \right).$$

PROOF.

By applying the diagonal lemma above to the formula

$$\varphi(\text{sub}(x, \bar{y}, q(y)), y),$$

I obtain a formula $\theta(y)$ such that the following holds:

$$E \vdash \forall y \left(\theta(y) \leftrightarrow \varphi(\text{sub}(\overline{\theta(y)}, \bar{y}, q(y)), y) \right).$$

This is the claim, since $\overline{\theta(y)}$ is defined as $\text{sub}(\overline{\theta(y)}, \bar{y}, q(y))$. □

I prove Yablo's paradox with weak assumptions.

The informal proof is based on an infinitely descending list of sentences with a top element. But much less is needed, as Ketland (2005) has shown.

I assume that the language \mathcal{L} contains a binary predicate symbol $<$ that satisfies the following conditions:

$$\text{(SER)} \quad \forall x \exists y x < y,$$

$$\text{(TRANS)} \quad \forall x \forall y \forall z (x < y \rightarrow (y < z \rightarrow x < z)).$$

That's an order of the entire universe. We don't need it; but I don't want to relativize the quantifiers. We don't need a primitive symbol for this. You can think of $<$ as a defined formula that satisfies these conditions.

Using a primitive symbol shows that nothing beyond SER and TRANS is needed in addition to ...

Of course, we need also a truth-theoretic assumption:

$$(UTS) \quad \forall y (\overline{\Box \varphi(y)} \leftrightarrow \varphi(y)).$$

THEOREM (YABLO'S PARADOX)

Assume that E contains all the following sentences:

$$\text{(SER)} \quad \forall x \exists y x < y,$$

$$\text{(TRANS)} \quad \forall x \forall y \forall z (x < y \rightarrow (y < z \rightarrow x < z)),$$

$$\text{(UTS)} \quad \forall y (\overline{\Box \varphi(y)} \leftrightarrow \varphi(y)).$$

Then E is inconsistent.

I understand the disappointment: UTS is obviously inconsistent.

Bear with me...

The point is the proof.

THEOREM (YABLO'S PARADOX)

Assume that E contains all the following sentences:

$$\text{(SER)} \quad \forall x \exists y x < y,$$

$$\text{(TRANS)} \quad \forall x \forall y \forall z (x < y \rightarrow (y < z \rightarrow x < z)),$$

$$\text{(UTS)} \quad \forall y (\overline{\Box \varphi(y)} \leftrightarrow \varphi(y)).$$

Then E is inconsistent.

I understand the disappointment: UTS is obviously inconsistent.

Bear with me...

The point is the proof.

PROOF.

I apply the diagonal lemma with a free variable to the formula

$$\forall z > y \neg \Box \text{sub}(v, \bar{y}, q(z)),$$

which has exactly x and y as free variables. This yields a formula $\theta(y)$ with the following property:

$$E \vdash \forall y \left(\theta(y) \leftrightarrow \forall z > y \neg \Box \text{sub}(\overline{\theta(y)}, \bar{y}, q(z)) \right).$$

That is

$$(10) \quad E \vdash \forall y \left(\theta(y) \leftrightarrow \forall z > y \neg \Box \overline{\theta(z)} \right).$$

The sentences $\theta(\bar{e})$ for arbitrary expressions e correspond to the Yablo sentences. Priest (1997) defined the Yablo sentences in this way in arithmetic, after Visser (1989) had used a similar method to obtain his paradox. The contradiction can now be derived in the following way in E: □

$$\begin{array}{ll}
E \vdash \forall y (\overline{\Box\theta(y)} \leftrightarrow \theta(y)) & \text{UTS} \\
\leftrightarrow \forall z > y \neg \overline{\Box\theta(z)} & (10) \\
\rightarrow \exists z > y \overline{\Box\theta(z)} & \text{SER} \\
\rightarrow \exists z > y \forall w > z \neg \overline{\Box\theta(w)} & \text{second line and TRANS} \\
\rightarrow \exists z > y \theta(z) & (10) \\
\rightarrow \exists z > y \overline{\Box\theta(z)} & \text{UTS}
\end{array}$$

From the second and last lines we get $\forall y \neg \overline{\Box\theta(y)}$.

$$\begin{array}{ll}
E \vdash \forall y \neg \theta(y) & \text{UTS} \\
E \vdash \forall y \exists z > y \overline{\Box\theta(z)} & (10)
\end{array}$$

The last line contradicts the previously derived $\forall y \neg \overline{\Box\theta(y)}$.

UTS is inconsistent by itself because of the liar paradox, but the same proof strategy can be used to prove that the following typed schema is sufficient.

$$(vUTS) \quad \forall y (\Box_y \overline{\varphi(y)} \leftrightarrow \varphi(y)),$$

where all terms occurring in index position in $\varphi(y)$ are variables x_1, x_2, \dots, x_k distinct from y (but not complex terms or constants) and all quantifiers $\forall x_1, \forall x_2, \dots, \forall x_k$ in $\varphi(y)$ are restricted by $x_i > y$, respectively.

See the book for a proof. This schema is consistent by itself.

UTS is inconsistent by itself because of the liar paradox, but the same proof strategy can be used to prove that the following typed schema is sufficient.

$$(vUTS) \quad \forall y (\Box_y \overline{\varphi(y)} \leftrightarrow \varphi(y)),$$

where all terms occurring in index position in $\varphi(y)$ are variables x_1, x_2, \dots, x_k distinct from y (but not complex terms or constants) and all quantifiers $\forall x_1, \forall x_2, \dots, \forall x_k$ in $\varphi(y)$ are restricted by $x_i > y$, respectively.

See the book for a proof. This schema is consistent by itself.

I get proper inconsistencies, not just ω -inconsistencies, because I quantify over indicies.

There are other versions of these paradoxes. Some are mere ω -inconsistencies.

methodology

What do these results tell us?

- (i) We cannot have a predicate with the mentioned properties. We cannot combine predicates with certain mixed axioms.
- (ii) We cannot define a predicate with certain properties, but could add a typed predicate axiomatically.
- (iii) There is some complexity or definability hierarchies hidden.
- (iv) There are more limitative results than just plain inconsistencies: internal inconsistencies, trivialities (e.g. predicates cannot apply to anything), ω -inconsistencies, and 'unintended' consequences of various kinds.

What do these results tell us?

- (i) We cannot have a predicate with the mentioned properties. We cannot combine predicates with certain mixed axioms.
- (ii) We cannot define a predicate with certain properties, but could add a typed predicate axiomatically.
- (iii) There is some complexity or definability hierarchies hidden.
- (iv) There are more limitative results than just plain inconsistencies: internal inconsistencies, trivialities (e.g. predicates cannot apply to anything), ω -inconsistencies, and 'unintended' consequences of various kinds.

What do these results tell us?

- (i) We cannot have a predicate with the mentioned properties. We cannot combine predicates with certain mixed axioms.
- (ii) We cannot define a predicate with certain properties, but could add a typed predicate axiomatically.
- (iii) There is some complexity or definability hierarchies hidden.
- (iv) There are more limitative results than just plain inconsistencies: internal inconsistencies, trivialities (e.g. predicates cannot apply to anything), ω -inconsistencies, and 'unintended' consequences of various kinds.

What do these results tell us?

- (i) We cannot have a predicate with the mentioned properties. We cannot combine predicates with certain mixed axioms.
- (ii) We cannot define a predicate with certain properties, but could add a typed predicate axiomatically.
- (iii) There is some complexity or definability hierarchies hidden.
- (iv) There are more limitative results than just plain inconsistencies: internal inconsistencies, trivialities (e.g. predicates cannot apply to anything), ω -inconsistencies, and 'unintended' consequences of various kinds.

The paradoxes that are not just plain inconsistencies teach us a lesson about alleged solutions:

A mere consistency proof doesn't tell us that there is no paradox.

The proposed strategy may still exclude the standard model, make the notion trivial, or generate off consequences.

The paradoxes that are not just plain inconsistencies teach us a lesson about alleged solutions:

A mere consistency proof doesn't tell us that there is no paradox.

The proposed strategy may still exclude the standard model, make the notion trivial, or generate off consequences.

What can go wrong?

outright inconsistency

ω -inconsistency or no ω -model

triviality: The modality applies to all (or no) sentence.

unacceptable consequences: The theory implies a contingent truth or decides a question it should not decide.

and there thousand ways things can go wrong.

What can go wrong?

outright inconsistency

ω -inconsistency or no ω -model

triviality: The modality applies to all (or no) sentence.

unacceptable consequences: The theory implies a contingent truth or decides a question it should not decide.

and there thousand ways things can go wrong.

What can go wrong?

outright inconsistency

ω -inconsistency or no ω -model

triviality: The modality applies to all (or no) sentence.

unacceptable consequences: The theory implies a contingent truth
or decides a question it should not decide.

and there thousand ways things can go wrong.

What can go wrong?

outright inconsistency

ω -inconsistency or no ω -model

triviality: The modality applies to all (or no) sentence.

unacceptable consequences: The theory implies a contingent truth or decides a question it should not decide.

and there thousand ways things can go wrong.

What can go wrong?

outright inconsistency

ω -inconsistency or no ω -model

triviality: The modality applies to all (or no) sentence.

unacceptable consequences: The theory implies a contingent truth or decides a question it should not decide.

and there thousand ways things can go wrong.

McGee's ω -inconsistency

For the paradox I require a modicum of arithmetic.

We use strings $xxx\dots$ as natural numbers and call expressions $\overline{xxx\dots}$ numerals; they act as constants for numbers. Moreover, \underline{n} , the numeral of n , stands for

$$\underbrace{\overline{x\dots x}}_n$$

For instance, $\underline{4}$, that is, \overline{xxxx} , is the numeral for 4.

DEFINITION

$\text{Nat}(x)$ is defined as $\text{sub}(x, \bar{x}, \underline{0}) = \underline{0}$.

LEMMA

$E \vdash \text{Nat}(\underline{n})$ for all natural numbers n .

DEFINITION

$\text{Nat}(x)$ is defined as $\text{sub}(x, \bar{x}, \underline{0}) = \underline{0}$.

LEMMA

$E \vdash \text{Nat}(\underline{n})$ for all natural numbers n .

I write $\forall n \varphi(n)$ for $\forall x (\text{Nat}(x) \rightarrow \varphi(x))$, for any variable x , and similarly $\exists n \varphi(n)$ for $\exists x (\text{Nat}(x) \wedge \varphi(x))$. This abbreviation is used in conjunction with the underdotting convention as in $\forall k \overline{\varphi(\dot{k})}$, which is short for

$$\forall x (\text{Nat}(x) \rightarrow \text{sub}(qx, \bar{x}, \overline{\varphi(x)})),$$

where x is a variable with no bound occurrences in φ .

LEMMA

Assume \underline{n} , \underline{k} , $\underline{n+k}$, and $\underline{n \cdot k}$ are numerals for n , k , $n+k$, and $n \cdot k$, respectively. Then the following holds:

- (i) $E \vdash \underline{n} \wedge \underline{k} = \underline{n+k}$,
- (ii) $E \vdash \text{sub}(\underline{k}, \bar{x}, \underline{n}) = \underline{n \cdot k}$.

In particular we have $E \vdash \underline{n} \wedge \underline{1} = \underline{n+1}$.

We expect that every number except $\underline{0}$ has a unique predecessor.
However, this relies on the linearity of expressions, which is expressed by axiom A8.

LEMMA

$$E \vdash \forall n \forall k (n \hat{=} \underline{1} = k \hat{=} \underline{1} \rightarrow n = k).$$

DEFINITION

E is ω -inconsistent if and only if there is a formula $\varphi(x)$ with the following properties:

- (i) $E \vdash \varphi(\underline{k})$ for all natural numbers k ,
- (ii) $E \vdash \neg \forall n \varphi(n)$.

LEMMA

$E \vdash \forall x (\text{Nat}(x) \rightarrow \text{Nat}(x \hat{=} \underline{1}))$.

LEMMA

The theory E proves $\forall n \varphi(n) \rightarrow \forall n \varphi(n \hat{=} \underline{1})$ for all formulæ $\varphi(x)$ of \mathcal{L} .

The sentence (S) says that at least one of the following sentences is not true:

(S)

(S) is true.

'(S) is true' is true.

'“(S) is true” is true' is true.

⋮

McGee's ω -inconsistency theorem can be obtained by formalizing (S).

Here is the theorem from (McGee 1985):

THEOREM (MCGEE'S ω -INCONSISTENCY THEOREM)

Assume that E is closed under the rule in (i) and contains the formula (v) and the schemas (ii)–(iv) for all sentences φ and ψ and all formulæ $\chi(x)$ having at most x free.

- (i) NEC,
- (ii) $\Box\overline{\varphi \rightarrow \psi} \rightarrow (\Box\overline{\varphi} \rightarrow \Box\overline{\psi})$,
- (iii) $\Box\overline{\neg\varphi} \rightarrow \neg\Box\overline{\varphi}$,
- (iv) $\forall x \Box\overline{\chi(x)} \rightarrow \Box\overline{\forall x \chi(x)}$,
- (v) $\forall n (n = \underline{0} \vee \exists k n = k \hat{\ } \underline{1}) \wedge \neg \exists k \underline{0} = k \hat{\ } \underline{1}$.

Then E is ω -inconsistent.

I only sketch a proof.

bibliography

References

- Roy Cook. *The Yablo Paradox: An Essay on Circularity*. Oxford University Press, 2014.
- Haskell B. Curry. The inconsistency of certain formal logics. *Journal of Symbolic Logic*, 7:115–117, 1942.
- Kurt Gödel. Über formal unentscheidbare sätze der principia mathematica und verwandter systeme I. *Monatshefte für Mathematik*, 38:173–198, 1931.
- Volker Halbach and Leon Horsten. Self-referential gettier sentences. In Mattia Petrolo and Giorgio Venturi, editors, *Paradoxes Between Truth and Proof*, pages ??–?? 2025. to appear.
- Leon Horsten and Hannes Leitgeb. No future. *Journal of Philosophical Logic*, 30:259–265, 2001.

- Paul Horwich. *Truth*. Oxford University Press, second edition, 1998.
- Jeffrey Ketland. Deflationism and Gödel phenomena – reply to Tennant. *Mind*, 114:75–88, 2005.
- Saul Kripke. Outline of a theory of truth. *Journal of Philosophy*, 72: 690–716, 1975.
- Saul Kripke. Ungroundedness in Tarskian languages. *Journal of Philosophical Logic*, 48:603–609, 2019.
- Vann McGee. How truthlike can a predicate be? A negative result. *Journal of Philosophical Logic*, 14:399–410, 1985.
- Vann McGee. Maximal consistent sets of instances of Tarski's schema (T). *Journal of Philosophical Logic*, 21:235–241, 1992.
- Richard Montague. Syntactical treatments of modality, with corollaries on reflexion principles and finite axiomatizability. *Acta Philosophica Fennica*, 16:153–67, 1963. Reprinted in Montague 1974, 286–302.

- Richard Montague. *Formal Philosophy: Selected Papers of Richard Montague*. Yale University Press, New Haven and London, 1974.
Edited and with an introduction by Richmond H. Thomason.
- Graham Priest. Yablo's paradox. *Analysis*, 57:236–242, 1997.
- Richmond H. Thomason. A note on syntactical treatments of modality. *Synthese*, 44:391–396, 1980.
- Albert Visser. Semantics and the liar paradox. In Dov Gabbay and Franz Guenther, editors, *Handbook of Philosophical Logic*, volume 4, pages 617–706. Reidel, Dordrecht, 1989.
- Stephen Yablo. Truth and reflection. *Journal of Philosophical Logic*, 14: 297–349, 1985.
- Stephen Yablo. Paradox without self-reference. *Analysis*, 53:251–252, 1993.